

Table 19.12

Non-Hodgkin Lymphoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2016

All Races, Males, All Ages

Year of Diagnosis

| SEER Location | 1975- 1979 | 1980- 1981 | 1982- 1983 | 1984- 1985 | 1986- 1987 | 1988- 1989 | 1990- 1991 | 1992- 1993 | 1994- 1995 | 1996- 1997 | 1998- 1999 | 2000- 2001 | 2002- 2003 | 2004- 2005 | 2006- 2007 | 2008- 2009 | 2010- 2011 | 2012- 2013 | 2014- 2016 |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| SEER 9 ^b | 5,222 | 2,506 | 2,704 | 3,176 | 3,627 | 3,994 | 4,570 | 4,729 | 5,325 | 5,260 | 5,263 | 5,562 | 5,775 | 6,226 | 6,401 | 6,704 | 6,998 | 7,072 | 11,154 |
| SEER 9 except SF-Oakland | 4,285 | 2,088 | 2,267 | 2,603 | 2,819 | 3,091 | 3,507 | 3,694 | 4,159 | 4,249 | 4,364 | 4,515 | 4,818 | 5,209 | 5,398 | 5,628 | 5,841 | 6,015 | 9,363 |
| San Francisco-Oakland SMSA ^c | 937 | 418 | 437 | 573 | 808 | 903 | 1,063 | 1,035 | 1,166 | 1,011 | 899 | 1,047 | 957 | 1,017 | 1,003 | 1,076 | 1,157 | 1,057 | 1,791 |
| San Francisco City/County | 245 | 96 | 123 | 210 | 326 | 393 | 467 | 410 | 450 | 354 | 213 | 256 | 220 | 283 | 219 | 216 | 240 | 225 | 362 |

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2016

All Races, Males, All Ages

Year of Diagnosis

| SEER Location | 1975- 1979 | 1980- 1981 | 1982- 1983 | 1984- 1985 | 1986- 1987 | 1988- 1989 | 1990- 1991 | 1992- 1993 | 1994- 1995 | 1996- 1997 | 1998- 1999 | 2000- 2001 | 2002- 2003 | 2004- 2005 | 2006- 2007 | 2008- 2009 | 2010- 2011 | 2012- 2013 | 2014- 2016 |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| SEER 9 ^b | 13.4 | 15.5 | 16.1 | 18.3 | 20.1 | 21.5 | 23.4 | 23.3 | 25.2 | 24.4 | 23.8 | 24.4 | 24.6 | 25.8 | 25.4 | 25.5 | 25.5 | 24.5 | 24.4 |
| SEER 9 except SF-Oakland | 13.2 | 15.4 | 16.1 | 18.1 | 18.8 | 20.1 | 21.7 | 21.8 | 23.6 | 23.6 | 23.5 | 23.6 | 24.4 | 25.6 | 25.4 | 25.3 | 25.2 | 24.7 | 24.3 |
| San Francisco-Oakland SMSA ^c | 14.7 | 15.8 | 16.0 | 19.4 | 26.7 | 28.5 | 31.7 | 30.5 | 33.5 | 28.8 | 25.2 | 28.6 | 25.6 | 26.6 | 25.6 | 26.3 | 26.8 | 23.5 | 24.6 |
| San Francisco City/County | 14.3 | 14.4 | 17.6 | 29.3 | 44.4 | 51.9 | 61.0 | 52.3 | 57.4 | 44.8 | 27.4 | 32.5 | 26.9 | 35.4 | 26.5 | 25.7 | 27.3 | 25.2 | 24.6 |

^a Rates are per 100,000 males and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Non-Hodgkin Lymphoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.