



Table 19.23  
Non-Hodgkin Lymphoma

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2012-2016

Males

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	7.25	0.03		
High Five States				
Kentucky	8.86	0.30	01(01,12)	22.13 <sup>c</sup>
Iowa	8.45	0.32	02(01,20)	16.47 <sup>c</sup>
Michigan	8.42	0.19	03(01,14)	16.15 <sup>c</sup>
Indiana	8.39	0.24	04(01,16)	15.67 <sup>c</sup>
Delaware	8.24	0.58	05(01,44)	13.64
Low Five States				
Arizona	6.35	0.19	47(32,51)	-12.44
Alaska	6.30	0.75	48(02,51)	-13.11
Hawaii	6.20	0.41	49(19,51)	-14.52
D.C.	6.14	0.71	50(05,51)	-15.35 <sup>c</sup>
New Mexico	5.85	0.33	51(34,51)	-19.28 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	6.88	0.25	36(16,49)	-5.13	Montana	7.09	0.50	26(02,51)	-2.18
Alaska	6.30	0.75	48(02,51)	-13.11	Nebraska	7.37	0.40	20(02,48)	1.65
Arizona	6.35	0.19	47(32,51)	-12.44	Nevada	6.53	0.32	44(19,51)	-9.94
Arkansas	7.21	0.31	22(07,47)	-0.61	New Hampshire	7.10	0.46	25(04,51)	-2.10
California	6.77	0.09	41(28,46)	-6.66	New Jersey	7.21	0.18	21(14,41)	-0.59
Colorado	6.43	0.24	46(24,51)	-11.33	New Mexico	5.85	0.33	51(34,51)	-19.28 <sup>c</sup>
Connecticut	7.05	0.28	28(12,48)	-2.73	New York	7.02	0.12	30(19,43)	-3.23
Delaware	8.24	0.58	05(01,44)	13.64	North Carolina	7.09	0.18	27(16,44)	-2.21
D.C.	6.14	0.71	50(05,51)	-15.35 <sup>c</sup>	North Dakota	6.82	0.61	39(02,51)	-5.98
Florida	6.83	0.11	37(24,46)	-5.85	Ohio	8.00	0.17	09(03,20)	10.29
Georgia	6.97	0.19	34(17,46)	-3.87	Oklahoma	8.06	0.30	08(01,26)	11.09
Hawaii	6.20	0.41	49(19,51)	-14.52	Oregon	7.94	0.28	12(01,28)	9.47
Idaho	7.75	0.45	16(01,46)	6.92	Pennsylvania	7.83	0.15	15(05,23)	7.94
Illinois	7.39	0.16	19(12,35)	1.90	Rhode Island	6.49	0.50	45(10,51)	-10.50
Indiana	8.39	0.24	04(01,16)	15.67 <sup>c</sup>	South Carolina	6.81	0.25	40(18,50)	-6.04
Iowa	8.45	0.32	02(01,20)	16.47 <sup>c</sup>	South Dakota	7.01	0.56	31(02,51)	-3.32
Kansas	7.13	0.32	24(09,48)	-1.74	Tennessee	8.20	0.23	06(01,20)	13.11
Kentucky	8.86	0.30	01(01,12)	22.13 <sup>c</sup>	Texas	6.98	0.12	32(20,43)	-3.68
Louisiana	8.14	0.29	07(01,22)	12.22	Utah	6.70	0.36	42(13,51)	-7.54
Maine	7.54	0.45	18(01,48)	3.95	Vermont	7.94	0.69	11(01,49)	9.50
Maryland	6.82	0.23	38(18,49)	-5.88	Virginia	6.94	0.20	35(18,47)	-4.29
Massachusetts	6.59	0.20	43(24,50)	-9.10	Washington	7.90	0.22	13(02,24)	8.97
Michigan	8.42	0.19	03(01,14)	16.15 <sup>c</sup>	West Virginia	7.74	0.38	17(01,42)	6.75
Minnesota	7.97	0.25	10(02,24)	9.95	Wisconsin	7.86	0.23	14(02,27)	8.44
Mississippi	7.17	0.34	23(07,48)	-1.10	Wyoming	6.98	0.71	33(01,51)	-3.77
Missouri	7.02	0.22	29(16,46)	-3.19					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.