Table 20.11 Cancer of the Oral Cavity and Pharynx by Subsite (Invasive)

SEER^a Relative Survival (Percent) by Cancer Site, 2009-2015

		Case	Survival Time					
	Number	Distribution	1-Year	2-Year	3-Year	4-Year	5-Year Percent	
Site	of Cases	(%)	Percent	Percent	Percent	Percent		
Oral Cavity & Pharynx	57854	100	84.9	75.9	71.1	67.8	65.3	
Lip	3393	6	98.2	95.4	93.7	91.8	89.7	
Tongue	17556	30	85.4	76.0	71.4	68.4	66.4	
Gum & Other Mouth	6868	12	80.7	70.7	65.5	61.8	59.0	
Floor of Mouth	2490	4	79.9	66.9	61.2	56.3	51.7	
Oropharynx & Tonsil	13929	24	87.4	79.9	75.3	72.1	70.0	
Hypopharynx	2797	5	67.1	50.1	42.1	37.1	34.6	
Salivary Gland	6169	11	88.6	81.5	76.5	73.7	71.4	
Nasopharynx	3523	6	84.2	74.7	68.8	65.0	61.3	
Other Oral Cavity & Pharynx	1129	2	68.8	59.2	53.5	49.6	46.6	

5-Year Relative Survival (Percent), 2009-2015

	Stage					All Stages			
					Unstaged/	White	White	Black	Black
Site	All Stages	Localized	Regional	Distant	Unknown	Male	Female	Male	Female
Oral Cavity & Pharynx	65.3	84.4	66.0	39.1	50.9	66.4	67.7	46.7	52.0
Lip	89.7	92.4	60.4 ^b	27.5 ^b	86.9	89.9	88.7	66.6°	-
Tongue	66.4	81.3	67.8	39.0	50.0	68.6	66.5	45.3	42.0
Gum & Other Mouth ^d	59.0	-	-	-	=	55.5	63.0	48.9	57.1
Floor of Mouth	51.7	76.9	38.2	19.8	40.0 ^b	50.8	57.7	39.0	38.2 ^b
Oropharynx & Tonsil ^d	70.0	-	-	-	-	72.6	68.0	51.7	45.5
Hypopharynx ^d	34.6	-	-	-	-	37.3	39.2	20.2	17.3 ^b
Salivary Gland	71.4	93.6	65.2	35.4	48.6	61.5	80.1	71.3	81.0
Nasopharynx ^d	61.3	-	-	-	-	57.0	61.5	53.7	49.9 ^b
Other Oral Cavity & Pharyn \mathbf{x}^{d}	46.6	-	-	-	-	52.9	37.9	29.4 ^b	32.6 ^b

SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Expected survival rates are derived from life tables by socio-economic status, geography and race developed by the SEER program. Based on follow-up of patients into 2016.

The standard error is between 5 and 10 percentage points.

The standard error is greater than 10 percentage points.

Survival by stage was not calculated for this site due to inconsistent coding of stage over time.

Statistic could not be calculated due to fewer than 25 cases during the time period.