

Table 23.7
Cancer of the Prostate (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2012-2016</u>						
Age at Diagnosis/Death:						
All ages	109.5	101.9	176.7	19.2	18.0	38.9
Under 65	44.3	39.9	82.7	1.7	1.4	4.0
65 and over	560.4	530.7	826.0	140.7	132.4	280.1
All ages (WHO world std) ^c	83.5	77.3	138.0	8.5	7.8	18.0
<u>Age-Specific Rates, 2012-2016</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	0.6	0.5	1.6	0.0	-	-
40-44	7.8	6.5	19.0	0.2	0.1	0.4
45-49	34.0	28.0	83.0	0.8	0.7	1.7
50-54	116.0	102.5	229.0	3.2	2.7	7.3
55-59	248.6	224.0	459.2	8.6	7.3	20.5
60-64	417.6	384.7	720.8	19.8	17.1	47.6
65-69	627.8	584.6	1,017.3	38.1	33.8	91.4
70-74	652.0	618.5	961.8	70.0	63.4	160.6
75-79	567.4	537.4	798.3	124.4	114.7	267.2
80-84	425.3	408.5	547.1	217.2	205.1	433.5
85+	367.0	360.5	494.5	452.9	438.9	787.6

^a SEER 21 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.