

Table 26.13
Cancer of the Thyroid (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years,
Lifetime Risk of Being Diagnosed with Cancer Given Alive and Cancer-Free at Current Age, and
Lifetime Risk of Dying from Cancer Given Alive at Current Age
Males, 2014-2016 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer
		+10 yrs	+20 yrs	+30 yrs	Ever				+10 yrs	+20 yrs	+30 yrs	Ever	
All Races	0	0.00	0.01	0.04	0.70	0.06	Asian/ Pacific Islander	0	0.00	0.01	0.04	0.73	0.08
	10	0.01	0.04	0.12	0.70	0.06		10	0.01	0.04	0.12	0.73	0.08
	20	0.03	0.11	0.22	0.70	0.06		20	0.03	0.11	0.23	0.72	0.08
	30	0.07	0.18	0.32	0.67	0.06		30	0.08	0.20	0.33	0.69	0.08
	40	0.11	0.25	0.41	0.61	0.06		40	0.12	0.26	0.41	0.62	0.08
	50	0.14	0.31	0.44	0.51	0.06		50	0.14	0.30	0.42	0.51	0.08
	60	0.18	0.32	0.38	0.39	0.06		60	0.17	0.29	0.37	0.38	0.08
	70	0.16	0.24	-	0.25	0.05		70	0.14	0.22	-	0.24	0.07
White	80	0.10	-	-	0.11	0.04	American Indian/ Alaska Native ^a	80	0.11	-	-	0.12	0.06
	0	0.00	0.01	0.05	0.74	0.06		0	0.00	0.01	0.03	0.37	0.04
	10	0.01	0.05	0.13	0.74	0.06		10	0.01	0.03	0.06	0.37	0.04
	20	0.04	0.12	0.23	0.74	0.06		20	0.02	0.05	0.09	0.36	0.04
	30	0.08	0.19	0.34	0.71	0.06		30	0.03	0.07	0.13	0.35	0.04
	40	0.12	0.27	0.44	0.64	0.06		40	0.04	0.10	0.19	0.33	0.04
	50	0.15	0.33	0.47	0.54	0.06		50	0.06	0.16	0.24	0.30	0.05
	60	0.19	0.34	0.40	0.41	0.06		60	0.10	0.20	0.26	0.26	0.05
Black	70	0.17	0.25	-	0.26	0.06	Hispanic ^b	70	0.11	0.18	-	0.19	0.03
	80	0.10	-	-	0.12	0.04		80	0.11	-	-	0.11	0.03
	0	0.00	0.01	0.01	0.33	0.04		0	0.00	0.01	0.03	0.55	0.08
	10	0.01	0.01	0.04	0.33	0.04		10	0.01	0.03	0.08	0.56	0.08
	20	0.01	0.04	0.09	0.33	0.04		20	0.02	0.07	0.14	0.55	0.08
	30	0.03	0.08	0.15	0.33	0.04		30	0.05	0.13	0.22	0.54	0.08
	40	0.05	0.13	0.21	0.31	0.04		40	0.08	0.18	0.30	0.49	0.08
	50	0.08	0.17	0.23	0.27	0.04		50	0.10	0.23	0.35	0.43	0.08
	60	0.10	0.17	0.20	0.21	0.04		60	0.13	0.27	0.33	0.34	0.08
	70	0.10	0.14	-	0.14	0.03		70	0.15	0.22	-	0.24	0.07
	80	0.06	-	-	0.07	0.02		80	0.09	-	-	0.11	0.06

Devcan 6.7.7, April 2019, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts). Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the PRCDA (Purchased/Referred Care Delivery Areas) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

- Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

- Statistic could not be calculated.

A percent of 0.00 represents a value that is below 0.005.