


Table 26.18
Cancer of the Thyroid (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2012-2016


Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	0.51	0.01		
High Five States				
D.C.	0.71	0.15	01(01,49)	39.11 ^c
Utah	0.65	0.08	02(01,39)	27.80 ^c
California	0.65	0.02	03(01,10)	26.78 ^c
Idaho	0.64	0.09	04(01,45)	26.29 ^c
Hawaii	0.60	0.08	05(01,47)	17.86 ^c
Low Five States				
West Virginia	0.42	0.06	45(06,49)	-17.86 ^c
Maine	0.42	0.07	46(04,49)	-18.22 ^c
Tennessee	0.40	0.03	47(22,49)	-20.78 ^c
Delaware	0.37	0.08	48(04,49)	-27.33 ^c
New Hampshire	0.33	0.07	49(12,49)	-34.25 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	0.43	0.04	42(11,49)	-15.25 ^c	Montana	0.44	0.08	40(01,49)	-12.71
Alaska	-	-	-(-,-)	-	Nebraska	0.52	0.07	17(01,49)	2.20
Arizona	0.46	0.03	37(10,48)	-8.76	Nevada	0.56	0.06	10(01,46)	10.48
Arkansas	0.48	0.05	32(04,49)	-5.48	New Hampshire	0.33	0.07	49(12,49)	-34.25 ^c
California	0.65	0.02	03(01,10)	26.78 ^c	New Jersey	0.49	0.03	27(08,46)	-4.64
Colorado	0.53	0.05	14(02,45)	3.78	New Mexico	0.59	0.07	07(01,46)	16.10 ^c
Connecticut	0.43	0.04	43(09,49)	-15.68 ^c	New York	0.53	0.02	15(06,35)	3.37
Delaware	0.37	0.08	48(04,49)	-27.33 ^c	North Carolina	0.52	0.03	18(05,42)	1.57
D.C.	0.71	0.15	01(01,49)	39.11 ^c	North Dakota	0.58	0.12	08(01,49)	13.75
Florida	0.44	0.02	41(23,47)	-13.16	Ohio	0.47	0.03	35(12,46)	-7.88
Georgia	0.42	0.03	44(18,49)	-16.73 ^c	Oklahoma	0.46	0.05	38(07,49)	-10.26
Hawaii	0.60	0.08	05(01,47)	17.86 ^c	Oregon	0.59	0.05	06(01,39)	16.53 ^c
Idaho	0.64	0.09	04(01,45)	26.29 ^c	Pennsylvania	0.50	0.02	21(08,42)	-1.75
Illinois	0.48	0.03	28(09,45)	-4.70	Rhode Island	0.57	0.10	09(01,49)	11.28
Indiana	0.48	0.04	34(07,47)	-6.26	South Carolina	0.51	0.04	19(03,46)	1.10
Iowa	0.51	0.05	20(02,48)	0.48	South Dakota	0.48	0.10	31(01,49)	-5.13
Kansas	0.48	0.05	33(04,49)	-5.56	Tennessee	0.40	0.03	47(22,49)	-20.78 ^c
Kentucky	0.50	0.04	23(04,47)	-2.49	Texas	0.54	0.02	12(05,32)	5.34
Louisiana	0.46	0.04	36(06,49)	-8.61	Utah	0.65	0.08	02(01,39)	27.80 ^c
Maine	0.42	0.07	46(04,49)	-18.22 ^c	Vermont	0.48	0.11	29(01,49)	-4.77
Maryland	0.50	0.04	24(05,47)	-2.50	Virginia	0.46	0.03	39(11,48)	-10.38
Massachusetts	0.50	0.04	22(06,46)	-2.33	Washington	0.56	0.04	11(02,38)	10.07
Michigan	0.49	0.03	25(07,45)	-2.76	West Virginia	0.42	0.06	45(06,49)	-17.86 ^c
Minnesota	0.52	0.04	16(03,45)	2.41	Wisconsin	0.53	0.04	13(03,43)	4.76
Mississippi	0.49	0.06	26(02,48)	-2.77	Wyoming	-	-	-(-,-)	-
Missouri	0.48	0.04	30(06,47)	-4.77					



Age-Adjusted Death
Rates per 100,000
Quantile Interval



- ^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- ^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).
- ^c Absolute percent difference between state rate and total U.S. rate is 15% or more.
- ^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.
- SE Standard error of the rate.
- PD Percent difference between state rate and total U.S. rate.
- Statistic not shown. Rate based on less than 16 cases for the time interval.