## Table 29.2

## Age-Adjusted SEER Cancer Incidence Rates<sup>a</sup>

By International Classification of Childhood Cancer(ICCC)<sup>b</sup> Selected Group and Subgroup<sup>c</sup> and Year of Diagnosis

Excluding benign brain and myelodysplastic syndromes

All Races, Males and Females, Ages 0-19

ICCC Group and Subgroup	<u>1975-2016</u>	<u>1975-1986</u>	1987-1996	<u>1997-2006</u>	2007-2016
All ICCC Groups Combined	162.5	146.3	159.2	165.1	180.2
I Leukemia I(a) Lymphoid leukemia	39.8 29.7	35.4 26.1	38.2 28.7	41.6 31.3	43.8 32.8
I(b) Acute myeloid leukemia	7.2	6.1	6.9	8.1	7.9
II Lymphomas and reticuloendothelial neoplasms II(a) Hodqkin lymphoma	25.0 13.1	24.6 14.3	24.4 13.4	23.5 11.4	27.2 13.0
II(b,c,e) Non-Hodgkin lymphoma	11.4	9.6	10.3	11.6	13.9
III CNS & misc intracranial & intraspinal neoplasms III(a) Ependymomas and choroid plexus tumor	28.6 2.5	24.1 1.9	29.9 2.6	29.4 2.5	31.3 2.9
III(b) Astrocytoma	14.5	12.1	15.7	14.5	16.1
III(c) Intracranial and intraspinal embryonal tumors III(d) Other gliomas	5.6 5.2	4.5 4.6	5.7 5.1	6.2 5.6	6.0 5.6
IV Neuroblastoma and other peripheral nervous cell tumors IV(a) Neuroblastoma and ganglioneuroblastoma	8.6 8.3	8.2 8.0	8.5 8.2	8.7 8.5	8.9 8.6
V Retinoblastoma	3.1	2.8	3.2	3.2	3.0
VI Renal tumors	6.7	6.5	6.8	6.6	6.9
VI(a) Nephroblastoma and other nonepithelial renal tumor	6.2	6.2	6.4	6.0	6.2
VII Hepatic tumors	2.0	1.4	1.6	2.3	2.7
VII(a) Hepatoblastoma	1.4	0.9	1.2	1.7	1.9
VIII Malignant bone tumors VIII(a) Osteosarcoma	8.7 4.8	8.4 4.3	8.8 4.9	8.5 4.7	9.3 5.3
VIII(c) Ewing tumor and related sarcomas of bone	3.0	3.2	2.8	2.8	3.1
IX Soft tissue and other extraosseous sarcomas	11.7	10.7	11.0	12.5	12.6
IX(a) Rhabdomyosarcoma	4.5	4.2	4.8	4.6	4.6
X Germ cell & trophoblastic tumors & neoplasms of gonads X(a) Intracranial and intraspinal germ-cell tumor	10.7 1.5	9.4 1.0	10.8 1.6	11.0 1.8	11.6 1.8
X(c) Malignant gonadal germ cell tumor	6.8	6.0	6.7	7.0	7.5
XI Other malignant epithelial neoplasms and melanomas <sup>d</sup>	17.1	14.0	15.2	17.1	22.3
XI(b) Thyroid carcinoma XI(d) Malignant melanoma	6.6 5.1	5.0 4.2	5.2 5.3	6.1 5.9	10.2 5.1

а SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Rates are per 1,000,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). International Classification of Childhood Cancer is based on ICD-O-3. Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. b International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005: Vol 103, No. 7 , pg 1457-1467. С Classifications are shown for invasive cases only except as noted. For ICCC groups and subgroups that include in situ

behavior, only invasive cases are shown except as noted.

d Includes in situ urinary bladder tumors.

Statistic could not be calculated. Rate based on less than 16 cases for the time interval.

\_