

Table 2.9
All Cancer Sites (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis
All Races, Males and Females

Survival Time	Year of Diagnosis																									
	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	69.1	70.6	73.2	76.3	76.6	77.0	77.6	77.9	78.8	79.0	79.5	79.7	79.6	80.0	80.3	80.9	81.5	81.6	81.9	81.8	82.2	81.8	82.1	81.9	82.7	82.8
2-year	59.8	61.3	64.4	68.8	69.4	70.1	70.9	71.5	72.3	72.7	73.4	74.0	73.5	74.0	74.3	75.4	75.8	75.9	76.1	76.3	76.5	75.9	76.3	76.3	76.3	77.0
3-year	54.8	56.2	59.8	64.9	65.8	66.7	67.5	68.1	69.0	69.5	70.3	70.9	70.4	71.0	71.3	72.3	72.7	72.7	73.1	73.2	73.4	72.6	73.0	73.1		
4-year	51.4	52.8	56.8	62.4	63.4	64.3	65.1	65.8	66.9	67.5	68.3	68.8	68.3	68.9	69.3	70.2	70.8	70.7	71.1	71.0	71.3	70.6	70.9			
5-year	48.9	50.2	54.4	60.4	61.6	62.5	63.3	64.0	65.2	66.0	66.8	67.3	66.8	67.3	67.7	68.7	69.4	69.1	69.6	69.5	69.7	69.0				
6-year	47.0	48.2	52.6	58.9	60.2	61.2	61.9	62.7	63.9	64.8	65.7	66.2	65.6	66.1	66.5	67.7	68.2	68.0	68.4	68.5	68.5					
7-year	45.4	46.5	51.1	57.6	59.1	60.1	61.0	61.7	62.9	63.7	64.7	65.2	64.5	65.1	65.6	66.7	67.2	66.9	67.5	67.5						
8-year	44.1	45.1	49.8	56.5	58.1	59.2	60.1	60.9	62.0	62.7	63.8	64.4	63.7	64.3	64.8	65.9	66.4	66.1	66.7							
9-year	42.9	43.9	48.6	55.5	57.2	58.2	59.2	59.9	61.2	62.0	63.1	63.6	62.9	63.5	64.0	65.2	65.8	65.3								
10-year	41.8	42.9	47.7	54.6	56.5	57.4	58.5	59.1	60.5	61.3	62.3	63.1	62.0	62.8	63.3	64.5	65.1									
11-year	40.9	41.9	46.8	53.7	55.6	56.8	57.8	58.4	59.8	60.5	61.7	62.4	61.4	62.1	62.7	63.9										
12-year	40.1	41.1	45.9	53.0	54.9	56.2	57.2	57.8	59.2	59.9	61.1	61.7	60.8	61.6	62.1											
13-year	39.3	40.3	45.1	52.3	54.0	55.5	56.5	57.1	58.6	59.2	60.4	61.2	60.3	61.0												
14-year	38.7	39.6	44.5	51.7	53.3	54.8	55.9	56.6	57.9	58.6	59.9	60.7	59.6													
15-year	38.0	39.0	43.8	51.1	52.7	54.3	55.4	56.1	57.4	58.0	59.4	60.3														
16-year	37.5	38.3	43.2	50.5	52.2	53.8	54.7	55.5	56.8	57.6	58.9															
17-year	36.9	37.7	42.7	49.8	51.6	53.2	54.2	55.0	56.3	57.1																
18-year	36.4	37.2	42.2	49.3	51.0	52.7	53.6	54.8	55.7																	
19-year	36.0	36.6	41.8	48.7	50.5	52.0	53.2	54.3																		
20-year	35.5	36.1	41.3	48.2	49.8	51.4	52.8																			
21-year	35.0	35.6	40.9	47.7	49.2	50.9																				
22-year	34.6	35.2	40.5	47.1	48.7																					
23-year	34.2	34.7	40.0	46.7																						
24-year	33.7	34.3	39.5																							
25-year	33.2	33.9	39.1																							
26-year	32.8	33.5	38.6																							
27-year	32.5	33.1	38.2																							
28-year	32.2	32.8	37.7																							
29-year	31.8	32.5																								
30-year	31.5	32.1																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 2.10
All Cancer Sites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	63.7	66.6	69.8	75.4	75.4	76.2	76.8	76.7	78.1	78.6	79.1	79.1	79.1	79.7	79.9	80.8	81.5	81.3	81.6	81.3	81.5	80.5	80.7	80.5	81.3	81.5
2-year	53.4	56.4	60.4	67.7	68.1	69.4	70.2	70.4	71.6	72.5	73.0	73.5	73.1	73.9	74.1	75.5	76.0	75.6	75.6	75.9	75.8	74.4	74.6	74.5	75.2	
3-year	48.2	51.2	55.5	64.1	64.7	66.1	67.1	67.2	68.5	69.5	70.4	70.8	70.2	71.0	71.3	72.6	73.3	72.7	73.0	73.0	72.7	71.2	71.2	71.3		
4-year	44.8	47.7	52.3	61.7	62.5	63.9	64.9	65.0	66.6	67.7	68.5	69.1	68.4	69.3	69.6	70.8	71.5	70.9	71.2	71.1	70.8	69.3	69.2			
5-year	42.2	45.0	49.9	59.8	60.9	62.3	63.2	63.5	65.2	66.3	67.4	67.9	67.2	68.0	68.2	69.5	70.3	69.6	69.8	69.7	69.5	67.8				
6-year	40.2	43.0	48.1	58.4	59.6	61.1	61.9	62.3	64.1	65.2	66.4	67.0	66.1	66.9	67.0	68.6	69.2	68.5	68.9	68.8	68.5					
7-year	38.5	41.3	46.4	57.1	58.6	60.1	61.1	61.3	63.2	64.3	65.6	66.3	65.2	66.0	66.1	67.8	68.3	67.7	68.2	68.0						
8-year	37.1	39.9	45.1	55.9	57.6	59.3	60.3	60.5	62.4	63.3	64.9	65.5	64.5	65.3	65.3	67.0	67.7	67.0	67.6							
9-year	35.8	38.6	43.8	54.9	56.9	58.4	59.4	59.6	61.6	62.7	64.4	64.9	63.6	64.5	64.7	66.3	67.1	66.4								
10-year	34.6	37.6	42.8	54.0	56.0	57.6	58.8	59.0	61.0	62.0	63.6	64.4	62.8	63.8	64.1	65.7	66.5									
11-year	33.5	36.6	41.9	53.1	55.3	56.8	58.2	58.3	60.5	61.1	63.0	63.6	62.2	63.2	63.6	65.2										
12-year	32.6	35.7	40.9	52.4	54.4	56.2	57.6	57.5	59.8	60.6	62.6	63.2	61.6	62.7	63.1											
13-year	31.8	34.9	40.1	51.7	53.6	55.7	56.8	56.9	59.3	59.8	61.9	62.6	61.2	62.2												
14-year	31.1	34.1	39.3	51.0	52.7	55.1	56.2	56.5	58.6	59.2	61.5	62.2	60.6													
15-year	30.4	33.4	38.6	50.3	52.0	54.5	55.7	56.1	58.0	58.6	61.0	61.8														
16-year	29.7	32.7	37.9	49.7	51.4	54.0	55.0	55.6	57.6	58.3	60.5															
17-year	29.0	32.0	37.4	49.0	50.8	53.4	54.3	55.1	57.2	57.9																
18-year	28.5	31.5	36.8	48.5	50.4	53.1	53.6	54.9	56.6																	
19-year	28.0	30.9	36.3	47.9	49.8	52.5	53.3	54.6																		
20-year	27.6	30.4	35.9	47.5	48.8	51.9	53.0																			
21-year	27.1	29.8	35.5	46.8	48.2	51.6																				
22-year	26.6	29.3	35.1	46.2	47.6																					
23-year	26.1	28.9	34.5	45.7																						
24-year	25.6	28.5	34.1																							
25-year	25.1	28.2	33.6																							
26-year	24.6	27.7	33.2																							
27-year	24.3	27.4	32.9																							
28-year	24.0	27.1	32.4																							
29-year	23.6	26.8																								
30-year	23.2	26.5																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 2.11
All Cancer Sites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

Survival Time	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	74.6	74.8	76.6	77.4	78.0	77.8	78.5	79.2	79.5	79.4	79.9	80.3	80.1	80.4	80.7	80.9	81.4	81.9	82.2	82.3	82.9	83.2	83.5	83.4	84.1	84.1
2-year	66.1	66.2	68.7	70.1	70.8	71.0	71.6	72.6	73.1	73.0	73.8	74.4	74.0	74.2	74.6	75.2	75.7	76.1	76.5	76.7	77.3	77.3	78.0	78.1	78.8	
3-year	61.3	61.2	64.2	65.9	67.0	67.3	67.9	69.1	69.5	69.6	70.2	70.9	70.5	70.9	71.3	72.0	72.1	72.7	73.2	73.3	74.0	74.0	74.8	74.8		
4-year	58.0	57.9	61.4	63.2	64.3	64.8	65.4	66.6	67.1	67.3	68.0	68.5	68.2	68.4	69.1	69.7	70.0	70.6	71.0	71.0	71.8	71.9	72.6			
5-year	55.6	55.3	59.0	61.2	62.4	62.8	63.5	64.6	65.3	65.7	66.2	66.6	66.3	66.6	67.3	67.9	68.3	68.6	69.3	69.3	70.0	70.1				
6-year	53.6	53.4	57.3	59.6	61.0	61.3	62.0	63.2	63.8	64.4	64.9	65.3	65.0	65.3	66.0	66.7	67.0	67.3	68.0	68.1	68.6					
7-year	52.1	51.7	55.8	58.2	59.6	60.2	60.9	62.1	62.6	63.2	63.7	64.1	63.8	64.2	65.0	65.6	66.0	66.1	66.8	67.0						
8-year	50.8	50.4	54.6	57.1	58.6	59.1	60.0	61.2	61.6	62.2	62.7	63.2	62.9	63.4	64.2	64.7	65.1	65.1	65.8							
9-year	49.7	49.2	53.5	56.1	57.7	58.1	58.9	60.2	60.8	61.3	61.7	62.3	62.1	62.5	63.3	63.9	64.2	64.2								
10-year	48.7	48.1	52.5	55.3	57.0	57.3	58.1	59.3	60.1	60.6	61.1	61.7	61.2	61.7	62.5	63.2	63.4									
11-year	47.8	47.1	51.7	54.4	56.1	56.7	57.4	58.6	59.2	59.8	60.4	61.1	60.6	61.0	61.8	62.6										
12-year	47.1	46.4	50.9	53.7	55.4	56.1	56.9	58.1	58.5	59.2	59.5	60.3	60.0	60.5	61.1											
13-year	46.4	45.6	50.1	53.0	54.5	55.4	56.3	57.3	57.9	58.5	58.8	59.7	59.4	59.7												
14-year	45.8	44.9	49.5	52.4	54.0	54.7	55.7	56.8	57.2	58.0	58.2	59.2	58.7													
15-year	45.1	44.3	48.9	51.9	53.5	54.1	55.1	56.1	56.7	57.4	57.8	58.7														
16-year	44.6	43.6	48.3	51.3	52.9	53.6	54.4	55.6	56.0	56.9	57.3															
17-year	44.1	43.0	47.8	50.7	52.4	52.9	54.1	55.1	55.5	56.4																
18-year	43.6	42.5	47.3	50.2	51.6	52.4	53.6	54.8	54.9																	
19-year	43.2	42.0	46.9	49.6	51.2	51.7	53.1	54.2																		
20-year	42.7	41.6	46.5	49.1	50.7	51.1	52.5																			
21-year	42.2	41.0	46.0	48.6	50.2	50.3																				
22-year	41.8	40.6	45.6	48.0	49.6																					
23-year	41.4	40.2	45.1	47.6																						
24-year	40.9	39.7	44.7																							
25-year	40.4	39.2	44.3																							
26-year	40.1	38.9	43.7																							
27-year	39.7	38.4	43.2																							
28-year	39.3	38.0	42.7																							
29-year	39.0	37.7																								
30-year	38.6	37.2																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 3.9
Cancer of the Brain and Other Nervous System (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	46.3	47.4	50.7	52.6	52.6	53.2	51.2	53.8	55.3	55.6	57.1	59.2	57.2	56.9	58.5	61.5	62.1	62.4	62.4	62.0	61.5	63.1	64.5	63.5	62.6	59.0
2-year	32.3	33.9	36.9	38.9	39.2	40.7	38.0	40.8	40.4	41.8	42.4	46.6	43.6	43.0	45.8	47.3	47.4	46.7	47.0	46.8	45.4	46.4	48.5	45.8	45.5	
3-year	27.9	29.4	32.7	34.9	36.0	36.6	34.1	35.7	36.1	38.4	39.0	40.6	38.1	38.2	40.0	40.6	41.0	41.0	41.4	39.9	39.8	38.7	40.2	38.2		
4-year	25.2	26.6	30.0	32.5	34.3	34.6	31.9	33.0	33.8	36.1	36.7	38.2	36.0	36.0	37.2	38.2	37.5	37.6	38.2	37.0	36.2	35.5	37.5			
5-year	23.1	24.6	28.0	30.8	33.3	32.5	30.3	31.1	32.4	34.9	34.8	36.1	34.2	35.0	34.9	35.9	35.5	36.1	36.3	34.3	33.9	33.4				
6-year	21.5	23.1	26.6	29.6	32.3	31.2	28.2	29.8	30.8	33.4	33.7	34.5	32.9	33.4	32.9	34.2	33.5	34.5	34.7	32.5	32.7					
7-year	20.5	21.9	25.5	28.3	31.3	30.3	27.5	28.8	30.1	31.7	32.6	33.6	31.4	32.0	31.9	33.4	32.5	33.1	33.6	31.3						
8-year	19.4	20.9	24.2	27.4	30.4	28.9	26.8	27.9	29.4	31.0	31.7	32.7	30.4	31.0	30.6	31.8	31.7	32.1	33.0							
9-year	18.4	19.9	23.4	26.6	29.9	27.7	25.6	27.2	28.3	30.1	31.1	31.7	29.6	30.2	29.8	30.7	30.7	31.0								
10-year	17.8	18.9	22.6	25.9	28.9	27.1	24.6	26.4	27.8	29.4	30.6	30.9	29.2	29.3	29.1	30.1	29.9									
11-year	17.2	18.2	22.0	25.2	28.1	26.1	23.9	25.9	26.9	28.3	29.7	30.0	28.9	28.4	28.1	29.4										
12-year	16.6	17.6	21.5	24.6	27.8	25.2	23.1	25.2	26.1	27.5	29.0	29.5	28.3	27.9	27.7											
13-year	16.2	17.0	20.9	24.1	26.6	24.8	22.8	24.4	25.6	26.8	28.4	29.0	27.6	27.3												
14-year	15.9	16.6	20.3	23.5	26.0	24.6	22.5	23.9	25.1	26.4	27.8	28.4	26.9													
15-year	15.4	16.1	19.6	22.9	25.3	24.0	22.4	22.6	24.9	26.1	27.4	28.0														
16-year	15.1	15.6	19.3	22.5	24.2	23.3	21.6	22.3	24.7	25.2	26.5															
17-year	14.7	15.3	18.8	22.0	24.0	23.2	21.2	22.1	24.4	25.0																
18-year	14.4	14.9	18.3	21.5	23.8	22.8	21.0	21.8	23.9																	
19-year	14.1	14.4	17.9	21.3	23.1	22.5	20.5	21.2																		
20-year	13.8	14.0	17.6	20.7	23.0	22.5	20.1																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 3.10
Cancer of the Brain and Other Nervous System (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	45.7	48.4	50.9	54.3	54.6	54.6	54.2	53.8	55.3	54.1	54.8	58.8	57.4	58.9	57.9	63.5	62.0	62.9	63.5	63.6	62.4	63.2	65.8	64.6	60.8	57.7
2-year	31.3	33.5	36.1	39.7	40.3	39.7	39.4	39.9	38.9	39.5	38.5	46.2	42.1	44.4	44.8	47.2	45.8	46.2	46.1	47.2	45.0	47.2	47.4	44.8	42.5	
3-year	26.4	29.3	31.6	35.4	37.2	35.8	35.0	35.3	34.3	35.8	35.3	39.2	36.3	39.3	39.2	39.8	39.4	40.4	41.1	39.6	39.4	38.8	38.4	36.9		
4-year	24.0	26.0	28.5	33.1	35.9	33.6	32.8	31.9	31.6	33.3	32.5	36.3	34.5	36.7	36.7	36.9	34.7	35.7	37.1	36.4	35.4	35.3	35.8			
5-year	21.6	23.6	26.4	31.1	34.8	31.3	30.5	30.2	29.9	32.1	30.8	33.6	33.2	35.7	34.5	35.4	33.3	34.3	35.1	33.8	32.8	33.1				
6-year	20.0	22.2	25.0	30.0	33.4	29.7	28.3	28.2	28.5	30.6	29.9	31.6	31.8	34.2	31.8	33.4	31.3	32.3	33.0	31.5	31.7					
7-year	19.0	21.2	23.8	28.6	32.1	28.7	27.2	27.1	27.9	28.4	28.8	30.5	30.2	32.6	30.4	32.6	30.5	31.2	32.2	30.1						
8-year	17.9	20.2	22.6	27.8	30.9	27.1	26.7	26.3	27.0	27.7	27.8	29.6	28.9	31.2	28.9	30.7	29.4	30.2	31.7							
9-year	16.9	19.1	22.0	26.9	30.1	26.0	26.0	25.2	25.4	26.7	27.3	28.7	28.4	30.1	28.3	29.5	28.5	29.0								
10-year	16.3	18.1	21.1	26.2	29.1	25.6	24.9	24.6	24.6	25.6	27.3	28.0	27.9	28.8	27.7	28.6	27.5									
11-year	15.6	17.5	20.6	25.4	28.3	24.8	24.5	23.9	23.2	24.8	26.3	26.8	27.9	27.7	27.1	27.9										
12-year	15.0	17.0	20.1	24.8	28.0	23.6	23.3	23.0	22.6	23.9	25.2	26.0	27.2	27.1	26.4											
13-year	14.6	16.5	19.6	24.4	26.5	23.3	23.2	22.7	22.4	23.7	24.9	25.2	26.3	25.9												
14-year	14.5	15.9	19.0	23.6	25.5	23.3	22.8	22.1	21.9	23.2	24.4	24.8	25.4													
15-year	14.0	15.5	18.5	22.9	24.8	22.8	22.7	21.1	21.7	22.9	24.3	24.4														
16-year	13.9	14.9	18.1	22.6	23.1	21.9	22.2	20.9	21.5	22.3	23.6															
17-year	13.5	14.5	17.8	22.0	22.8	21.6	21.8	20.7	21.2	21.8																
18-year	13.3	14.2	17.2	21.7	22.6	21.4	21.8	20.2	20.5																	
19-year	12.9	13.8	16.8	21.4	21.8	21.1	21.4	19.7																		
20-year	12.7	13.5	16.6	21.0	21.6	21.0	20.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 3.11
Cancer of the Brain and Other Nervous System (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	47.0	46.1	50.4	50.4	49.9	51.3	47.3	53.9	55.4	57.5	60.0	59.6	57.1	54.3	59.2	58.9	62.3	61.7	61.1	60.0	60.4	63.0	62.8	62.0	65.0	60.8
2-year	33.6	34.3	37.9	37.9	37.7	42.0	36.2	41.9	42.5	44.7	47.4	47.1	45.3	41.2	46.9	47.5	49.4	47.3	48.0	46.2	45.9	45.5	49.9	47.1	49.5	
3-year	29.9	29.5	34.0	34.3	34.4	37.6	33.0	36.3	38.4	41.6	43.8	42.3	40.3	36.8	40.9	41.8	43.1	41.8	41.9	40.3	40.2	38.6	42.6	40.1		
4-year	26.8	27.3	31.8	31.8	32.2	35.7	30.8	34.4	36.6	39.7	42.1	40.6	37.7	35.1	37.8	39.9	41.0	39.9	39.6	37.7	37.2	35.7	39.8			
5-year	25.1	25.9	30.0	30.3	31.3	34.1	29.8	32.1	35.5	38.4	39.8	39.2	35.3	34.1	35.4	36.6	38.3	38.2	37.8	34.9	35.2	33.6				
6-year	23.4	24.2	28.6	29.0	30.7	33.2	28.2	31.7	33.9	36.9	38.7	38.0	34.1	32.4	34.2	35.1	36.3	37.0	36.6	33.6	33.8					
7-year	22.4	22.9	27.5	27.9	30.1	32.3	27.9	30.7	32.9	35.9	37.4	37.4	32.8	31.3	33.7	34.3	35.2	35.4	35.4	32.7						
8-year	21.4	21.7	26.2	26.9	29.7	31.1	26.7	29.8	32.4	35.2	36.7	36.5	32.1	30.8	32.5	33.2	34.5	34.4	34.5							
9-year	20.4	20.9	25.2	26.2	29.6	29.9	25.1	29.4	32.1	34.4	35.9	35.4	30.9	30.2	31.5	32.2	33.5	33.3								
10-year	19.7	19.9	24.5	25.6	28.6	29.2	24.1	28.4	32.0	34.0	34.8	34.5	30.6	29.8	30.7	31.9	32.8									
11-year	19.2	18.9	23.8	25.0	27.6	27.8	23.2	28.3	31.7	32.8	34.1	34.0	30.2	29.3	29.3	31.4										
12-year	18.7	18.4	23.2	24.3	27.4	27.2	22.8	27.5	30.8	32.0	33.7	33.8	29.5	29.0	29.3											
13-year	18.3	17.7	22.4	23.9	26.7	26.8	22.4	26.3	29.6	30.8	32.9	33.6	29.1	28.8												
14-year	17.7	17.5	21.8	23.2	26.5	26.2	22.2	25.9	29.3	30.4	32.2	32.9	28.7													
15-year	17.2	16.9	21.1	22.8	25.9	25.6	21.9	24.3	29.1	30.1	31.1	32.4														
16-year	16.7	16.4	20.8	22.4	25.6	25.0	20.8	23.8	28.9	29.0	30.2															
17-year	16.2	16.2	20.2	21.9	25.4	25.0	20.4	23.6	28.6	29.0																
18-year	15.9	15.8	19.6	21.4	25.3	24.7	19.8	23.6	28.2																	
19-year	15.7	15.2	19.2	21.0	24.7	24.4	19.3	23.0																		
20-year	15.2	14.8	18.9	20.4	24.6	24.3	19.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 4.14
Cancer of the Female Breast (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

Survival Time	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	94.4	95.1	96.3	97.1	97.1	96.9	97.3	97.5	97.5	97.5	97.3	97.5	97.8	97.8	97.7	97.7	97.7	97.7	97.8	97.6	97.8	98.1	98.0	97.6	98.1	97.7
2-year	88.7	89.9	92.3	93.8	94.0	93.6	94.3	95.0	95.2	95.3	95.0	95.5	95.5	95.6	95.7	95.7	96.1	95.7	96.0	95.8	96.1	96.0	96.2	96.0	96.4	
3-year	83.1	84.6	88.3	90.5	91.4	90.9	92.0	92.9	93.1	93.4	92.8	93.6	93.3	93.3	93.8	93.8	94.0	93.8	94.0	94.0	94.5	94.1	94.5	94.3		
4-year	78.5	80.2	85.1	87.8	88.8	88.9	89.9	91.1	91.3	91.6	91.0	91.8	91.6	91.4	91.9	92.3	92.5	92.4	92.6	92.3	92.8	92.6	93.1			
5-year	74.6	76.3	82.2	85.6	86.8	86.6	88.4	89.5	89.6	90.1	89.4	90.2	89.8	90.0	90.5	90.8	91.2	90.7	91.4	91.0	91.5	91.2				
6-year	71.4	73.4	79.8	83.8	85.4	85.0	86.9	88.3	88.1	89.0	88.2	88.8	88.6	88.7	89.2	89.7	89.9	89.5	90.5	90.1	90.3					
7-year	68.7	70.7	77.9	82.1	83.8	83.8	85.9	87.1	87.0	87.7	86.9	87.7	87.5	87.5	88.1	88.8	89.0	88.4	89.4	89.3						
8-year	66.5	68.4	76.3	81.0	82.6	82.4	85.0	85.8	85.9	86.4	86.0	86.5	86.4	86.6	87.4	87.8	88.1	87.4	88.6							
9-year	64.3	66.4	74.6	79.8	81.4	81.3	83.7	84.6	85.0	85.3	85.2	85.6	85.9	85.7	86.6	86.9	87.1	86.5								
10-year	62.5	64.5	73.3	78.7	80.6	80.4	82.6	83.5	84.2	84.5	84.4	85.1	84.8	84.9	85.8	86.0	86.2									
11-year	60.8	62.7	72.1	77.7	79.5	79.5	81.8	82.7	83.0	83.6	83.7	84.5	84.0	84.4	85.1	85.3										
12-year	59.3	61.6	70.9	76.7	78.5	78.6	81.1	82.1	82.5	82.6	82.5	83.5	83.4	83.9	84.2											
13-year	58.1	60.4	69.8	75.8	77.5	77.6	80.1	81.1	81.8	81.8	81.9	82.7	82.9	83.0												
14-year	57.1	59.2	69.0	75.0	76.9	76.9	79.4	80.5	80.8	81.2	81.1	82.2	82.2													
15-year	56.1	58.3	68.1	74.2	76.0	76.2	78.5	79.5	80.3	80.5	80.5	81.7														
16-year	55.2	57.3	67.3	73.3	75.0	75.6	77.5	78.9	79.5	80.0	79.9															
17-year	54.2	56.3	66.5	72.4	74.2	74.9	77.1	78.2	79.0	79.4																
18-year	53.4	55.6	65.7	71.7	73.0	74.2	76.4	77.7	78.1																	
19-year	52.6	55.0	65.1	71.1	72.5	73.2	75.5	76.7																		
20-year	51.9	54.2	64.5	70.3	71.9	72.4	74.6																			
21-year	51.1	53.5	63.8	69.4	71.2	71.2																				
22-year	50.6	52.8	63.2	68.7	70.5																					
23-year	50.1	52.2	62.6	68.2																						
24-year	49.4	51.6	62.0																							
25-year	48.7	51.1	61.4																							
26-year	48.4	50.7	60.6																							
27-year	48.0	50.1	59.7																							
28-year	47.6	49.6	59.1																							
29-year	47.1	48.8																								
30-year	46.7	48.2																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 4.15
Cancer of the Female Breast (Ages <50, Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

Survival Time	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	96.6	97.4	97.8	98.0	97.9	97.7	97.9	98.3	98.6	98.4	98.2	98.5	98.8	98.5	98.8	98.8	99.0	98.7	98.8	98.6	99.1	98.8	98.5	98.8	98.9	98.5
2-year	90.8	91.2	92.5	93.8	93.6	93.8	94.6	95.1	95.0	95.5	94.5	95.7	96.1	95.9	96.1	96.8	97.1	96.3	96.5	96.3	97.1	96.6	96.6	96.6	96.9	
3-year	84.5	85.5	87.4	89.7	90.0	90.7	91.6	92.2	92.2	93.0	91.7	93.3	93.7	93.3	93.8	94.8	95.1	94.1	94.3	94.0	95.2	94.4	94.9	94.8		
4-year	80.0	80.9	83.4	86.3	86.7	88.2	88.7	89.5	90.1	90.6	89.3	90.6	91.8	91.2	92.2	93.1	93.5	92.5	92.6	92.0	93.1	92.6	93.2			
5-year	76.1	77.0	80.1	83.8	84.2	85.5	86.7	87.3	88.1	89.0	87.3	88.7	89.8	89.4	90.7	91.2	91.9	91.0	91.2	90.7	91.5	91.1				
6-year	73.2	74.2	77.9	81.8	82.3	83.3	84.7	85.4	86.1	87.3	85.3	86.9	88.1	87.9	89.4	89.7	90.5	89.6	89.8	89.6	90.2					
7-year	70.9	71.6	75.8	80.0	80.3	81.7	83.2	83.7	84.8	85.7	83.7	85.4	86.8	86.4	88.3	88.8	89.1	88.4	88.5							
8-year	68.9	69.5	74.2	78.5	78.8	80.1	81.5	82.4	83.6	83.8	82.2	84.3	85.7	85.3	87.4	87.7	88.2	87.5	87.7							
9-year	67.0	67.8	72.6	77.4	77.7	79.1	80.5	80.9	82.6	82.6	81.1	83.3	85.0	84.4	86.5	87.0	87.1	86.5								
10-year	65.6	66.4	71.5	76.3	76.4	77.7	79.3	79.8	81.6	81.6	80.3	82.5	84.1	83.5	85.7	86.1	86.4									
11-year	64.2	65.1	70.4	75.4	75.6	76.8	78.2	79.0	80.4	80.8	79.4	81.8	83.1	83.0	85.2	85.8										
12-year	62.9	64.1	69.3	74.5	74.7	76.1	77.4	78.2	80.0	80.1	78.6	80.7	82.6	82.4	84.4											
13-year	61.7	63.1	68.3	73.5	73.8	74.8	76.4	77.6	79.3	79.4	77.8	80.1	82.0	81.9												
14-year	61.0	62.0	67.6	72.8	73.1	74.2	75.4	77.3	78.1	79.0	77.4	79.5	81.4													
15-year	60.2	61.2	66.9	72.2	72.4	73.5	74.9	76.7	77.6	78.4	76.9	79.2														
16-year	59.5	60.5	66.2	71.5	71.6	73.1	74.2	76.1	77.1	77.9	76.4															
17-year	59.0	59.9	65.6	70.8	70.6	72.5	73.7	75.5	76.2	77.3																
18-year	58.4	59.3	65.0	70.2	69.6	72.1	72.8	75.2	75.6																	
19-year	57.8	58.8	64.4	69.7	68.7	71.6	72.3	74.8																		
20-year	57.1	58.3	64.0	69.2	68.2	71.0	71.7																			
21-year	56.4	57.5	63.4	68.7	67.7	70.1																				
22-year	55.9	56.8	62.9	68.4	66.9																					
23-year	55.3	56.2	62.5	68.0																						
24-year	54.7	55.7	62.1																							
25-year	54.1	55.2	61.6																							
26-year	53.8	54.7	61.0																							
27-year	53.2	54.2	60.4																							
28-year	52.8	53.6	59.9																							
29-year	52.5	53.1																								
30-year	52.1	52.7																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 4.16
Cancer of the Female Breast (Ages 50+, Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	93.7	94.4	95.8	96.7	96.8	96.7	97.1	97.2	97.2	97.2	97.0	97.2	97.5	97.5	97.4	97.4	97.3	97.4	97.5	97.3	97.4	97.9	97.8	97.2	97.8	97.5
2-year	88.1	89.6	92.3	93.8	94.1	93.6	94.2	94.9	95.2	95.3	95.2	95.4	95.3	95.5	95.6	95.3	95.7	95.5	95.9	95.7	95.7	95.9	96.1	95.8	96.2	
3-year	82.7	84.3	88.7	90.7	91.9	91.0	92.2	93.2	93.4	93.6	93.2	93.7	93.2	93.3	93.8	93.5	93.7	93.7	93.9	94.0	94.2	94.0	94.4	94.2		
4-year	78.0	80.0	85.7	88.4	89.6	89.2	90.3	91.6	91.8	92.0	91.6	92.2	91.5	91.5	91.9	92.1	92.2	92.4	92.5	92.4	92.7	92.6	93.0			
5-year	74.1	76.1	82.9	86.3	87.8	87.1	88.9	90.3	90.1	90.6	90.1	90.7	89.8	90.2	90.5	90.7	91.0	90.7	91.5	91.0	91.5	91.3				
6-year	70.8	73.1	80.5	84.6	86.6	85.6	87.8	89.4	88.8	89.6	89.2	89.5	88.9	88.9	89.2	89.8	89.8	89.5	90.7	90.3	90.3					
7-year	68.0	70.4	78.6	82.9	85.1	84.6	86.9	88.4	87.8	88.4	88.0	88.5	87.8	88.0	88.0	88.8	89.0	88.4	89.7	89.6						
8-year	65.6	68.0	77.1	81.9	84.1	83.3	86.3	87.2	86.7	87.4	87.3	87.3	86.6	87.1	87.4	87.8	88.1	87.4	88.9							
9-year	63.3	65.9	75.3	80.7	82.8	82.2	84.9	86.1	85.9	86.3	86.8	86.5	86.3	86.2	86.7	86.9	87.1	86.5								
10-year	61.3	63.8	73.9	79.6	82.2	81.4	83.9	84.9	85.1	85.5	86.0	86.1	85.1	85.4	85.9	86.0	86.2									
11-year	59.6	61.9	72.7	78.5	81.0	80.6	83.2	84.1	84.0	84.6	85.2	85.5	84.4	85.0	85.1	85.1										
12-year	58.0	60.6	71.4	77.6	80.0	79.6	82.5	83.6	83.4	83.6	84.0	84.5	83.8	84.5	84.1											
13-year	56.7	59.4	70.3	76.7	78.8	78.7	81.5	82.5	82.7	82.8	83.5	83.6	83.3	83.5												
14-year	55.6	58.1	69.4	75.7	78.3	78.0	80.9	81.7	81.7	82.1	82.4	83.2	82.6													
15-year	54.4	57.2	68.4	74.9	77.4	77.4	79.9	80.5	81.4	81.3	81.7	82.5														
16-year	53.4	56.0	67.5	74.0	76.3	76.7	78.8	79.8	80.4	80.7	81.2															
17-year	52.1	54.8	66.6	72.9	75.5	75.9	78.3	79.2	80.1	80.1																
18-year	51.3	54.1	65.8	72.1	74.3	74.9	77.9	78.5	79.0																	
19-year	50.4	53.3	65.2	71.4	74.1	73.7	76.7	77.2																		
20-year	49.6	52.5	64.5	70.5	73.4	72.8	75.6																			
21-year	48.8	51.7	63.7	69.3	72.5	71.3																				
22-year	48.3	51.0	63.0	68.4	71.9																					
23-year	47.7	50.4	62.4	67.7																						
24-year	47.0	49.7	61.6																							
25-year	46.1	49.2	60.8																							
26-year	45.8	48.9	59.7																							
27-year	45.4	48.3	58.7																							
28-year	45.1	47.7	57.9																							
29-year	44.5	46.6																								
30-year	44.0	45.7																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 5.9
Cancer of the Cervix Uteri (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	86.5	87.2	87.0	87.7	89.3	88.6	89.1	88.0	89.7	88.3	85.8	87.8	86.7	86.2	87.1	86.3	88.3	87.1	88.8	87.8	85.1	87.1	86.5	86.9	88.3	87.2
2-year	77.9	77.3	77.9	79.5	82.2	81.8	80.0	81.4	82.8	81.0	79.0	79.7	79.1	77.3	77.2	78.6	78.7	79.5	79.9	80.0	76.3	78.0	79.8	80.5	81.3	
3-year	73.2	72.7	73.1	75.2	78.8	77.8	75.9	76.8	78.4	76.5	75.5	74.9	75.8	73.1	72.3	75.8	74.8	75.7	74.5	73.9	72.4	73.9	74.8	76.3		
4-year	70.3	69.7	70.5	72.2	76.0	74.7	73.3	74.1	76.1	74.3	72.4	72.5	72.5	70.2	69.9	73.7	73.0	72.8	71.4	71.4	69.6	70.9	71.3			
5-year	68.3	67.6	68.5	70.4	74.9	73.6	71.8	72.6	74.4	72.2	70.4	69.9	70.4	67.5	68.0	72.0	71.2	70.1	68.8	69.5	67.9	69.3				
6-year	67.1	65.7	67.2	69.0	73.6	72.9	70.2	71.0	73.4	71.1	69.1	68.9	68.8	66.0	67.1	70.1	70.5	67.7	67.8	68.2	65.5					
7-year	66.0	64.1	66.1	68.1	72.5	71.7	69.8	69.9	73.0	70.3	67.8	68.2	67.5	65.3	66.8	69.1	69.4	66.6	66.3	67.3						
8-year	65.1	62.8	65.0	67.2	71.8	70.8	69.3	69.5	72.6	70.0	67.4	67.6	66.2	63.6	65.8	68.6	68.2	65.8	65.9							
9-year	64.3	62.2	64.4	66.2	70.8	69.3	67.7	68.8	72.2	68.9	66.9	67.3	65.8	62.7	65.1	67.7	67.5	65.0								
10-year	63.1	61.4	63.7	65.6	70.0	68.8	67.1	67.8	70.9	68.3	66.7	66.6	65.5	61.8	64.5	67.5	67.0									
11-year	62.5	60.8	63.1	65.0	69.6	68.2	66.2	67.1	69.5	67.9	65.9	66.2	64.6	61.3	63.8	66.8										
12-year	61.7	60.0	62.3	64.4	68.6	67.1	66.0	66.1	68.8	66.9	64.9	65.5	64.3	60.7	63.6											
13-year	61.1	59.5	61.6	63.8	67.1	66.5	65.7	64.8	67.8	66.0	64.0	65.2	64.1	59.9												
14-year	60.5	58.7	61.0	63.2	66.6	65.8	65.4	64.6	67.2	65.6	63.6	64.7	63.5													
15-year	59.8	57.9	60.4	62.7	66.3	65.1	65.0	64.3	67.0	64.9	62.7	63.8														
16-year	59.3	57.2	59.7	62.0	65.4	64.7	63.4	64.1	66.7	63.9	62.2															
17-year	58.6	56.3	58.9	61.6	64.6	63.8	62.6	63.8	66.5	63.5																
18-year	58.2	55.5	58.7	60.8	64.1	62.9	62.6	63.2	65.9																	
19-year	57.4	54.9	58.3	60.3	63.7	62.2	62.0	63.1																		
20-year	56.6	54.2	57.3	59.9	63.3	61.0	61.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 6.15
Cancer of the Colon and Rectum (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																													
	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016				
1-year	73.9	77.0	79.9	80.8	79.5	81.1	80.2	81.9	82.4	81.9	83.0	83.2	82.5	83.4	83.9	83.5	84.1	84.6	84.3	84.2	83.8	84.0	84.1	84.5	85.0	84.5				
2-year	63.0	66.8	71.1	72.5	71.4	73.3	72.3	74.6	74.7	74.4	75.5	76.9	75.8	76.6	76.9	77.2	77.3	78.3	77.3	77.4	76.9	76.5	77.5	78.1	78.0					
3-year	56.9	60.5	65.7	67.2	65.9	68.5	67.3	69.5	70.1	70.2	71.1	72.2	70.8	71.5	72.6	72.2	72.4	73.4	72.6	72.3	72.0	71.6	72.3	72.9						
4-year	53.0	56.5	61.9	63.3	62.0	65.0	64.2	66.3	66.9	67.0	68.2	69.0	67.7	68.4	68.9	68.5	69.2	70.1	69.1	68.9	68.5	68.3	68.6							
5-year	50.4	53.7	59.3	60.7	59.8	62.5	61.4	63.2	64.6	64.8	66.0	66.2	65.4	65.8	66.2	66.2	66.8	67.5	66.6	66.4	65.9	65.5								
6-year	48.6	51.8	57.5	59.0	57.3	61.2	59.7	61.4	63.1	63.1	64.1	64.4	63.6	63.9	64.6	64.7	64.8	65.6	64.8	64.2	64.2									
7-year	47.4	50.3	56.0	57.5	55.6	60.2	58.8	60.1	62.0	61.7	62.3	63.1	62.1	62.7	63.0	63.0	63.4	63.9	63.2	62.8										
8-year	46.4	49.3	55.0	56.3	54.9	59.5	58.0	59.3	60.7	60.3	61.3	62.1	61.4	61.8	62.0	61.8	62.2	62.8	61.9											
9-year	45.8	48.4	54.1	55.4	54.6	58.7	57.1	58.0	59.8	59.5	60.4	60.8	60.2	61.0	61.3	60.8	61.6	61.7												
10-year	45.1	47.8	53.3	54.6	53.9	58.1	56.4	57.3	59.2	59.0	59.6	60.5	59.1	60.1	60.4	60.4	60.4													
11-year	44.5	47.2	52.9	54.1	53.1	57.5	55.6	56.5	58.6	58.1	59.2	59.8	58.3	58.9	59.9	60.2														
12-year	44.1	46.8	52.5	53.6	52.4	57.4	55.3	56.2	57.9	57.7	58.3	58.8	58.0	58.9	59.1															
13-year	43.7	46.4	51.7	53.0	51.7	56.8	55.0	55.7	57.7	57.2	57.7	58.4	57.0	57.9																
14-year	43.2	45.9	51.4	52.6	51.2	55.9	54.1	55.6	57.0	56.8	57.1	57.5	56.2																	
15-year	42.8	45.6	50.9	52.2	50.6	55.3	53.6	55.5	56.6	56.6	56.6	56.6	56.6																	
16-year	42.4	45.2	50.7	52.2	50.1	55.0	52.8	55.2	55.7	56.0	55.9																			
17-year	42.0	44.7	50.0	51.6	49.6	53.9	52.3	54.8	54.7	55.6																				
18-year	41.5	44.3	49.7	51.1	49.0	53.8	51.7	54.5	54.2																					
19-year	41.5	43.9	49.5	50.5	48.2	53.1	51.5	54.4																						
20-year	41.4	43.7	49.0	49.9	47.3	52.3	51.1																							
21-year	40.9	42.9	49.0	49.4	46.5	51.9																								
22-year	40.6	42.4	48.9	48.7	45.9																									
23-year	40.1	42.3	48.4	48.3																										
24-year	39.8	41.9	47.8																											
25-year	39.5	41.4	47.5																											
26-year	39.0	41.0	46.9																											
27-year	38.5	40.6	46.4																											
28-year	38.4	40.2	45.5																											
29-year	38.0	40.0																												
30-year	37.5	39.7																												

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 6.16
Cancer of the Colon and Rectum (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	74.0	77.9	80.9	81.7	81.0	82.5	81.2	82.0	83.5	83.5	85.0	83.1	83.2	84.4	84.5	84.7	85.5	86.2	85.6	85.4	84.2	84.7	84.5	85.5	86.3	84.3
2-year	63.0	67.2	72.4	73.3	72.8	74.9	73.5	74.5	75.4	76.3	77.4	76.6	76.3	77.9	77.9	78.3	78.5	79.3	78.6	78.6	77.2	77.1	77.6	79.0	79.2	
3-year	56.3	60.6	66.8	67.9	67.0	69.7	68.3	68.9	70.6	72.0	72.8	72.1	71.1	72.4	73.2	73.0	73.4	74.1	73.6	73.1	71.6	71.7	71.8	73.6		
4-year	52.2	56.2	62.5	63.5	63.2	65.3	65.3	65.4	67.6	69.0	69.9	68.7	67.9	69.1	68.8	68.8	69.7	70.5	69.6	69.0	68.2	68.2	67.9			
5-year	49.3	53.2	59.7	60.5	60.1	62.4	62.3	62.4	65.0	66.1	67.3	65.6	65.4	66.2	65.8	66.5	66.9	67.8	66.8	66.3	65.4	65.5				
6-year	47.2	51.2	57.8	58.6	57.4	61.0	60.3	60.2	63.8	63.8	64.9	63.4	62.9	64.2	63.7	64.7	64.4	65.6	65.0	63.6	63.8					
7-year	45.9	49.5	56.3	56.8	55.7	59.8	59.7	58.7	62.6	62.0	63.1	61.8	61.5	62.5	61.7	63.1	62.9	63.9	63.5	62.1						
8-year	44.8	48.5	55.4	55.4	55.2	59.4	58.4	57.5	61.2	60.2	62.4	60.4	60.3	61.5	60.6	61.8	61.6	62.8	62.3							
9-year	44.1	47.5	54.3	54.5	54.7	58.9	57.3	56.1	60.1	59.8	61.4	59.4	59.0	60.8	59.8	60.6	60.8	61.0	61.5							
10-year	43.3	46.7	53.6	53.6	54.1	58.0	56.7	55.5	59.2	58.7	60.6	59.3	57.7	59.8	59.2	60.0	59.5									
11-year	42.8	46.1	53.2	53.2	53.1	57.3	55.9	54.5	58.7	57.9	59.7	58.4	56.7	58.7	58.5	59.4										
12-year	42.1	45.7	52.6	52.7	52.3	57.0	55.7	54.0	57.9	57.4	59.2	57.4	56.1	58.6	57.8											
13-year	41.7	45.4	51.9	51.9	51.8	56.7	55.3	53.4	57.3	57.0	58.3	56.9	55.2	57.8												
14-year	41.0	44.7	51.8	51.4	50.8	55.7	54.6	53.2	56.4	56.5	57.9	56.0	54.3													
15-year	40.8	44.6	51.2	50.9	49.8	55.4	54.3	53.1	56.1	56.2	57.4	55.0														
16-year	40.3	44.3	51.1	50.9	49.3	55.2	53.2	52.8	54.9	55.9	56.8															
17-year	39.8	43.7	50.6	50.4	48.9	54.2	52.0	52.2	53.9	55.7																
18-year	39.4	43.3	50.2	49.9	48.0	54.2	51.3	52.2	53.6																	
19-year	39.4	42.9	49.9	49.6	47.4	53.9	51.2	52.2																		
20-year	39.2	42.6	49.5	49.1	45.9	53.9	51.0																			
21-year	39.0	41.7	49.4	48.6	45.3	53.5																				
22-year	38.5	41.1	49.4	48.0	45.1																					
23-year	38.1	41.0	48.8	47.5																						
24-year	37.7	40.4	48.1																							
25-year	37.3	40.2	47.6																							
26-year	36.4	39.4	46.9																							
27-year	35.9	39.0	46.4																							
28-year	35.7	38.7	45.1																							
29-year	35.3	38.4																								
30-year	34.4	37.7																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 6.17
Cancer of the Colon and Rectum (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

Survival Time	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	73.9	76.2	78.8	79.9	78.1	79.7	79.1	81.8	81.3	80.2	81.0	83.3	81.7	82.4	83.3	82.4	82.6	82.9	83.0	82.9	83.4	83.4	83.7	83.3	83.6	84.8
2-year	63.0	66.4	69.8	71.6	70.1	71.6	71.1	74.6	73.9	72.4	73.6	77.2	75.3	75.2	75.9	76.1	76.0	77.2	76.0	76.2	76.5	75.8	77.5	77.1	76.7	
3-year	57.4	60.3	64.6	66.5	64.8	67.3	66.2	70.0	69.5	68.4	69.4	72.4	70.5	70.5	72.0	71.4	71.4	72.6	71.6	71.4	72.5	71.4	72.9	72.3		
4-year	53.9	56.7	61.3	63.1	60.9	64.7	63.1	67.3	66.3	65.0	66.5	69.3	67.6	67.6	69.0	68.2	68.6	69.8	68.5	68.8	68.8	68.4	69.4			
5-year	51.4	54.0	58.9	60.9	59.4	62.7	60.5	64.1	64.2	63.5	64.7	66.8	65.3	65.3	66.7	65.9	66.7	67.1	66.4	66.6	66.4	65.5				
6-year	49.8	52.3	57.2	59.5	57.3	61.5	59.1	62.7	62.4	62.4	63.2	65.5	64.3	63.5	65.4	64.7	65.2	65.6	64.7	64.9	64.7					
7-year	48.8	51.0	55.7	58.1	55.4	60.6	57.9	61.4	61.3	61.3	61.3	64.4	62.8	63.0	64.4	62.8	64.0	63.8	62.8	63.5						
8-year	47.9	50.1	54.7	57.1	54.6	59.4	57.6	61.0	60.1	60.4	60.1	63.8	62.5	62.1	63.4	61.8	62.9	62.8	61.5							
9-year	47.3	49.3	53.9	56.3	54.4	58.5	56.9	59.8	59.5	59.3	59.2	62.2	61.5	61.1	62.9	61.1	62.3	61.9								
10-year	46.7	48.7	53.0	55.6	53.8	58.1	56.1	59.1	59.2	59.3	58.7	61.6	60.6	60.3	61.7	60.8	61.4									
11-year	46.1	48.2	52.5	55.0	53.0	57.7	55.4	58.3	58.4	58.2	58.7	61.0	60.0	59.1	61.4	60.8										
12-year	45.9	47.8	52.3	54.4	52.4	57.6	54.8	58.3	57.9	57.8	57.4	60.3	59.8	59.1	60.5											
13-year	45.5	47.4	51.4	54.1	51.7	56.9	54.5	57.9	57.9	57.2	56.9	59.9	58.9	58.1												
14-year	45.1	47.0	51.0	53.8	51.5	56.1	53.7	57.7	57.7	57.1	56.3	59.0	58.2													
15-year	44.7	46.6	50.5	53.5	51.3	55.3	52.6	57.7	57.1	57.1	55.7	58.2														
16-year	44.3	45.9	50.1	53.5	50.8	54.7	52.3	57.5	56.6	55.9	55.1															
17-year	43.9	45.6	49.5	52.7	50.1	53.6	52.3	57.3	55.6	55.4																
18-year	43.4	45.3	49.2	52.3	49.7	53.4	52.1	56.5	54.5																	
19-year	43.4	44.8	49.0	51.4	49.0	52.2	51.7	56.3																		
20-year	43.2	44.6	48.6	50.7	48.5	50.5	51.2																			
21-year	42.8	44.0	48.5	50.1	47.6	50.1																				
22-year	42.6	43.5	48.2	49.4	46.6																					
23-year	42.0	43.4	47.9	49.1																						
24-year	41.6	43.2	47.5																							
25-year	41.5	42.5	47.3																							
26-year	41.2	42.3	46.8																							
27-year	40.8	42.0	46.4																							
28-year	40.7	41.5	45.7																							
29-year	40.2	41.5																								
30-year	40.1	41.3																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 7.9
Cancer of the Corpus And Uterus, NOS (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	93.7	91.3	92.2	92.1	92.9	91.5	93.1	93.3	93.9	93.6	92.1	92.0	92.8	92.8	92.8	92.8	93.1	93.3	92.8	93.3	92.2	93.1	92.7	93.2	93.0	92.1
2-year	90.1	86.3	87.5	88.1	88.4	88.0	88.9	89.7	89.5	89.5	88.1	88.2	88.6	88.3	89.0	89.5	89.5	88.8	88.3	89.0	88.0	88.8	87.9	88.4	89.4	
3-year	87.8	83.4	84.9	85.6	86.0	85.7	86.2	86.6	87.2	87.4	85.7	85.6	86.1	85.4	85.8	87.2	87.3	86.1	85.8	87.0	85.5	85.7	85.5	85.5		
4-year	86.3	82.0	83.4	84.1	84.7	84.6	85.2	84.5	85.5	86.0	84.2	83.7	84.6	83.8	84.2	85.6	85.7	84.5	84.5	84.9	83.8	84.2	84.0			
5-year	85.8	80.9	82.3	83.3	84.1	83.2	84.1	83.2	84.3	85.2	83.1	82.8	83.4	82.6	82.9	84.9	83.9	82.9	83.2	83.5	82.6	83.0				
6-year	85.1	80.3	81.7	82.2	83.4	82.1	83.3	82.5	83.7	84.4	82.2	82.4	82.5	81.7	81.9	84.4	83.3	82.0	81.9	82.9	81.4					
7-year	84.7	79.8	81.3	81.9	82.9	81.3	82.6	82.2	82.6	83.9	81.5	81.3	82.0	80.7	81.3	83.8	82.4	81.3	81.3	82.3						
8-year	84.5	79.1	80.8	81.2	82.5	80.4	81.6	81.9	82.0	83.3	80.8	81.0	81.9	80.3	80.9	83.5	81.9	80.8	80.8							
9-year	84.5	78.7	80.5	81.0	81.5	79.7	81.0	81.2	82.0	83.2	80.1	80.3	81.4	79.9	80.4	83.2	81.5	80.4								
10-year	84.3	78.4	80.3	80.5	81.2	79.5	81.0	81.0	81.3	82.9	80.0	80.1	80.3	79.6	79.8	82.5	81.3									
11-year	84.1	78.1	80.0	80.4	81.1	79.5	81.0	81.0	81.2	81.9	79.2	80.1	80.1	79.1	78.9	82.3										
12-year	83.9	77.8	79.6	79.9	80.9	79.5	81.0	80.8	80.9	81.8	78.6	80.0	79.4	78.9	78.8											
13-year	83.9	77.5	79.2	79.6	80.7	79.2	81.0	80.6	80.4	81.5	78.1	80.0	78.8	78.2												
14-year	83.7	77.1	78.9	79.2	80.2	78.9	81.0	79.7	80.0	81.4	78.1	79.6	78.1													
15-year	83.7	76.8	78.6	79.1	79.7	78.4	81.0	79.7	80.0	80.2	78.1	79.4														
16-year	83.4	76.2	78.1	78.8	79.4	77.5	80.8	78.8	79.2	79.8	78.1															
17-year	83.4	75.9	78.0	78.3	79.0	76.5	80.2	78.0	78.9	79.3																
18-year	83.4	75.5	77.8	77.8	78.1	75.7	79.4	78.0	78.4																	
19-year	83.4	74.8	77.7	77.2	77.5	75.4	78.9	77.7																		
20-year	83.4	74.7	77.2	76.5	76.6	74.5	78.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 8.9
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	25.8	30.0	34.9	38.2	40.6	42.1	40.0	40.8	44.3	45.9	46.2	41.8	47.0	47.9	45.3	47.4	50.2	47.7	48.5	49.6	49.4	50.8	51.9	50.9	50.7	54.8
2-year	11.5	14.3	19.0	21.9	22.6	24.4	23.9	23.9	26.7	28.3	29.3	26.0	30.4	30.2	31.0	30.8	33.1	32.3	30.7	33.9	32.4	34.3	36.2	35.2	33.6	
3-year	7.7	10.0	13.5	16.3	16.8	18.7	17.5	18.4	20.7	23.7	23.0	20.9	24.0	25.8	24.3	25.1	27.1	24.5	24.6	27.8	24.7	26.9	29.0	27.8		
4-year	6.0	8.4	10.8	13.7	13.2	16.0	14.3	14.2	18.3	20.6	19.2	17.9	20.1	22.3	21.4	21.8	23.6	21.9	21.7	24.9	22.1	23.3	25.1			
5-year	4.9	7.0	9.5	12.2	11.6	15.1	12.4	12.6	17.4	18.9	17.8	16.8	17.2	20.2	18.9	20.1	21.8	18.8	20.7	22.6	18.9	20.6				
6-year	4.3	6.2	8.5	11.1	10.8	14.0	11.3	11.6	15.9	17.9	15.9	15.3	14.6	19.6	17.3	18.7	19.6	18.0	19.6	20.3	17.8					
7-year	3.8	5.6	7.7	10.2	10.2	13.0	10.6	11.3	14.7	16.6	14.8	14.9	13.3	18.3	15.7	17.9	18.4	16.8	18.7	19.3						
8-year	3.4	4.9	7.0	9.5	9.6	12.2	9.8	9.8	14.2	15.8	13.8	13.7	13.0	17.9	14.7	16.7	18.4	16.1	17.5							
9-year	3.2	4.5	6.4	9.0	8.9	11.5	9.3	9.6	13.5	15.6	13.6	13.2	12.9	16.5	13.7	15.9	17.6	15.7								
10-year	2.9	4.3	6.0	8.3	8.6	10.7	9.2	9.4	12.9	15.1	13.3	13.0	11.9	15.5	13.3	15.2	16.8									
11-year	2.7	4.0	5.4	7.8	7.6	9.7	9.0	9.1	12.7	14.3	12.3	13.0	11.9	14.9	13.2	14.0										
12-year	2.6	3.8	4.9	7.1	7.2	9.4	7.6	8.5	11.9	14.0	12.0	12.4	11.5	14.6	12.8											
13-year	2.4	3.5	4.6	7.0	6.8	9.4	7.0	8.5	11.0	12.2	11.8	11.4	11.3	13.8												
14-year	2.4	3.4	4.3	6.6	5.7	9.0	6.4	8.4	10.4	12.0	11.2	10.9	10.6													
15-year	2.4	3.4	4.2	6.2	5.5	8.8	6.0	7.6	10.2	11.6	11.2	10.9														
16-year	2.3	3.3	3.6	5.8	4.9	8.0	5.9	7.5	9.9	10.5	11.2															
17-year	2.3	3.0	3.2	5.5	4.9	7.1	5.9	7.5	9.7	10.1																
18-year	2.2	2.7	3.1	5.3	4.9	7.1	5.6	7.5	9.2																	
19-year	2.2	2.5	2.9	4.9	3.8	6.3	5.0	7.5																		
20-year	2.1	2.5	2.6	4.8	3.0	5.3	5.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 8.10
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	23.8	29.3	34.0	38.6	40.3	43.3	40.4	41.0	43.6	47.4	47.1	43.0	46.4	47.3	46.1	48.2	51.8	48.6	49.8	48.5	49.0	51.5	52.6	49.9	51.3	55.9
2-year	10.0	13.8	18.4	21.7	22.8	24.9	24.7	24.3	26.3	28.7	29.7	26.6	29.9	30.2	31.1	31.3	33.4	32.8	30.9	33.1	32.2	34.1	36.8	34.2	34.3	
3-year	6.8	9.3	13.1	16.4	16.7	18.4	18.3	18.6	20.6	23.6	23.2	21.3	23.3	25.7	24.2	25.8	27.4	24.8	25.3	26.9	23.8	26.3	29.3	26.6		
4-year	5.2	7.5	10.3	13.4	13.0	15.2	15.2	14.2	18.1	19.7	19.4	18.1	19.6	22.4	20.8	22.9	23.7	22.0	22.0	23.9	21.0	22.4	25.3			
5-year	4.2	6.0	9.3	12.0	11.8	14.0	13.0	12.1	17.2	17.4	18.2	16.8	16.8	20.0	18.4	21.0	21.9	18.8	21.0	22.0	17.6	19.5				
6-year	3.7	5.2	8.4	10.8	11.0	13.1	11.8	11.5	16.1	16.6	16.3	15.4	14.0	19.3	17.0	19.4	19.4	17.8	20.3	19.3	16.7					
7-year	3.3	4.5	7.6	10.2	10.7	12.1	11.5	11.2	15.0	15.5	15.4	15.3	12.7	18.3	15.3	18.7	17.9	16.6	19.3	18.2						
8-year	3.0	4.0	6.9	9.3	9.9	11.4	10.3	9.4	14.7	14.8	14.6	13.7	12.2	17.8	14.1	17.4	17.9	15.7	17.9							
9-year	2.8	3.7	6.5	8.8	9.6	10.8	10.0	9.3	14.1	14.5	14.2	13.3	12.2	17.1	13.5	16.3	17.3	15.7								
10-year	2.7	3.5	6.2	7.9	9.3	10.1	9.7	9.3	13.3	14.2	13.8	13.3	11.6	16.0	13.1	15.5	16.6									
11-year	2.5	3.3	5.7	7.5	8.4	9.1	9.5	9.0	13.2	13.3	13.0	13.3	11.6	15.3	13.0	14.4										
12-year	2.4	3.3	5.1	6.8	7.7	8.8	8.4	8.8	12.8	12.8	12.7	12.4	11.6	14.8	12.7											
13-year	2.3	3.1	4.6	6.8	7.1	8.8	7.8	8.8	11.8	10.7	12.6	11.4	11.4	14.1												
14-year	2.2	3.0	4.2	6.6	6.3	8.5	6.9	8.8	11.0	10.6	11.8	10.9	11.3													
15-year	2.2	3.0	4.0	6.1	5.9	8.5	6.4	7.8	10.8	9.9	11.8	10.9														
16-year	2.2	3.0	3.4	5.9	5.3	8.4	6.3	7.8	10.5	8.8	11.8															
17-year	2.2	2.9	2.9	5.5	5.3	7.3	6.2	7.8	10.4	8.7																
18-year	2.2	2.6	2.8	5.3	5.3	7.3	5.9	7.8	9.8																	
19-year	2.1	2.4	2.5	5.1	3.9	6.1	5.4	7.8																		
20-year	2.0	2.3	2.3	5.0	3.1	5.2	5.3																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 8.11
Cancer of the Esophagus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	31.0	31.6	37.1	37.2	41.4	38.4	38.8	39.9	46.6	41.2	42.9	38.2	49.0	50.1	42.2	44.7	44.5	44.3	44.2	54.0	50.8	48.2	49.0	54.7	48.4	50.0
2-year	15.3	15.6	20.7	22.4	22.2	22.9	21.6	22.5	28.3	27.0	27.9	24.3	32.4	30.1	31.0	29.1	32.2	30.2	30.2	37.3	33.2	35.1	33.9	39.1	30.8	
3-year	10.0	12.0	14.5	16.2	17.0	19.8	15.1	17.6	21.0	24.1	22.7	19.6	26.3	25.9	24.5	22.3	25.8	23.1	22.4	31.8	28.0	29.4	27.7	32.1		
4-year	7.9	10.5	12.3	14.5	13.7	18.2	11.7	14.3	18.7	23.4	18.6	17.3	21.9	21.9	23.5	18.3	22.8	21.5	21.0	29.3	26.1	26.9	24.2			
5-year	6.5	9.3	10.0	12.9	10.7	18.2	10.6	14.1	17.7	23.3	16.1	16.9	18.8	20.8	20.6	16.6	21.4	18.8	19.6	25.1	23.8	24.9				
6-year	5.9	8.6	8.5	11.8	9.9	16.3	9.8	11.9	15.0	22.2	14.2	14.8	16.6	20.5	18.4	15.9	20.4	18.4	17.3	24.6	22.1					
7-year	5.2	8.3	8.0	10.2	9.0	15.7	8.1	11.7	13.2	20.2	12.3	13.4	15.4	18.0	16.9	15.1	19.8	17.6	16.5	24.1						
8-year	4.4	7.1	7.3	10.0	8.7	14.2	8.1	10.8	11.9	19.1	10.7	13.1	15.4	17.7	16.6	14.4	19.3	17.5	15.9							
9-year	4.0	6.5	6.1	9.5	7.1	13.7	7.4	10.5	10.8	18.7	10.7	12.3	14.9	14.4	14.7	14.2	18.7	15.7								
10-year	3.5	6.4	5.3	9.2	6.6	12.3	7.4	9.5	10.8	17.8	10.7	12.0	12.7	13.6	13.9	13.8	17.1									
11-year	3.3	5.4	4.8	8.8	5.6	11.4	7.4	9.3	10.0	17.1	9.6	12.0	11.9	13.4	13.9	12.6										
12-year	3.1	5.2	4.6	7.9	5.6	11.0	5.2	7.6	8.1	17.1	9.1	12.0	10.2	13.4	13.3											
13-year	2.8	4.7	4.6	7.0	5.6	11.0	4.6	7.6	7.4	16.6	8.5	11.4	10.2	12.7												
14-year	2.8	4.5	4.5	6.4	4.2	10.5	4.6	6.8	7.4	16.3	8.5	10.8	7.4													
15-year	2.8	4.5	4.5	6.1	4.2	8.7	4.6	6.8	7.4	16.3	8.5	10.5														
16-year	2.7	3.8	4.2	5.5	3.7	6.8	4.6	6.4	7.3	15.6	8.5															
17-year	2.4	3.2	3.9	5.4	3.7	6.3	4.6	6.4	6.9	14.4																
18-year	2.2	3.0	3.8	5.2	3.7	6.3	4.6	6.4	6.3																	
19-year	2.2	2.8	3.7	4.5	3.4	6.3	3.7	6.4																		
20-year	2.2	2.8	3.3	4.2	2.6	5.5	3.7																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 9.9
Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	86.9	89.4	90.7	92.1	92.8	91.6	93.5	94.6	91.0	92.4	94.0	92.9	93.2	91.3	93.5	92.4	93.5	94.7	92.3	91.9	93.7	94.5	93.8	93.9	93.0	94.3
2-year	80.8	83.6	86.3	87.7	88.6	88.2	90.6	90.5	87.8	90.6	90.8	90.0	92.2	89.4	91.1	90.1	92.5	92.8	89.6	90.8	92.5	92.9	92.4	93.3	92.4	
3-year	76.5	80.1	83.0	85.1	86.7	86.8	89.5	88.3	85.4	88.7	88.5	88.4	90.1	87.3	90.5	89.0	90.8	91.7	88.4	89.8	91.4	91.4	91.3	92.5		
4-year	73.7	77.4	80.6	83.2	83.5	84.6	88.6	86.6	84.2	87.9	86.6	86.2	88.1	86.1	89.6	88.5	89.2	90.8	87.3	88.7	90.1	90.7	91.0			
5-year	71.4	74.8	79.0	81.6	82.2	83.3	87.3	84.9	83.4	86.4	85.2	85.2	87.5	85.9	88.3	88.1	88.6	90.5	85.7	86.8	89.8	90.3				
6-year	69.0	72.3	77.5	80.0	80.2	81.8	86.1	83.8	82.2	85.6	84.3	84.7	87.0	84.9	87.9	86.9	87.1	89.7	84.6	85.6	88.6					
7-year	67.1	70.7	75.2	78.6	79.0	80.1	85.4	83.0	80.9	84.9	84.3	83.4	86.4	84.3	87.4	86.3	85.4	88.0	83.8	85.5						
8-year	65.0	69.3	74.1	77.8	78.5	79.4	84.9	82.4	80.0	83.9	83.8	82.3	85.7	83.4	85.8	86.2	84.2	87.4	82.9							
9-year	62.9	67.9	73.3	76.9	77.3	78.3	84.8	81.5	79.0	83.0	82.8	81.3	84.8	83.1	85.5	85.8	84.2	86.6								
10-year	61.7	66.8	72.6	76.1	76.8	76.8	84.1	80.2	78.3	83.0	81.9	80.5	84.5	82.6	85.4	85.3	84.2									
11-year	60.4	65.8	71.7	75.5	75.7	75.8	83.6	79.6	78.0	82.9	81.3	79.0	84.1	82.4	84.9	84.9										
12-year	59.5	64.7	70.9	74.8	75.6	74.9	83.2	79.2	77.6	82.4	81.3	78.3	83.5	81.9	84.6											
13-year	58.7	63.8	69.7	74.1	75.5	74.4	82.0	79.0	76.4	82.0	79.6	77.5	82.3	81.2												
14-year	57.5	62.7	68.7	73.6	75.5	73.6	80.5	78.4	76.3	81.2	79.5	77.5	81.9													
15-year	56.5	61.9	67.5	73.1	75.4	73.3	79.7	77.2	75.7	78.9	78.0	76.9														
16-year	55.7	60.9	66.9	72.6	75.2	73.3	79.5	77.1	74.1	77.9	77.2															
17-year	54.7	59.5	66.3	72.0	74.9	72.6	78.4	76.7	73.4	77.3																
18-year	53.9	58.6	65.6	71.3	73.9	72.1	77.4	76.1	73.2																	
19-year	53.2	58.1	64.9	70.4	73.9	71.8	76.6	75.6																		
20-year	52.1	57.0	64.2	69.9	73.5	70.9	76.3																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 9.10
Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	85.9	89.4	89.8	91.2	92.2	92.7	92.1	94.3	89.3	90.9	92.9	90.8	93.0	89.5	93.6	91.5	92.7	93.5	93.1	90.3	92.9	93.5	91.9	93.5	92.0	92.4
2-year	80.2	83.6	84.8	85.9	88.0	88.7	88.9	91.0	86.4	88.4	89.1	88.5	92.2	87.9	91.0	89.0	91.6	91.4	89.5	89.5	91.8	92.1	90.0	92.9	90.8	
3-year	75.3	80.4	81.4	83.1	85.4	87.1	87.7	88.8	83.6	85.5	86.6	86.7	89.9	86.2	90.2	88.5	89.5	90.6	88.0	88.1	90.7	90.8	89.0	92.6		
4-year	71.7	77.3	78.4	80.8	81.1	85.1	87.2	86.5	82.8	83.8	85.2	84.9	86.8	85.7	88.9	87.4	87.9	89.2	87.2	87.2	89.5	90.2	88.7			
5-year	69.2	74.1	76.5	79.1	80.2	83.1	86.1	84.5	82.5	81.9	83.7	83.8	86.1	85.7	87.6	86.8	87.2	89.2	84.9	85.3	88.9	89.7				
6-year	66.5	71.4	74.7	77.2	78.9	81.9	83.8	83.6	80.5	81.0	82.2	83.0	86.1	83.7	86.9	85.4	85.3	88.2	84.0	84.0	87.5					
7-year	64.3	69.8	71.9	76.1	77.6	79.8	83.2	82.2	78.7	80.3	81.9	81.2	85.2	83.5	85.8	85.0	83.4	85.5	82.8	83.4						
8-year	62.4	68.3	70.6	74.9	76.5	78.7	82.5	81.8	77.5	79.2	81.4	80.1	84.3	82.6	83.2	85.0	82.3	84.7	81.8							
9-year	59.9	67.1	69.4	73.9	75.4	77.6	82.5	80.4	76.1	78.9	79.7	79.2	82.9	82.6	82.9	84.9	82.3	83.5								
10-year	58.9	65.8	68.7	73.3	75.4	75.4	81.7	78.3	75.3	78.9	78.3	78.6	82.3	81.5	82.9	84.5	82.3									
11-year	57.5	64.8	67.3	72.7	74.3	74.3	81.1	77.7	75.2	78.0	77.4	76.1	81.8	81.3	82.9	84.3										
12-year	56.4	63.5	66.2	72.0	74.3	72.7	81.0	77.3	74.2	77.4	77.1	75.4	81.0	80.7	82.9											
13-year	55.8	62.8	64.5	71.1	74.3	72.2	78.9	77.1	71.7	76.7	75.9	74.1	79.7	79.6												
14-year	54.4	61.4	63.4	70.7	74.3	70.8	77.6	76.1	71.5	75.7	75.9	74.1	78.9													
15-year	53.2	60.7	62.3	70.1	74.3	70.3	76.5	73.4	70.4	73.3	74.7	73.4														
16-year	52.5	59.7	61.4	69.6	74.3	70.3	76.2	73.3	70.1	72.2	74.0															
17-year	51.2	58.3	60.9	69.0	74.3	69.9	75.3	72.7	68.5	71.6																
18-year	50.2	57.3	60.1	68.3	73.2	69.1	73.6	71.7	68.4																	
19-year	49.6	56.8	59.6	67.5	73.2	68.8	72.4	70.6																		
20-year	48.6	55.5	58.6	67.3	72.8	68.0	71.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 9.11
Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	88.2	89.4	91.8	93.0	93.7	90.0	95.1	94.9	92.9	94.5	95.3	95.2	93.4	93.1	93.4	93.4	94.5	96.1	91.4	93.8	94.8	95.6	96.4	94.4	94.4	96.5
2-year	81.6	83.7	88.3	89.9	89.4	87.4	92.3	89.9	89.4	93.5	92.7	91.6	92.2	90.9	91.1	91.3	93.5	94.2	89.7	92.5	93.3	93.6	95.6	93.7	94.4	
3-year	78.2	79.7	85.0	87.6	88.1	86.3	91.3	87.7	87.5	93.1	90.8	90.2	90.2	88.4	90.7	89.5	92.5	93.0	88.8	91.8	92.2	91.9	94.0	92.0		
4-year	76.3	77.6	83.2	86.0	86.6	84.0	89.8	86.8	85.7	93.1	88.5	87.6	89.6	86.3	90.4	89.5	90.7	92.6	87.3	90.6	90.8	91.2	93.9			
5-year	74.4	75.7	82.1	84.6	84.7	83.6	88.6	85.3	84.4	92.5	87.1	86.7	89.1	86.2	89.1	89.5	90.2	91.8	86.3	88.6	90.8	90.9				
6-year	72.3	73.4	81.0	83.4	81.8	81.5	88.5	84.0	83.9	92.0	86.8	86.6	88.1	85.9	89.1	88.6	89.3	91.4	85.2	87.8	89.8					
7-year	70.8	71.9	79.3	81.7	80.7	80.5	87.7	83.7	83.2	91.1	86.8	85.8	87.8	85.1	89.1	87.6	87.9	91.2	84.9	87.8						
8-year	68.5	70.7	78.4	81.3	80.7	80.2	87.2	83.2	82.6	90.3	86.8	84.7	87.2	84.1	89.0	87.1	86.7	90.7	83.9							
9-year	67.0	69.0	78.1	80.3	79.6	78.9	86.7	82.7	82.2	88.5	86.7	83.4	87.1	83.3	88.5	86.8	86.4	90.1								
10-year	65.5	68.2	77.6	79.3	78.3	78.4	86.6	82.3	81.3	88.5	86.6	82.5	86.9	83.3	87.8	86.0	86.3									
11-year	64.3	67.2	77.2	78.9	77.5	77.6	86.2	81.7	80.8	88.5	86.5	82.2	86.8	83.3	87.0	85.3										
12-year	63.6	66.4	76.8	78.1	77.0	77.5	85.2	81.3	80.8	88.5	86.5	81.2	86.5	83.0	86.7											
13-year	62.5	65.2	76.1	77.6	76.2	77.2	85.1	80.9	80.8	88.5	84.2	81.1	85.5	82.8												
14-year	61.6	64.3	75.2	77.1	76.2	76.9	83.6	80.9	80.8	88.5	84.0	81.1	85.3													
15-year	60.8	63.4	74.1	76.6	76.0	76.9	82.9	80.9	80.8	86.5	82.3	80.6														
16-year	60.1	62.4	73.6	76.1	75.9	76.9	82.6	80.9	78.4	85.6	81.2															
17-year	59.5	61.1	72.9	75.5	75.3	75.9	81.6	80.9	78.4	85.2																
18-year	58.7	60.4	72.5	75.0	74.6	75.7	81.3	80.9	78.2																	
19-year	58.0	59.6	71.6	73.9	74.4	75.6	80.8	80.8																		
20-year	56.8	59.0	71.2	73.0	74.3	74.6	80.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 10.7
Kaposi Sarcoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	94.4	65.4	59.0	59.0	61.8	69.0	72.8	73.8	76.8	74.1	80.5	78.1	75.3	82.6	82.1	81.7	82.8	80.0	82.3	83.5	85.2	80.6	86.6	87.3	83.7	82.2
2-year	89.6	44.5	32.8	29.4	46.8	60.4	65.1	65.5	65.1	66.3	73.4	71.1	72.5	73.5	78.8	79.0	80.4	76.1	79.5	77.7	81.5	80.5	83.9	80.1	79.2	
3-year	88.7	36.1	19.8	18.3	42.9	54.1	62.1	61.1	61.5	60.1	67.6	67.6	66.9	70.9	75.0	76.8	78.8	74.9	76.4	75.0	80.1	79.0	82.5	79.8		
4-year	83.8	32.0	13.8	14.0	38.8	51.6	58.9	59.0	56.7	57.0	65.6	66.4	65.0	69.3	74.2	76.1	76.8	74.9	74.7	73.4	79.2	79.0	78.1			
5-year	83.8	28.0	10.6	11.8	35.7	48.4	56.7	58.5	55.8	54.2	64.1	60.8	64.7	68.2	72.3	74.0	74.9	74.5	74.7	73.3	77.7	77.6				
6-year	76.9	24.5	8.9	10.9	35.1	45.6	54.6	55.7	53.4	52.8	63.8	59.1	64.4	67.0	70.7	73.1	73.5	73.9	74.1	71.7	76.2					
7-year	76.8	23.5	7.7	10.3	33.4	44.9	49.9	53.1	52.8	52.8	63.0	59.1	60.7	66.8	68.7	72.0	72.5	73.9	72.9	71.7						
8-year	76.8	20.8	6.8	9.9	32.2	44.5	47.2	50.4	52.4	52.4	61.6	57.3	58.5	66.7	67.5	70.2	72.5	71.8	69.6							
9-year	76.8	19.2	6.1	9.5	31.0	42.7	44.8	49.9	52.4	50.8	59.6	57.2	56.8	65.5	65.7	68.3	71.8	71.3								
10-year	76.8	17.2	5.9	9.4	29.9	42.7	43.7	49.9	51.5	48.9	56.9	57.2	56.3	63.0	65.1	66.3	68.4									
11-year	74.7	15.9	5.7	9.2	28.8	42.3	42.4	49.2	50.4	46.7	56.9	57.2	55.6	63.0	65.1	63.0										
12-year	74.7	15.6	5.6	9.2	28.0	41.9	41.2	49.2	50.4	46.1	56.1	57.2	54.7	62.5	64.1											
13-year	74.7	14.7	5.2	9.2	27.5	41.9	40.2	47.9	50.4	45.6	53.7	57.2	54.7	60.4												
14-year	74.7	14.7	5.0	8.9	26.9	40.6	40.2	45.6	49.2	45.6	53.7	57.2	54.1													
15-year	74.7	14.7	4.9	8.7	26.3	40.4	40.2	44.5	48.5	45.6	53.7	57.2														
16-year	74.7	14.7	4.8	8.5	25.8	40.1	40.2	42.9	47.9	45.6	53.2															
17-year	74.7	14.4	4.8	8.4	25.5	38.9	38.8	42.2	45.2	43.3																
18-year	74.7	13.2	4.8	8.2	25.4	38.2	37.7	42.1	45.2																	
19-year	74.7	13.2	4.6	8.0	25.4	38.2	37.7	40.5																		
20-year	74.7	13.2	4.5	7.8	24.9	37.9	37.7																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 10.8
Kaposi SarcomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	95.1	63.6	58.7	58.6	60.9	68.8	72.8	73.0	76.9	73.8	79.8	76.9	75.8	81.3	83.0	81.1	82.7	81.2	82.0	83.4	84.1	81.8	87.1	85.8	82.7	83.9
2-year	89.1	41.8	32.1	28.8	45.5	59.8	64.4	64.0	64.8	65.6	73.7	70.2	72.7	72.0	79.4	78.2	80.5	77.2	80.0	77.6	81.0	81.6	84.9	78.3	78.7	
3-year	86.8	33.7	19.0	17.6	41.9	53.8	60.9	59.6	61.2	59.5	68.7	67.4	67.4	69.0	75.9	76.1	78.5	76.0	76.5	74.5	79.2	80.5	83.2	77.4		
4-year	83.7	29.6	13.0	13.3	37.6	51.2	57.3	57.0	56.0	56.7	66.2	66.9	65.2	67.1	75.0	75.6	76.1	76.0	75.1	72.5	78.2	80.5	79.2			
5-year	83.6	25.9	9.7	11.0	35.1	48.1	55.9	56.1	54.9	54.2	64.0	60.7	64.8	65.6	72.9	73.3	73.9	75.4	75.1	72.5	76.9	79.0				
6-year	74.2	22.1	8.0	10.2	34.4	45.4	53.7	54.0	52.0	53.2	64.0	59.0	64.4	63.9	71.1	72.3	72.1	74.7	75.0	71.6	76.6					
7-year	74.1	20.8	6.8	9.6	32.5	44.6	49.5	52.4	51.3	53.2	63.8	59.0	60.3	63.9	68.9	71.1	71.0	74.7	73.5	71.6						
8-year	74.1	18.7	6.0	9.2	31.3	44.2	46.9	50.8	51.2	52.7	62.6	58.1	58.5	63.9	67.6	69.1	71.0	72.4	70.5							
9-year	74.1	17.2	5.4	8.9	30.3	42.3	44.4	50.2	51.2	51.0	60.8	57.8	58.0	63.9	66.1	67.8	70.1	71.9								
10-year	74.1	15.5	5.2	8.7	29.2	42.2	43.1	50.2	50.5	49.6	57.7	57.8	57.5	62.8	65.4	65.6	67.6									
11-year	69.0	14.1	5.0	8.5	28.3	41.9	42.2	49.4	49.2	47.3	57.6	57.8	56.8	62.8	65.4	62.1										
12-year	68.9	13.9	4.9	8.4	27.4	41.4	41.0	49.4	49.2	46.5	56.2	57.8	55.9	62.4	64.4											
13-year	68.9	13.1	4.6	8.4	26.8	41.4	40.0	47.9	48.9	46.0	54.2	57.8	55.9	60.1												
14-year	68.9	13.1	4.4	8.1	26.4	40.4	39.9	45.5	47.1	46.0	54.2	57.8	55.3													
15-year	68.9	13.1	4.4	7.9	25.9	40.2	39.9	44.3	46.1	46.0	54.2	57.1														
16-year	68.9	12.9	4.3	7.8	25.3	39.9	39.9	42.6	45.9	46.0	53.7															
17-year	68.9	12.9	4.3	7.6	25.2	38.7	38.5	41.9	44.1	43.1																
18-year	68.9	12.4	4.3	7.4	25.2	38.0	37.2	41.6	44.1																	
19-year	68.9	12.4	4.1	7.2	25.2	38.0	37.2	40.6																		
20-year	68.9	12.4	4.0	7.0	24.7	37.6	37.2																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 10.9
Kaposi Sarcoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	92.6	90.0	80.5	80.5	87.2	78.4	73.3	85.1	74.0	78.0	86.3	88.8	64.5	100.0	67.3	89.8	83.7	52.1	86.0	81.6	98.3	64.4	76.5	100.0	89.8	64.3
2-year	90.4	84.0	79.1	63.7	85.9	78.4	73.3	85.1	67.4	76.5	67.5	78.2	64.5	97.2	67.3	89.8	78.0	52.1	71.8	76.9	86.1	64.4	68.4	97.7	81.5	
3-year	90.4	73.4	74.3	57.8	74.5	64.5	73.3	85.0	63.3	67.3	52.1	66.4	55.9	97.2	60.9	82.8	78.0	52.1	71.8	76.9	86.1	59.3	68.4	97.7		
4-year	83.5	69.5	73.6	54.5	74.5	64.5	73.3	85.0	63.3	61.4	52.1	54.7	55.9	97.2	60.9	82.8	78.0	52.1	67.6	76.9	86.1	59.3	63.4			
5-year	81.4	61.0	73.6	53.3	53.7	55.2	66.8	85.0	63.3	52.5	52.1	54.7	55.9	97.2	60.9	82.8	78.0	52.1	59.4	76.9	80.7	54.2				
6-year	81.4	61.0	73.6	49.7	53.7	49.3	66.8	81.4	63.3	43.3	52.1	54.7	55.9	97.2	60.9	82.8	78.0	52.1	59.4	70.2	61.3					
7-year	81.4	61.0	73.6	49.7	53.7	49.3	54.6	54.4	63.3	43.3	50.9	54.7	55.9	97.2	60.9	82.8	78.0	52.1	59.4	70.2						
8-year	81.4	54.1	66.7	49.7	53.7	49.3	47.6	19.5	63.3	43.3	45.9	43.8	55.9	77.2	60.9	82.8	78.0	52.1	56.1							
9-year	81.4	51.0	66.1	48.3	50.7	49.3	47.6	19.5	63.3	43.3	38.0	43.8	21.2	77.2	58.6	75.5	78.0	52.1								
10-year	81.4	45.4	59.8	48.3	48.1	49.3	47.6	19.5	63.0	35.5	38.0	43.8	21.2	51.4	58.6	75.5	74.8									
11-year	81.4	42.7	59.8	48.3	40.3	49.3	41.0	19.5	63.0	35.5	38.0	43.8	21.2	51.4	58.6	75.5										
12-year	81.4	41.9	57.0	48.3	40.3	49.3	41.0	19.5	63.0	35.5	38.0	43.8	21.2	51.4	58.6											
13-year	81.4	41.0	52.4	48.3	40.3	38.4	41.0	19.5	63.0	35.5	0.0	43.8	21.2	51.4												
14-year	81.4	41.0	39.3	48.3	39.0	38.4	41.0	19.5	63.0	35.5	0.0	43.8	21.2													
15-year	81.4	41.0	33.4	48.3	27.5	38.4	41.0	19.5	63.0	35.5	0.0	43.8														
16-year	81.4	41.0	33.4	48.3	27.5	38.4	41.0	19.5	63.0	35.5	0.0															
17-year	81.4	29.3	33.4	48.3	15.6	38.4	41.0	19.5	48.3	35.5																
18-year	80.2	16.6	33.4	48.3	0.0	38.4	41.0	19.5	48.3																	
19-year	80.2	16.6	33.4	48.3	0.0	38.4	41.0	0.0																		
20-year	80.2	16.6	33.4	48.3	0.0	38.4	41.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 11.9
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	68.8	70.4	72.9	76.4	77.0	77.0	76.3	77.7	78.0	79.7	81.2	79.6	81.8	85.0	83.9	85.0	86.4	85.2	86.5	86.4	85.9	87.3	87.8	86.4	87.1	88.2
2-year	60.3	61.7	65.0	69.2	69.7	70.5	70.0	71.2	71.8	73.8	73.9	74.7	76.2	79.6	80.1	81.8	82.5	81.2	82.2	81.6	81.0	83.4	83.6	82.1	83.6	
3-year	56.3	57.0	60.8	65.8	65.5	67.3	66.1	67.5	67.6	69.3	70.8	71.3	73.4	76.4	77.7	78.5	80.3	78.9	79.4	79.3	77.9	80.7	81.3	79.6		
4-year	53.0	53.9	57.8	63.2	63.3	64.3	63.7	64.5	64.5	67.5	68.1	69.2	71.1	73.5	76.0	75.9	77.7	76.4	77.6	77.0	75.1	78.9	78.9			
5-year	50.9	52.0	55.8	60.7	61.4	62.5	61.7	63.3	62.5	65.4	66.4	67.5	69.0	71.1	74.4	73.8	75.1	74.4	76.2	74.8	73.3	77.2				
6-year	49.3	49.7	53.8	59.1	59.4	60.5	59.7	61.2	60.5	63.8	64.9	65.5	67.9	69.1	72.8	72.5	73.7	73.2	74.4	73.9	72.1					
7-year	47.5	48.6	52.3	57.5	58.0	59.0	58.1	59.5	59.1	62.2	63.1	63.6	66.2	67.8	71.9	70.9	72.6	71.7	73.2	72.0						
8-year	46.2	47.4	51.1	56.1	56.8	58.6	56.0	58.0	57.3	60.2	61.9	61.8	64.8	65.8	70.9	69.5	71.0	70.6	72.0							
9-year	45.0	46.0	49.6	55.0	55.3	57.8	54.1	56.6	55.8	58.9	60.8	60.4	63.4	64.1	69.8	68.3	70.0	69.1								
10-year	44.0	44.9	48.0	53.3	54.3	55.9	53.0	55.5	53.9	56.8	59.5	59.4	62.3	62.5	69.0	67.5	68.5									
11-year	43.1	43.6	46.5	51.8	52.3	55.0	52.1	55.0	52.8	55.8	57.6	58.5	60.9	61.4	66.9	66.3										
12-year	42.2	42.8	45.3	50.7	50.7	53.5	51.4	54.3	52.0	55.1	56.2	57.7	59.5	60.3	66.1											
13-year	41.5	42.0	44.3	49.5	49.2	51.8	50.4	53.4	51.4	54.1	54.4	56.2	58.0	59.1												
14-year	40.5	41.4	43.0	48.2	48.2	50.3	49.8	52.2	50.1	52.7	53.6	55.9	56.8													
15-year	39.5	41.0	42.1	47.5	47.2	49.0	48.9	50.7	48.8	52.2	53.0	54.9														
16-year	39.2	40.4	41.1	46.8	46.9	48.1	47.6	49.6	47.8	51.2	52.0															
17-year	38.7	39.8	40.2	46.0	45.7	47.9	46.3	48.3	46.8	49.6																
18-year	37.6	39.3	39.3	45.0	44.6	47.3	45.2	47.5	46.4																	
19-year	36.7	38.3	38.5	44.3	44.0	46.3	44.7	46.5																		
20-year	36.1	37.3	37.9	43.4	43.1	45.0	43.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 11.10
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	69.4	70.5	73.4	77.4	79.2	77.9	77.9	78.2	77.5	79.1	80.6	80.2	80.6	85.5	84.1	84.9	86.4	85.1	86.3	86.4	86.1	87.2	87.8	85.8	87.4	88.3
2-year	60.7	61.9	65.5	69.5	71.3	71.7	71.0	72.1	70.9	72.8	73.5	75.0	75.8	80.4	80.7	81.4	82.4	80.6	81.9	80.8	81.0	83.1	83.8	81.3	84.0	
3-year	56.4	57.4	61.4	66.3	67.2	68.0	66.8	68.0	66.8	67.9	70.3	71.0	73.5	77.2	78.0	77.9	79.7	78.1	78.9	78.2	77.9	80.1	81.3	78.7		
4-year	52.9	54.4	58.3	63.5	64.4	64.9	64.2	65.0	63.5	65.8	67.5	68.4	70.7	74.8	75.8	74.9	76.7	75.5	77.2	75.9	74.9	78.0	78.6			
5-year	51.1	52.4	56.2	60.9	62.4	63.1	61.7	63.7	61.5	62.9	65.7	67.2	69.1	71.9	74.1	72.8	73.7	73.2	75.9	73.4	72.8	76.5				
6-year	49.5	50.2	53.9	59.0	59.5	61.7	59.4	61.3	58.8	61.6	63.6	65.1	68.1	69.5	71.8	71.0	72.6	72.1	74.0	72.2	71.6					
7-year	47.3	48.9	52.1	57.4	58.5	60.1	57.6	59.3	57.7	59.7	61.0	63.3	66.3	67.8	70.7	69.1	71.8	70.9	72.7	70.3						
8-year	45.8	47.8	50.8	56.1	57.2	59.2	54.5	57.3	55.7	57.7	59.7	61.5	64.6	66.0	69.4	67.5	69.8	69.8	71.6							
9-year	44.8	46.3	48.8	55.2	56.0	58.6	52.6	56.3	53.8	56.0	58.8	60.4	63.1	64.0	68.6	67.0	69.0	68.7								
10-year	43.4	45.3	47.5	53.5	54.7	56.2	52.1	55.2	51.6	54.3	57.4	59.2	62.1	62.5	68.2	66.0	67.1									
11-year	42.4	44.1	45.6	52.0	53.3	55.5	50.6	54.5	50.2	53.2	55.8	58.3	60.5	61.2	66.2	65.1										
12-year	41.3	43.3	44.1	50.9	51.0	53.7	49.4	53.3	49.3	52.9	54.6	57.2	58.9	60.6	65.4											
13-year	40.5	42.6	42.8	49.6	49.2	52.3	48.8	53.0	49.1	52.1	53.2	55.4	57.2	59.2												
14-year	39.5	41.6	41.7	48.5	49.1	50.6	48.3	52.0	48.1	50.3	52.6	55.1	56.2													
15-year	38.4	41.3	40.8	47.8	47.7	48.6	46.7	50.2	46.9	50.3	51.7	54.9														
16-year	38.0	40.4	39.6	47.2	47.7	47.3	45.3	48.9	46.1	48.9	50.6															
17-year	37.4	40.1	38.8	46.4	46.3	47.1	43.3	47.9	44.5	47.6																
18-year	36.5	39.6	37.7	45.3	44.9	47.1	41.7	47.2	44.0																	
19-year	35.6	38.7	36.9	44.5	44.2	46.4	40.8	46.7																		
20-year	34.7	37.5	36.3	43.7	43.4	44.8	40.1																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 11.11
Cancer of the Kidney and Renal Pelvis (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	67.9	70.2	72.2	74.6	73.5	75.7	73.9	76.9	78.7	80.5	82.3	78.7	83.7	84.2	83.4	85.3	86.5	85.4	86.7	86.4	85.6	87.6	87.7	87.3	86.4	88.0
2-year	59.6	61.4	64.2	68.6	67.1	68.5	68.6	69.7	73.3	75.4	74.6	74.2	76.8	78.4	79.1	82.3	82.7	82.2	82.8	82.9	80.8	83.9	83.3	83.5	82.9	
3-year	56.2	56.3	59.8	65.1	62.8	66.2	65.0	66.6	68.9	71.6	71.8	71.9	73.2	75.3	77.3	79.4	81.4	80.3	80.3	81.0	77.9	81.5	81.3	81.1		
4-year	53.1	53.2	57.1	62.7	61.6	63.4	62.8	63.8	66.1	70.1	69.0	70.4	71.7	71.5	76.2	77.5	79.4	77.8	78.1	79.0	75.6	80.6	79.4			
5-year	50.6	51.3	55.0	60.5	59.7	61.6	61.4	62.6	64.1	69.5	67.7	68.1	68.7	69.9	75.0	75.2	77.5	76.4	76.8	77.2	74.2	78.5				
6-year	49.0	49.0	53.7	59.1	58.8	58.6	60.2	61.1	63.0	67.5	66.9	66.0	67.5	68.6	74.5	74.6	75.6	74.8	75.0	76.7	73.1					
7-year	47.7	48.0	52.5	57.5	57.2	57.2	58.8	59.7	61.2	66.3	66.3	63.9	66.2	67.8	73.8	73.7	73.9	73.0	74.0	74.5						
8-year	46.9	46.8	51.6	56.1	56.2	57.2	57.9	58.8	59.9	64.2	65.5	62.3	65.2	65.4	73.5	72.3	72.9	71.7	72.7							
9-year	45.3	45.5	50.7	54.7	54.1	56.6	56.2	57.1	58.9	63.5	63.9	60.3	63.9	64.0	71.6	70.1	71.7	69.8								
10-year	44.9	44.3	48.7	52.9	53.6	55.4	54.4	56.0	57.6	61.0	62.7	59.5	62.6	62.5	70.0	69.7	70.8									
11-year	44.2	42.8	48.0	51.5	50.7	54.1	54.2	55.5	56.8	60.0	60.4	58.7	61.3	61.5	68.1	67.8										
12-year	43.6	42.0	47.2	50.3	50.1	52.9	54.2	55.5	56.1	58.5	58.8	58.5	60.4	59.7	67.0											
13-year	43.0	41.1	46.5	49.2	48.9	51.0	52.8	54.0	54.9	57.2	56.2	57.4	59.2	59.0												
14-year	42.1	40.7	45.1	47.8	46.7	49.7	51.9	52.4	53.2	56.6	55.3	56.6	57.9													
15-year	41.2	40.3	43.9	47.0	46.0	49.3	51.9	51.3	51.7	55.3	55.0	54.8														
16-year	41.1	40.2	43.3	46.1	45.0	49.1	50.9	50.4	50.5	54.9	54.1															
17-year	40.4	39.3	42.3	45.5	44.6	49.0	50.8	48.9	50.3	52.9																
18-year	39.4	38.7	41.6	44.6	44.1	47.6	50.2	47.9	50.1																	
19-year	38.4	37.6	40.9	43.9	43.6	46.1	50.1	45.9																		
20-year	38.2	37.0	40.2	43.0	42.4	45.4	48.3																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 12.9
Cancer of the Larynx (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	88.3	88.4	88.2	87.7	89.2	88.1	85.8	86.1	85.5	85.3	85.3	84.0	84.0	85.4	86.6	86.4	88.2	84.2	86.9	85.7	86.9	87.6	84.2	85.6	85.5	84.8
2-year	79.1	79.5	79.0	78.0	80.0	78.9	77.3	76.1	74.8	76.8	75.1	74.8	74.6	75.2	78.1	74.7	75.6	76.2	76.7	77.2	73.0	78.6	76.1	77.7	77.5	
3-year	73.3	74.1	73.6	71.8	74.9	74.1	72.3	70.8	70.7	71.0	70.0	70.0	66.7	68.6	72.6	68.4	70.6	71.2	71.7	70.5	67.4	72.9	69.3	71.0		
4-year	68.9	70.0	68.8	67.4	69.3	69.7	68.8	67.0	68.6	66.7	66.6	67.0	62.6	64.4	70.0	62.7	65.9	67.0	68.1	65.5	64.0	66.6	64.7			
5-year	65.5	66.7	65.8	64.5	66.1	65.8	65.2	63.4	63.8	63.3	62.9	62.7	60.3	62.4	65.7	59.4	62.1	64.5	64.1	60.7	61.6	63.2				
6-year	63.1	63.9	62.5	61.0	63.3	63.4	62.3	59.1	60.5	60.4	60.5	60.1	56.0	59.8	61.0	56.8	59.5	61.1	61.4	58.6	57.8					
7-year	60.8	61.3	60.1	59.2	60.0	61.2	59.7	55.5	57.2	56.5	57.0	56.4	54.2	57.0	58.4	53.7	55.8	58.1	58.0	56.1						
8-year	58.8	59.7	57.2	56.5	57.4	57.9	57.0	52.7	54.1	53.1	54.6	53.1	52.0	53.9	54.8	51.2	53.4	54.2	55.1							
9-year	56.4	57.3	54.6	53.9	54.3	54.0	55.5	50.3	52.1	50.8	52.0	52.2	49.8	52.0	51.8	48.3	51.5	50.9								
10-year	54.2	55.4	52.3	51.4	51.3	51.8	53.5	48.3	50.5	48.9	50.4	48.9	47.6	51.0	48.8	45.6	50.1									
11-year	52.7	53.0	50.2	49.4	47.8	49.1	50.9	47.0	47.7	47.6	47.0	45.9	44.8	48.6	46.3	42.4										
12-year	51.0	50.3	48.2	47.4	46.1	46.5	46.3	45.0	45.0	44.4	45.2	42.4	43.1	46.0	43.3											
13-year	49.7	48.0	47.0	45.0	44.3	44.5	45.1	43.2	42.1	43.0	43.3	39.4	41.9	42.9												
14-year	48.0	46.5	44.3	43.1	41.5	41.5	43.2	41.6	41.1	40.4	41.2	37.9	39.7													
15-year	45.8	44.8	41.9	41.4	39.0	38.8	42.3	39.6	39.2	38.6	39.3	36.2														
16-year	43.7	42.1	40.3	39.5	38.0	37.6	41.1	39.0	37.1	36.2	37.3															
17-year	41.4	40.5	38.9	37.6	37.5	35.5	38.4	36.0	34.0	35.8																
18-year	40.2	38.8	37.4	35.5	36.4	34.0	37.3	33.2	33.1																	
19-year	38.4	37.3	34.8	33.8	35.8	31.2	36.9	32.8																		
20-year	36.5	35.6	33.6	31.3	35.4	28.8	36.3																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 12.10
Cancer of the Larynx (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	88.6	88.5	88.5	88.4	91.2	88.0	88.3	87.1	86.9	85.4	85.7	84.6	84.1	85.9	86.3	86.7	88.9	84.3	87.7	86.2	87.1	86.6	84.6	86.9	86.1	83.2
2-year	79.2	79.7	79.4	79.1	81.8	79.1	80.0	76.9	76.1	77.4	75.2	75.0	75.2	75.3	77.1	75.6	75.7	76.8	77.1	77.9	73.0	78.6	77.1	79.0	79.0	
3-year	73.6	74.5	74.2	72.6	76.6	74.2	75.0	71.6	71.6	71.7	70.9	70.9	68.3	68.7	72.6	69.4	70.2	72.9	71.6	71.4	67.3	73.1	71.6	72.8		
4-year	69.2	70.6	69.6	68.4	71.5	69.8	71.2	68.3	69.6	67.3	66.8	68.2	63.2	63.9	70.1	64.0	65.9	68.2	67.6	66.2	63.1	67.3	67.1			
5-year	65.8	67.3	66.6	65.5	68.2	66.3	67.8	65.4	65.3	64.2	63.1	63.5	60.5	62.1	65.2	61.4	61.6	65.6	64.1	62.4	60.0	63.8				
6-year	63.4	64.4	63.3	62.1	65.6	63.7	65.0	62.2	62.0	61.6	60.8	61.2	56.2	60.4	60.3	58.4	59.1	62.5	61.8	60.1	56.3					
7-year	61.3	61.8	61.1	60.4	62.0	61.3	62.9	58.7	58.6	58.5	58.5	57.1	54.5	58.6	57.2	55.2	55.9	59.8	58.3	57.3						
8-year	59.1	60.4	58.4	57.9	59.4	57.9	59.7	55.6	55.4	55.9	56.1	54.2	53.2	55.8	53.0	52.1	53.8	55.6	55.3							
9-year	56.8	57.9	56.2	55.3	56.6	53.6	57.7	53.0	53.4	53.9	54.4	53.0	50.9	53.9	50.3	49.4	52.0	53.0								
10-year	54.8	56.3	54.0	52.6	53.7	51.3	55.9	50.8	51.4	51.8	53.0	49.7	49.0	53.7	48.1	46.9	50.6									
11-year	53.5	53.8	52.2	50.6	50.0	48.8	53.8	49.9	48.8	49.8	49.1	47.0	45.9	51.1	45.4	43.4										
12-year	52.0	51.0	50.3	48.7	49.2	46.0	48.8	48.6	46.0	47.0	47.4	43.7	44.0	48.6	42.4											
13-year	50.8	48.8	49.3	46.5	47.3	43.8	47.0	46.5	43.8	46.2	45.5	41.2	43.1	44.9												
14-year	49.1	47.0	46.9	44.6	44.9	41.7	44.7	45.5	42.5	43.9	43.7	39.6	40.8													
15-year	46.9	45.2	44.4	42.8	42.1	39.2	44.4	43.6	41.1	41.7	42.1	37.4														
16-year	44.7	42.9	42.9	40.9	40.9	37.7	43.4	43.4	38.3	39.1	40.0															
17-year	42.6	41.7	41.5	39.2	40.2	35.4	39.7	40.0	36.5	38.9																
18-year	41.2	39.8	39.6	37.1	39.9	34.2	38.7	36.9	36.1																	
19-year	39.2	38.2	36.8	35.1	39.9	31.8	38.7	36.4																		
20-year	37.6	36.4	35.8	32.6	39.9	29.5	38.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 12.11
Cancer of the Larynx (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	86.8	88.1	86.7	85.0	80.7	88.8	76.8	83.1	81.0	84.7	83.2	81.8	83.8	83.4	87.6	85.0	84.8	83.5	83.0	83.6	86.0	91.5	82.1	80.1	82.5	91.8
2-year	78.6	78.6	77.6	73.6	72.0	78.4	67.3	73.4	70.5	74.3	74.3	73.5	71.8	74.7	81.0	70.3	74.9	73.9	74.5	74.3	73.1	78.7	71.9	72.4	71.2	
3-year	71.7	72.0	71.2	68.4	67.5	73.4	62.3	68.3	66.9	68.0	66.0	66.1	59.9	67.9	72.5	63.7	72.0	64.0	72.2	66.5	67.7	72.3	58.9	63.5		
4-year	67.0	67.5	65.6	63.3	60.1	69.4	59.4	62.9	64.8	63.9	65.5	61.8	59.6	65.6	69.6	56.7	65.6	61.8	70.7	61.9	66.9	63.3	54.1			
5-year	63.8	64.2	62.6	60.1	57.4	63.6	55.7	57.1	58.6	59.7	61.7	59.4	58.5	63.0	67.2	50.2	63.3	59.2	64.4	53.4	66.9	60.4				
6-year	61.3	61.3	59.4	56.7	53.8	62.3	52.1	49.6	55.2	55.8	59.1	55.6	55.4	57.2	63.8	49.4	61.0	54.9	59.6	51.9	64.0					
7-year	58.4	59.1	56.1	54.3	51.4	59.9	48.4	45.6	52.5	48.3	50.2	53.2	52.7	51.4	62.3	46.3	55.2	50.8	56.8	50.1						
8-year	57.1	56.6	52.5	50.9	49.1	57.8	47.1	44.0	49.2	41.7	47.3	48.5	46.8	47.5	61.2	46.3	51.7	48.4	53.8							
9-year	54.6	54.4	48.2	48.5	45.1	55.0	47.1	41.8	47.7	38.2	41.3	48.5	44.7	45.4	57.1	43.1	49.4	42.3								
10-year	51.2	51.7	45.5	46.5	41.6	53.0	44.6	40.5	47.0	37.4	38.3	45.3	41.1	40.8	51.7	39.5	47.8									
11-year	48.4	49.3	42.4	44.4	38.8	50.1	40.2	37.6	43.7	37.4	37.5	41.1	39.9	39.9	49.5	37.5										
12-year	45.7	46.9	39.9	42.4	34.0	48.4	37.3	34.3	41.4	34.0	35.3	37.0	38.7	36.8	46.4											
13-year	44.1	44.4	38.1	39.1	32.4	46.0	37.3	33.3	36.6	30.7	33.0	32.3	36.3	35.7												
14-year	42.8	43.4	34.4	37.6	28.2	40.7	36.9	29.7	36.2	27.1	29.5	31.3	34.5													
15-year	40.5	42.8	32.7	36.1	27.1	37.3	34.8	27.9	32.8	26.2	27.0	30.5														
16-year	38.6	38.9	30.9	34.0	27.0	36.9	33.2	26.4	32.7	24.7	25.2															
17-year	36.1	35.6	29.4	31.6	26.3	35.3	33.1	23.1	26.0	23.9																
18-year	35.3	34.6	29.0	29.6	22.7	33.3	31.8	22.4	22.4																	
19-year	34.2	33.1	27.4	28.7	19.2	29.1	29.5	22.2																		
20-year	31.8	31.8	25.3	26.4	18.5	26.6	28.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 13.17
LeukemiaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	60.7	62.6	64.8	66.8	68.2	68.2	67.8	66.7	66.0	66.9	70.6	72.2	73.4	75.0	74.1	76.5	77.0	76.7	77.4	78.3	79.6	79.2	79.0	79.2	81.0	79.1
2-year	50.1	52.5	55.2	57.5	58.9	59.9	59.7	58.1	58.5	59.0	63.1	65.0	66.3	67.7	68.4	70.7	70.7	70.4	71.0	72.7	73.5	72.8	73.4	73.8	75.5	
3-year	43.8	46.5	49.6	52.9	54.3	55.4	55.0	54.1	54.8	55.1	59.1	62.0	62.4	64.2	64.9	67.5	67.6	66.6	68.6	69.6	70.6	69.5	70.3	70.5		
4-year	38.9	41.7	45.5	49.1	50.7	51.9	51.0	50.4	51.7	52.0	56.8	59.7	59.6	61.9	62.8	64.8	65.4	64.1	66.1	67.9	68.7	67.8	67.6			
5-year	34.6	37.6	42.3	46.1	48.1	48.7	48.3	48.2	48.5	49.9	54.4	57.7	57.5	59.6	61.0	62.7	63.8	62.1	63.7	66.3	66.9	66.1				
6-year	31.4	35.0	39.8	43.6	45.7	46.5	45.8	46.2	46.3	48.6	52.4	56.1	55.7	57.2	59.4	60.8	62.3	60.6	61.5	65.2	66.0					
7-year	29.1	32.5	37.2	41.0	44.3	45.1	43.9	44.7	44.5	47.3	51.1	54.5	53.8	56.1	58.2	59.3	61.6	58.8	60.4	64.1						
8-year	27.0	30.4	35.0	38.8	41.7	43.9	43.3	43.2	43.0	45.8	49.7	53.6	52.0	54.7	56.3	57.4	60.5	57.8	58.6							
9-year	25.0	28.5	33.0	37.3	40.6	42.1	41.9	42.2	42.2	44.9	48.3	52.0	50.9	53.1	54.7	56.2	59.5	56.8								
10-year	23.7	26.9	31.6	36.1	39.2	40.7	40.7	41.1	40.7	43.9	47.1	50.6	49.1	51.5	53.4	54.6	58.5									
11-year	22.4	25.7	29.9	34.7	37.8	39.8	38.9	39.8	39.3	42.7	46.0	49.3	47.5	49.9	52.9	53.5										
12-year	21.2	24.6	28.8	33.6	37.1	38.7	37.8	39.1	38.4	42.1	45.0	48.5	46.3	49.3	52.3											
13-year	20.5	23.6	28.0	32.7	36.7	38.3	37.1	38.1	37.3	40.5	44.2	46.8	45.7	48.0												
14-year	19.8	22.9	27.3	31.9	35.6	37.0	36.9	37.1	36.8	39.3	43.4	45.9	44.9													
15-year	19.2	22.3	26.7	31.4	35.3	36.2	36.4	36.0	36.1	38.5	42.4	44.9														
16-year	18.5	21.4	26.1	30.7	35.0	35.9	36.3	35.1	35.5	37.7	41.9															
17-year	18.1	20.7	25.6	30.0	34.0	35.5	35.3	34.7	34.8	37.4																
18-year	17.7	20.1	25.3	29.5	33.0	35.1	34.7	34.5	34.6																	
19-year	17.2	19.5	24.8	29.2	32.6	34.3	34.5	34.5																		
20-year	16.7	19.1	24.5	28.7	32.0	33.8	33.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 13.18
LeukemiaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Survival Time																											
1-year	60.3	62.8	65.5	67.8	69.3	69.7	68.7	66.6	66.9	68.8	72.7	73.0	74.4	76.6	76.0	77.0	78.1	77.6	79.0	79.4	80.8	79.9	80.2	81.3	82.0	79.8	
2-year	49.6	52.4	56.1	58.3	60.4	61.6	61.0	58.6	59.2	60.8	64.8	65.2	66.9	68.6	70.1	70.8	71.9	71.6	72.8	74.2	74.9	73.5	75.1	75.7	76.6		
3-year	42.7	46.0	49.9	53.3	55.6	57.3	55.3	54.6	55.3	57.2	61.0	62.5	62.8	65.3	66.5	67.5	69.0	67.9	70.1	71.6	72.2	70.2	71.8	72.4			
4-year	37.9	41.2	46.0	49.5	51.7	53.6	50.7	50.6	52.0	53.8	58.1	59.8	60.2	62.7	65.0	64.9	66.8	65.1	68.0	69.3	70.0	68.2	68.9				
5-year	33.4	37.0	42.9	46.2	48.2	49.6	47.9	48.5	48.1	52.3	55.3	57.1	57.8	60.1	62.9	62.7	65.3	63.3	65.9	67.7	68.7	66.4					
6-year	30.3	34.3	40.3	43.6	45.2	47.1	45.9	46.2	46.2	50.8	52.9	55.5	55.7	57.5	60.9	60.4	63.6	61.6	63.3	66.6	67.4						
7-year	27.7	31.8	37.4	40.9	44.0	45.7	43.9	44.4	44.1	49.8	51.5	53.1	53.6	56.5	59.6	59.1	62.7	59.6	62.1	65.9							
8-year	25.4	29.4	35.4	38.6	41.8	44.6	43.1	43.1	42.4	47.8	50.0	52.1	51.4	55.0	57.2	56.5	61.6	58.5	60.4								
9-year	23.3	27.4	33.3	37.1	40.5	42.4	42.1	42.0	41.6	46.3	48.6	50.8	50.3	52.8	55.3	54.9	60.6	57.1									
10-year	21.9	26.0	31.7	35.9	38.5	40.8	41.3	41.0	40.2	45.4	47.1	49.1	48.2	51.1	54.0	52.8	59.7										
11-year	20.5	24.9	30.2	34.7	37.7	39.8	39.3	39.3	38.5	44.4	46.1	47.5	46.5	49.7	53.3	51.5											
12-year	19.4	24.3	29.1	33.5	37.1	39.1	38.1	38.4	37.6	43.7	45.2	46.9	45.7	49.2	52.7												
13-year	18.6	23.4	28.3	32.5	36.7	39.0	37.4	37.0	36.4	41.9	44.1	45.7	45.0	48.0													
14-year	17.9	22.7	27.7	31.6	35.7	37.8	37.0	35.8	35.9	41.0	43.3	44.9	44.6														
15-year	17.5	22.1	27.0	31.0	35.2	37.2	36.5	35.0	35.3	39.9	42.2	43.7															
16-year	16.8	21.3	26.3	30.4	34.7	37.0	36.2	34.2	34.8	38.8	41.7																
17-year	16.4	20.4	26.0	29.7	33.9	36.7	34.9	33.6	34.6	38.7																	
18-year	16.0	19.9	25.7	29.1	32.6	36.5	34.5	33.2	34.4																		
19-year	15.5	19.2	25.4	28.9	32.2	35.6	34.2	33.2																			
20-year	14.7	18.7	24.9	28.5	31.7	34.8	33.6																				

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 13.19
LeukemiaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	61.2	62.3	63.8	65.4	66.6	66.3	66.5	66.8	64.6	64.4	67.6	71.1	72.0	72.8	71.5	75.8	75.5	75.4	75.1	76.7	78.0	78.3	77.3	76.4	79.6	77.9
2-year	50.9	52.6	54.1	56.4	56.8	57.8	58.1	57.5	57.5	56.5	60.7	64.7	65.4	66.4	66.1	70.5	69.0	68.6	68.5	70.7	71.7	71.9	71.1	71.2	74.1	
3-year	45.1	47.1	49.1	52.3	52.5	52.9	54.6	53.4	54.0	52.3	56.6	61.4	61.9	62.8	62.5	67.5	65.8	64.9	66.4	66.9	68.4	68.7	68.1	68.0		
4-year	40.2	42.5	44.9	48.6	49.4	49.8	51.5	50.0	51.2	49.6	55.0	59.6	58.8	60.8	59.8	64.7	63.4	62.7	63.4	66.1	66.8	67.3	65.8			
5-year	36.1	38.4	41.4	45.8	48.0	47.5	48.9	47.9	49.0	46.8	53.1	58.4	57.1	58.8	58.4	62.8	61.7	60.4	60.7	64.4	64.4	65.8				
6-year	32.9	36.0	39.1	43.6	46.3	45.7	45.7	46.1	46.3	45.8	51.7	56.9	55.7	56.7	57.4	61.3	60.4	59.1	59.0	63.3	63.9					
7-year	31.0	33.5	36.8	41.1	44.7	44.2	43.9	45.0	45.1	44.0	50.5	56.3	54.0	55.7	56.1	59.5	60.0	57.9	58.1	61.5						
8-year	29.1	31.6	34.4	39.1	41.6	42.8	43.4	43.4	43.8	43.1	49.1	55.7	52.7	54.2	55.0	58.5	58.5	56.7	56.3							
9-year	27.3	29.9	32.6	37.5	40.9	41.6	41.4	42.5	42.9	43.1	47.9	53.7	51.7	53.4	53.8	57.7	57.9	56.3								
10-year	26.0	28.2	31.5	36.3	39.9	40.5	39.8	41.0	41.4	41.8	47.0	52.5	50.3	52.0	52.6	56.8	57.0									
11-year	24.8	26.9	29.5	34.7	37.8	39.7	38.4	40.4	40.4	40.4	45.7	51.8	48.7	50.3	52.3	56.0										
12-year	23.5	25.1	28.5	33.8	37.2	38.2	37.4	39.9	39.4	39.8	44.7	50.7	47.0	49.4	51.6											
13-year	22.8	23.9	27.5	32.9	36.5	37.4	36.7	39.3	38.5	38.6	44.3	48.3	46.5	48.0												
14-year	22.1	23.2	26.7	32.3	35.5	36.0	36.4	38.7	38.1	37.0	43.4	47.2	45.3													
15-year	21.5	22.5	26.4	31.8	35.3	34.8	36.1	37.1	36.9	36.7	42.7	46.6														
16-year	20.7	21.7	25.8	31.0	35.3	34.4	36.0	36.3	36.4	36.2	42.0															
17-year	20.2	21.0	25.1	30.4	34.2	34.0	35.8	36.1	35.1	35.5																
18-year	19.8	20.5	24.8	30.1	33.6	33.4	34.8	36.1	34.8																	
19-year	19.3	19.9	24.1	29.6	33.0	32.6	34.8	36.1																		
20-year	19.1	19.6	23.8	29.1	32.2	32.3	33.5																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 14.9
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	13.7	14.9	17.8	21.4	23.4	24.7	27.7	26.8	27.7	29.0	31.2	32.9	36.3	38.3	39.1	39.2	42.1	44.8	44.6	48.4	47.0	46.5	49.0	49.1	52.5	52.5
2-year	6.6	8.3	9.8	12.4	13.7	15.0	17.4	17.0	17.8	19.9	21.7	22.3	24.8	26.3	27.0	28.8	30.3	32.5	31.2	36.1	33.5	32.4	35.3	36.8	39.3	
3-year	4.8	6.1	7.3	8.7	9.5	10.9	12.5	13.1	14.3	15.4	16.0	17.6	20.5	20.7	21.2	23.4	24.0	25.8	24.3	27.9	26.3	24.8	27.3	29.5		
4-year	4.1	4.6	6.3	6.9	6.9	9.2	10.1	10.7	12.0	12.4	13.8	15.5	17.1	17.6	18.4	19.9	21.0	21.6	19.8	22.6	22.7	20.5	23.2			
5-year	3.4	3.8	5.3	5.6	5.7	8.2	8.2	9.4	10.1	11.7	12.0	14.1	15.6	14.7	16.8	17.1	18.5	18.7	17.0	19.4	20.3	18.0				
6-year	3.0	3.6	5.0	5.0	4.9	7.1	7.4	8.4	9.1	10.8	11.0	12.0	15.0	13.3	15.4	15.6	16.5	16.9	15.3	17.3	18.6					
7-year	2.8	3.1	4.8	4.6	4.4	6.7	6.7	7.0	8.3	10.0	9.9	11.4	13.3	12.7	14.1	14.6	14.6	15.6	14.5	15.8						
8-year	2.7	2.9	4.4	4.4	4.2	6.5	5.9	6.6	7.8	9.6	8.9	10.4	12.3	12.1	13.5	13.7	13.7	14.3	13.1							
9-year	2.5	2.9	4.2	4.3	4.1	6.0	5.7	5.6	7.5	9.4	8.5	10.1	11.7	11.6	12.4	13.0	13.6									
10-year	2.5	2.9	4.0	4.2	3.6	5.7	5.5	5.5	7.4	9.0	7.9	9.3	11.2	10.8	11.7	12.6	12.0									
11-year	2.5	2.7	4.0	4.0	3.2	5.6	5.5	4.8	6.9	8.7	7.8	8.7	10.9	10.3	11.6	12.1										
12-year	2.5	2.7	3.8	3.9	3.1	5.3	5.5	4.6	6.8	8.6	7.6	8.1	10.6	9.9	11.6											
13-year	2.3	2.5	3.8	3.8	3.1	5.1	5.5	4.5	6.6	8.4	7.5	7.6	10.2	9.7												
14-year	2.3	2.5	3.7	3.7	3.1	4.9	5.5	4.5	6.1	8.4	7.2	7.4	10.0													
15-year	2.2	2.5	3.7	3.7	3.0	4.1	5.1	4.5	5.9	8.3	6.8	7.2														
16-year	2.2	2.4	3.7	3.6	3.0	4.0	4.8	4.5	5.4	8.2	6.6															
17-year	2.1	2.4	3.7	3.5	2.9	4.0	4.6	4.4	5.3	8.1																
18-year	2.0	2.3	3.7	3.4	2.8	4.0	4.4	4.1	5.3																	
19-year	2.0	2.2	3.7	3.3	2.8	4.0	4.4	4.1																		
20-year	2.0	2.2	3.7	3.3	2.8	3.9	4.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 14.10
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	11.3	12.8	14.8	20.2	21.9	24.7	26.8	25.4	28.7	28.5	30.7	32.2	35.3	37.5	40.0	39.4	41.3	45.5	45.5	49.2	46.9	46.5	48.5	49.5	52.6	52.5
2-year	4.8	6.5	7.3	11.4	12.2	16.0	16.6	16.2	17.8	19.3	20.6	21.3	24.8	25.6	27.8	29.7	29.8	33.0	31.8	36.8	33.5	32.3	35.0	37.1	39.3	
3-year	3.2	4.9	5.2	8.1	9.0	11.3	12.1	12.1	13.8	15.6	15.3	18.0	20.1	19.9	21.7	23.8	24.2	26.1	24.5	29.0	26.6	24.4	26.8	29.9		
4-year	2.7	3.4	4.5	6.3	6.2	9.4	9.7	10.6	11.4	12.4	12.9	15.8	16.9	17.5	18.6	20.0	20.5	22.0	19.8	23.0	23.0	20.2	22.7			
5-year	2.0	2.8	3.5	5.2	5.0	8.0	8.0	8.9	10.1	11.4	11.1	14.5	15.8	14.4	17.0	17.0	18.2	19.0	17.4	19.9	20.4	17.7				
6-year	1.7	2.6	3.3	4.8	4.3	7.0	7.5	7.7	9.0	10.6	10.2	12.2	15.0	13.0	15.6	15.4	16.1	17.0	15.6	17.4	18.8					
7-year	1.4	2.2	3.3	4.3	3.7	6.6	6.9	6.2	8.3	10.1	9.3	11.3	13.4	12.3	14.1	14.5	14.0	15.9	14.6	16.1						
8-year	1.4	1.9	3.0	4.1	3.5	6.5	5.7	6.1	7.7	10.0	8.1	10.2	12.2	11.7	13.3	13.7	13.2	14.9	12.9							
9-year	1.3	1.9	3.0	3.9	3.4	6.0	5.6	5.3	7.5	9.9	7.6	9.7	11.4	11.3	12.2	12.9	12.7	14.2								
10-year	1.3	1.9	2.9	3.8	2.8	5.6	5.6	5.3	7.4	9.6	7.1	8.7	10.7	10.4	11.7	12.7	11.9									
11-year	1.3	1.8	2.9	3.7	2.3	5.3	5.6	4.4	6.9	9.3	7.1	7.9	10.5	9.8	11.6	12.1										
12-year	1.3	1.7	2.7	3.6	2.1	5.1	5.6	4.3	6.9	9.2	7.0	7.4	10.1	9.6	11.6											
13-year	1.1	1.4	2.7	3.5	2.1	5.0	5.6	4.3	6.8	8.8	6.8	7.0	9.8	9.4												
14-year	1.1	1.4	2.6	3.5	2.1	4.6	5.5	4.2	6.3	8.8	6.2	6.6	9.5													
15-year	1.0	1.4	2.6	3.5	2.1	4.0	5.2	4.2	6.3	8.8	5.6	6.5														
16-year	1.0	1.4	2.6	3.4	2.1	4.0	4.6	4.1	5.9	8.6	5.5															
17-year	1.0	1.4	2.6	3.4	2.1	4.0	4.3	4.0	5.8	8.5																
18-year	1.0	1.4	2.6	3.3	2.1	4.0	3.9	3.7	5.7																	
19-year	1.0	1.4	2.6	3.2	2.1	4.0	3.9	3.7																		
20-year	1.0	1.4	2.6	3.2	2.1	4.0	3.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 14.11
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	18.4	19.0	23.7	24.0	26.3	24.6	29.6	29.7	25.6	30.1	32.4	34.5	38.6	40.7	37.0	38.7	44.5	43.0	42.3	46.4	47.3	46.6	50.4	48.2	52.4	52.4
2-year	10.0	11.7	14.5	14.3	16.5	12.9	18.9	18.7	17.9	21.2	24.5	24.7	24.7	28.0	25.1	26.5	32.0	31.4	29.5	34.0	33.5	32.8	36.3	35.7	39.3	
3-year	7.8	8.3	11.4	10.1	10.4	9.9	13.2	15.2	15.3	15.1	17.8	16.7	21.5	22.9	20.2	22.5	23.6	24.9	23.7	25.0	25.6	25.9	28.7	28.3		
4-year	6.6	6.8	9.9	8.1	8.3	8.7	11.0	10.8	13.3	12.3	15.9	14.9	17.7	18.1	17.8	19.6	22.3	20.7	19.7	21.3	21.9	21.4	24.4			
5-year	6.1	5.7	8.8	6.3	7.0	8.5	8.4	10.2	10.3	12.1	14.0	13.3	15.1	15.5	16.2	17.3	19.4	17.9	15.8	18.0	20.0	19.2				
6-year	5.4	5.3	8.4	5.6	6.0	7.3	7.3	9.9	9.2	11.1	12.7	11.8	14.9	14.3	14.8	15.9	17.8	16.9	14.6	17.1	18.0					
7-year	5.4	4.9	7.7	5.1	5.7	6.7	6.2	8.7	8.2	9.7	11.3	11.6	13.1	13.9	13.9	14.9	16.1	15.0	14.2	14.7						
8-year	5.2	4.7	7.0	5.1	5.4	6.4	6.2	7.5	8.0	9.0	10.8	10.8	12.6	13.1	13.9	13.7	15.1	13.0	13.6							
9-year	4.9	4.6	6.6	5.0	5.4	5.9	5.8	6.2	7.4	8.1	10.6	10.8	12.1	12.3	12.8	13.1	13.6	12.2								
10-year	4.9	4.5	6.2	5.0	5.1	5.9	5.2	5.9	7.4	7.9	9.5	10.5	12.1	12.1	11.6	12.6	12.2									
11-year	4.9	4.5	6.0	4.5	4.8	5.9	5.2	5.6	6.9	7.4	9.2	10.3	11.9	11.7	11.5	12.2										
12-year	4.5	4.5	6.0	4.4	4.8	5.7	5.2	5.3	6.4	7.4	9.0	9.8	11.7	10.6	11.1											
13-year	4.4	4.4	5.8	4.4	4.8	5.4	5.2	5.0	6.0	7.4	9.0	9.0	11.2	10.4												
14-year	4.4	4.4	5.8	4.2	4.8	5.4	5.2	5.0	5.4	7.4	9.0	9.0	11.2													
15-year	4.4	4.3	5.8	4.0	4.5	4.3	5.0	5.0	5.0	7.3	9.0	8.6														
16-year	4.3	4.3	5.8	4.0	4.5	3.9	5.0	5.0	4.3	7.3	8.9															
17-year	4.0	4.1	5.8	3.8	4.1	3.9	5.0	5.0	4.3	7.2																
18-year	3.8	3.9	5.7	3.6	3.7	3.9	5.0	5.0	4.3																	
19-year	3.8	3.6	5.7	3.6	3.7	3.5	5.0	5.0																		
20-year	3.8	3.6	5.6	3.4	3.7	3.5	5.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 15.15
Cancer of the Lung and Bronchus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

Survival Time	1975-	1980-	1985-	1990-																										
	1979	1984	1989	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016				
1-year	35.3	37.1	38.4	39.3	40.5	39.8	40.1	39.7	41.6	40.7	40.0	40.5	42.0	42.0	43.3	43.8	44.5	45.0	45.6	44.7	46.9	45.8	47.9	48.0	50.2	51.8				
2-year	21.1	21.9	22.6	23.7	24.9	24.6	24.6	25.2	25.9	25.8	25.5	26.5	26.6	27.7	28.4	29.5	30.1	30.9	31.4	31.2	32.9	31.8	33.6	35.0	37.2					
3-year	16.3	16.9	17.1	18.3	19.1	19.3	19.3	19.9	19.8	20.2	19.8	20.7	20.8	22.3	22.6	23.5	24.0	24.8	25.3	25.0	26.5	25.6	27.4	28.9						
4-year	14.0	14.5	14.6	15.5	16.5	16.3	16.6	17.0	17.0	17.5	16.8	17.5	17.9	18.9	19.5	20.0	20.6	21.4	21.7	21.6	23.0	22.2	24.2							
5-year	12.5	12.9	13.0	13.8	14.5	14.6	14.7	14.6	15.2	15.7	15.0	15.6	16.0	16.8	17.4	17.6	18.4	18.9	19.6	19.4	20.5	19.9								
6-year	11.4	11.7	11.9	12.7	13.2	13.2	13.3	13.3	13.7	14.2	13.5	14.1	14.5	15.4	15.7	16.2	16.8	17.3	17.9	17.8	18.7									
7-year	10.7	10.8	10.9	11.6	12.3	12.1	12.2	12.2	12.7	13.0	12.4	12.9	13.3	14.1	14.6	15.1	15.4	16.0	16.6	16.2										
8-year	10.0	10.0	10.1	10.7	11.5	11.2	11.3	11.2	11.8	12.0	11.5	11.9	12.1	13.2	13.5	14.1	14.2	14.6	15.3											
9-year	9.4	9.2	9.4	9.9	10.7	10.3	10.4	10.5	11.1	11.2	10.6	11.1	11.1	11.1	12.5	12.5	12.7	13.3	13.6											
10-year	8.8	8.6	8.8	9.2	10.0	9.5	9.8	10.0	10.5	10.4	10.1	10.4	10.3	11.6	11.6	11.7	12.4													
11-year	8.3	8.2	8.2	8.6	9.4	8.8	9.2	9.4	9.7	9.8	9.4	9.5	9.8	10.8	10.9	11.2														
12-year	7.8	7.7	7.7	8.0	8.9	8.3	8.5	8.9	9.1	9.1	8.8	8.9	9.3	10.1	10.3															
13-year	7.4	7.2	7.2	7.5	8.2	7.9	8.0	8.1	8.6	8.6	8.2	8.3	8.7	9.4																
14-year	7.1	6.8	6.8	7.1	7.7	7.6	7.5	7.6	8.0	8.1	7.7	7.9	7.9																	
15-year	6.6	6.4	6.4	6.7	7.3	7.0	7.1	7.1	7.6	7.7	7.2	7.6																		
16-year	6.3	6.0	6.1	6.3	6.9	6.7	6.6	6.8	7.2	7.5	6.9																			
17-year	6.0	5.6	5.7	5.9	6.4	6.3	6.2	6.3	6.8	7.1																				
18-year	5.7	5.3	5.3	5.5	5.9	6.0	5.7	6.1	6.4																					
19-year	5.4	5.0	5.0	5.2	5.5	5.6	5.4	5.8																						
20-year	5.2	4.6	4.7	4.9	5.2	5.3	5.0																							
21-year	4.9	4.3	4.5	4.6	5.0																									
22-year	4.7	4.1	4.2	4.4	4.7																									
23-year	4.4	3.8	4.0	4.1																										
24-year	4.2	3.6	3.8																											
25-year	4.0	3.4	3.6																											
26-year	3.8	3.2	3.5																											
27-year	3.6	3.1	3.3																											
28-year	3.5	2.9	3.1																											
29-year	3.3	2.7																												
30-year	3.2	2.6																												

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 15.16
Cancer of the Lung and Bronchus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	33.4	35.2	36.4	37.1	38.2	37.9	37.8	36.7	38.4	38.5	36.7	36.9	39.5	38.8	39.7	41.1	41.0	41.8	42.5	41.0	42.6	41.6	42.7	43.9	46.9	47.8
2-year	19.5	20.2	21.0	21.6	23.1	22.9	22.9	22.7	22.8	23.7	22.2	23.2	23.8	24.7	24.7	26.9	26.2	27.7	27.5	27.4	28.6	28.0	28.0	30.9	33.3	
3-year	14.8	15.3	15.6	16.4	17.3	17.7	17.7	17.4	16.7	18.2	17.2	18.0	18.2	19.5	19.5	20.8	20.9	22.0	21.9	21.6	22.5	22.1	22.1	25.2		
4-year	12.5	12.9	13.1	13.7	14.8	14.6	15.0	14.7	14.1	15.6	14.4	15.1	15.5	16.7	16.6	17.3	17.6	18.9	18.4	18.6	19.0	18.9	19.4			
5-year	11.1	11.4	11.6	12.1	13.2	13.1	13.3	12.4	12.5	14.1	12.7	13.6	14.1	14.9	14.8	15.1	15.7	16.7	16.5	16.5	16.8	16.8				
6-year	10.1	10.4	10.6	11.0	11.8	11.7	11.9	11.0	11.2	12.6	11.6	12.2	12.6	13.5	13.2	13.9	14.3	15.2	15.1	15.2	15.3					
7-year	9.4	9.5	9.7	10.0	11.2	10.6	10.8	10.1	10.3	11.4	10.5	11.2	11.6	12.3	12.2	12.8	12.8	14.1	13.7	13.9						
8-year	8.8	8.7	9.0	9.1	10.2	9.8	10.1	9.0	9.5	10.7	9.6	10.1	10.8	11.4	11.2	11.9	11.7	12.9	12.6							
9-year	8.2	7.9	8.4	8.4	9.6	9.1	9.2	8.5	8.8	10.0	8.9	9.3	9.9	10.8	10.4	10.7	10.9	12.0								
10-year	7.6	7.4	7.9	7.9	8.8	8.2	8.6	8.0	8.3	9.2	8.4	8.7	8.9	10.0	9.5	9.6	10.0									
11-year	7.2	7.0	7.3	7.2	8.3	7.4	8.0	7.5	7.7	8.5	7.8	8.1	8.4	9.4	8.8	8.9										
12-year	6.7	6.6	6.8	6.7	8.0	6.9	7.3	7.0	7.2	7.8	7.3	7.4	8.1	8.7	8.3											
13-year	6.3	6.2	6.3	6.2	7.3	6.7	6.8	6.3	6.8	7.3	6.5	6.9	7.6	8.2												
14-year	6.0	5.8	6.0	5.8	6.9	6.3	6.4	5.9	6.3	6.7	6.2	6.5	6.9													
15-year	5.6	5.5	5.6	5.5	6.4	5.8	6.1	5.6	5.9	6.5	5.5	6.3														
16-year	5.3	5.0	5.3	5.1	6.2	5.5	5.7	5.2	5.8	6.3	5.3															
17-year	5.1	4.6	4.9	4.8	5.8	5.2	5.3	4.9	5.4	6.0																
18-year	4.7	4.4	4.5	4.5	5.3	4.9	4.9	4.6	5.1																	
19-year	4.5	4.1	4.2	4.3	4.6	4.6	4.5	4.5																		
20-year	4.2	3.8	3.9	4.0	4.3	4.5	4.0																			
21-year	4.1	3.5	3.6	3.7	4.1	4.1																				
22-year	3.9	3.3	3.4	3.5	3.8																					
23-year	3.7	3.1	3.2	3.2																						
24-year	3.4	2.8	3.0																							
25-year	3.3	2.7	2.8																							
26-year	3.1	2.5	2.6																							
27-year	2.8	2.5	2.5																							
28-year	2.8	2.4	2.3																							
29-year	2.7	2.2																								
30-year	2.6	2.1																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 15.17
Cancer of the Lung and Bronchus (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	40.4	41.3	42.0	42.7	43.6	42.3	43.0	43.5	45.6	43.6	44.1	44.7	45.0	45.7	47.2	46.9	48.4	48.6	48.9	48.8	51.5	50.4	53.4	52.3	53.7	56.0
2-year	25.4	25.7	25.5	26.7	27.5	26.9	26.8	28.5	29.6	28.5	29.6	30.5	29.9	31.1	32.3	32.6	34.4	34.5	35.5	35.5	37.4	35.9	39.6	39.2	41.4	
3-year	20.3	20.5	19.7	21.1	21.6	21.4	21.3	23.0	23.6	22.7	23.0	24.0	23.8	25.4	26.1	26.6	27.4	27.8	29.0	28.7	30.8	29.5	32.9	32.8		
4-year	17.8	17.8	17.2	18.3	18.8	18.4	18.5	19.9	20.5	19.9	19.9	20.4	20.7	21.5	22.8	23.0	23.8	24.1	25.2	24.9	27.1	25.8	29.4			
5-year	16.1	16.0	15.4	16.4	16.5	16.5	16.4	17.4	18.4	17.7	17.8	17.9	18.3	19.0	20.2	20.4	21.3	21.2	23.0	22.5	24.4	23.2				
6-year	15.0	14.6	14.1	15.1	15.0	15.2	15.0	16.1	16.8	16.1	15.8	16.4	16.8	17.6	18.5	18.8	19.6	19.6	21.0	20.7	22.3					
7-year	14.0	13.6	12.9	13.9	13.9	14.1	14.0	14.9	15.6	15.0	14.7	15.0	15.3	16.3	17.1	17.7	18.2	17.9	19.6	18.8						
8-year	13.1	12.8	12.0	13.0	13.1	13.1	12.8	13.9	14.7	13.6	13.8	14.0	13.6	15.2	16.0	16.5	17.0	16.4	18.1							
9-year	12.4	11.9	11.2	12.1	12.3	12.0	11.9	13.1	13.8	12.8	12.7	13.1	12.7	14.3	14.8	15.0	16.0	15.4								
10-year	11.6	11.1	10.4	11.3	11.6	11.1	11.2	12.4	13.1	12.0	12.2	12.4	11.9	13.3	13.9	14.2	15.0									
11-year	11.1	10.6	9.8	10.6	11.0	10.7	10.6	11.9	12.2	11.3	11.5	11.3	11.4	12.3	13.1	13.7										
12-year	10.7	9.9	9.3	9.9	10.1	10.1	9.9	11.3	11.4	10.7	10.7	10.6	10.7	11.6	12.5											
13-year	10.3	9.3	8.8	9.4	9.4	9.6	9.4	10.5	10.7	10.2	10.2	9.9	9.9	10.7												
14-year	9.7	8.9	8.2	8.9	8.7	9.1	8.9	9.7	10.0	9.6	9.6	9.5	9.1													
15-year	9.2	8.4	7.8	8.4	8.4	8.7	8.3	9.1	9.6	9.2	9.2	9.0														
16-year	8.9	8.0	7.4	7.9	7.9	8.2	7.6	8.6	8.9	9.0	8.9															
17-year	8.4	7.7	7.1	7.4	7.3	7.8	7.2	8.1	8.5	8.4																
18-year	8.0	7.2	6.7	7.0	6.8	7.4	6.7	7.8	8.1																	
19-year	7.7	6.6	6.3	6.5	6.5	6.9	6.4	7.3																		
20-year	7.4	6.3	6.1	6.2	6.3	6.4	6.2																			
21-year	6.9	5.9	5.9	5.9	6.1	6.2																				
22-year	6.5	5.6	5.5	5.6	5.8																					
23-year	6.2	5.3	5.4	5.4																						
24-year	6.0	5.1	5.1																							
25-year	5.8	4.8	4.9																							
26-year	5.6	4.6	4.8																							
27-year	5.4	4.2	4.6																							
28-year	5.2	4.0	4.4																							
29-year	4.9	3.7																								
30-year	4.6	3.6																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 16.9
Melanoma of the Skin (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																									
	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	94.2	95.6	96.3	96.7	97.3	97.6	96.6	96.6	97.7	97.8	97.4	97.9	97.7	97.7	97.8	97.6	97.8	97.9	98.1	97.8	97.8	98.1	98.6	97.9	98.3	98.6
2-year	89.8	91.7	93.2	93.9	94.4	95.7	94.2	94.5	95.6	96.2	95.5	95.5	96.1	96.1	96.3	96.1	96.3	95.9	96.1	96.7	96.4	96.6	97.9	96.7	97.3	
3-year	86.1	88.0	90.8	92.0	92.7	93.8	92.3	92.7	94.0	94.5	94.1	94.3	94.9	94.6	95.0	94.7	94.8	94.8	94.8	95.3	95.0	96.0	96.7	96.0		
4-year	83.4	85.5	89.2	90.3	91.3	93.2	91.3	91.5	93.2	93.2	92.7	93.2	93.7	93.6	94.3	93.7	94.1	94.2	93.9	94.6	94.3	95.5	96.2			
5-year	82.0	83.5	87.7	89.4	90.2	92.2	89.8	90.9	92.4	92.1	92.1	92.9	93.3	93.1	93.3	93.0	93.5	93.6	93.4	93.9	93.8	95.1				
6-year	80.2	82.3	86.6	88.5	90.0	91.5	89.3	90.4	91.7	91.8	91.3	92.5	92.5	93.1	93.0	92.3	92.9	92.8	93.0	93.7	93.5					
7-year	79.0	81.7	85.9	87.9	90.0	91.2	88.6	90.3	91.6	91.5	91.1	92.0	91.9	92.5	92.9	92.1	92.9	92.3	92.7	93.6						
8-year	77.9	80.9	85.3	87.3	89.9	90.9	87.8	89.9	91.2	91.4	90.9	91.2	91.8	92.4	92.8	92.1	92.5	92.2	92.7							
9-year	77.1	80.3	85.1	87.1	89.5	90.4	87.6	89.7	91.0	91.4	90.8	90.7	91.8	92.2	92.8	92.1	92.3	92.0								
10-year	76.6	79.8	84.9	86.9	89.4	89.9	87.6	89.3	91.0	91.3	90.2	90.6	91.5	92.2	92.8	92.1	92.1									
11-year	76.0	79.4	84.4	86.6	89.2	89.7	87.6	89.3	91.0	90.9	90.0	90.5	91.5	92.2	92.8	92.1										
12-year	75.6	79.2	84.2	86.5	89.2	89.7	87.1	88.8	91.0	90.9	90.0	90.5	91.5	92.2	92.8											
13-year	75.3	78.8	84.0	86.5	88.7	89.6	86.9	88.8	91.0	90.9	89.9	90.5	91.4	92.2												
14-year	75.1	78.6	83.9	86.4	88.7	89.5	86.7	88.7	91.0	90.9	89.9	90.5	91.3													
15-year	74.7	78.4	83.9	86.4	88.7	89.5	86.6	88.6	91.0	90.9	89.9	90.5														
16-year	74.5	78.4	83.8	86.4	88.7	89.5	86.6	88.0	91.0	90.9	89.9															
17-year	74.5	78.1	83.8	86.4	88.7	89.2	86.6	88.0	91.0	90.9																
18-year	74.5	78.1	83.8	86.4	88.7	89.2	86.6	88.0	91.0																	
19-year	74.5	78.1	83.8	86.4	88.7	89.2	86.6	88.0																		
20-year	74.5	78.1	83.8	86.4	88.7	89.2	86.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 16.10
Melanoma of the Skin (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	92.3	94.0	95.2	96.1	96.3	96.6	96.3	96.2	97.2	97.4	96.5	96.8	97.0	97.5	97.5	97.1	97.5	97.0	97.4	97.3	97.2	97.3	98.4	97.2	97.7	98.2
2-year	86.6	88.7	91.4	92.6	92.2	94.4	93.2	93.8	94.3	95.5	93.7	94.1	94.8	95.5	95.9	95.4	95.4	94.6	94.6	96.1	95.4	95.5	97.0	95.7	96.4	
3-year	82.3	84.3	88.3	90.3	90.0	92.1	90.8	92.0	92.5	93.5	91.5	92.5	93.5	92.7	94.1	93.6	93.5	93.2	93.3	94.2	93.5	94.7	95.6	94.8		
4-year	79.0	81.3	86.2	88.4	88.1	91.2	89.8	90.4	91.4	92.1	89.8	90.8	91.8	91.8	92.9	92.7	92.8	92.3	92.1	93.7	92.5	93.9	95.2			
5-year	77.5	78.6	84.4	87.3	86.8	90.3	88.0	90.1	90.5	91.0	89.1	90.2	91.8	91.1	91.5	91.8	91.9	91.9	91.3	92.7	91.8	93.2				
6-year	75.4	77.3	83.2	86.3	86.4	89.6	87.2	89.6	89.1	90.4	88.2	90.0	90.7	91.1	91.0	91.0	91.1	90.5	91.0	92.3	91.4					
7-year	73.8	76.9	82.2	85.5	86.4	89.4	87.0	89.6	89.0	90.4	88.0	89.6	89.7	90.2	90.6	90.7	90.9	90.0	90.9	92.2						
8-year	72.6	76.0	81.6	84.7	86.2	88.9	86.1	89.3	88.8	90.1	88.0	88.5	89.4	89.9	90.1	90.7	90.5	89.7	90.9							
9-year	71.7	75.2	81.3	84.4	85.8	88.4	86.0	88.8	88.5	90.1	88.0	87.8	89.4	89.6	90.1	90.4	90.1	89.3								
10-year	71.2	74.8	81.3	84.2	85.7	87.9	85.5	88.3	88.5	90.0	87.4	87.8	89.4	89.6	89.8	90.4	89.9									
11-year	70.4	74.4	81.1	83.9	85.7	87.7	85.5	88.2	88.5	89.4	87.1	87.8	89.3	89.6	89.8	90.4										
12-year	69.9	74.3	80.9	83.8	85.7	87.7	84.7	87.6	88.5	89.3	87.1	87.8	89.3	89.6	89.8											
13-year	69.6	74.0	80.8	83.8	85.7	87.7	84.7	87.5	88.5	89.3	87.1	87.8	89.3	89.6												
14-year	69.6	73.6	80.4	83.6	85.7	87.7	84.2	87.3	88.5	89.3	87.1	87.8	89.3													
15-year	69.3	73.6	80.2	83.6	85.7	87.7	84.2	87.3	88.5	89.3	87.1	87.8														
16-year	69.1	73.6	80.0	83.6	85.5	87.7	84.2	87.1	88.5	89.3	87.1															
17-year	69.1	73.6	80.0	83.6	85.5	87.7	84.2	87.1	88.5	89.3																
18-year	69.1	73.6	80.0	83.6	85.5	87.7	84.2	87.1	88.5																	
19-year	69.1	73.6	80.0	83.6	85.5	87.7	84.2	87.1																		
20-year	69.1	73.6	80.0	83.6	85.5	87.7	84.2																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 16.11
Melanoma of the Skin (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	96.1	97.3	97.6	97.4	98.5	98.8	97.0	97.1	98.2	98.4	98.5	99.0	98.6	97.9	98.2	98.1	98.2	98.9	99.1	98.3	98.5	99.0	99.0	98.7	99.0	98.9
2-year	92.9	94.7	95.3	95.5	96.8	97.3	95.4	95.3	97.0	97.1	97.6	97.2	97.6	96.9	96.8	97.0	97.3	97.4	97.9	97.4	97.6	98.0	98.9	98.1	98.3	
3-year	89.9	91.9	93.5	93.9	95.7	95.9	94.1	93.6	95.7	95.7	97.1	96.4	96.6	96.5	96.2	95.9	96.5	96.7	96.8	96.5	96.9	97.5	98.0	97.5		
4-year	87.8	89.9	92.4	92.5	94.9	95.3	92.9	92.6	95.3	94.7	96.2	96.0	95.7	95.6	96.0	95.0	95.8	96.2	96.1	95.7	96.6	97.4	97.4			
5-year	86.4	88.6	91.3	91.8	94.1	94.6	91.8	91.8	94.6	93.4	95.6	95.7	95.1	95.4	95.4	94.5	95.4	95.5	95.9	95.3	96.3	97.4				
6-year	85.0	87.4	90.3	91.2	94.1	93.8	91.6	91.2	94.5	93.4	95.1	95.3	94.5	95.4	95.1	93.8	95.1	95.4	95.3	95.3	95.9					
7-year	84.0	86.6	89.9	90.7	94.1	93.2	90.4	91.0	94.4	92.9	94.7	94.8	94.3	95.0	95.1	93.7	95.1	95.0	94.8	95.0						
8-year	83.1	86.0	89.3	90.3	94.1	93.2	89.7	90.5	93.8	92.8	94.2	94.3	94.1	95.0	95.1	93.7	94.8	95.0	94.5							
9-year	82.4	85.5	89.1	90.2	93.7	92.9	89.5	90.5	93.7	92.8	93.7	94.0	94.0	94.9	95.1	93.7	94.7	95.0								
10-year	81.8	84.8	88.8	90.1	93.7	92.4	89.5	90.3	93.6	92.7	93.4	93.5	93.8	94.9	95.1	93.7	94.3									
11-year	81.5	84.5	88.1	89.8	92.8	92.0	89.5	90.2	93.6	92.6	93.4	93.3	93.8	94.9	95.1	93.7										
12-year	81.1	84.3	87.9	89.6	92.8	91.5	89.5	90.2	93.6	92.5	93.3	93.3	93.5	94.9	95.1											
13-year	80.8	83.7	87.5	89.6	92.0	91.5	89.1	90.1	93.6	92.5	93.3	93.3	93.5	94.9												
14-year	80.3	83.6	87.5	89.6	91.8	91.4	89.1	90.1	93.6	92.5	93.3	93.2	93.0													
15-year	79.9	83.1	87.5	89.6	91.8	90.7	89.0	89.8	93.6	92.5	93.3	93.2														
16-year	79.8	82.9	87.5	89.6	91.8	90.2	88.9	89.0	93.6	92.5	93.3															
17-year	79.8	82.7	87.5	89.6	91.8	90.0	88.9	89.0	93.6	92.5																
18-year	79.7	82.5	87.5	89.6	91.8	89.9	88.9	89.0	93.6																	
19-year	79.6	82.5	87.5	89.6	91.8	89.8	88.9	89.0																		
20-year	79.6	82.5	87.5	89.6	91.8	89.7	88.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 17.5
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																									
	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	34.9	37.6	35.6	35.3	36.3	41.5	35.3	37.4	31.7	40.0	40.4	32.1	47.1	39.5	40.4	44.5	43.9	45.3	45.4	33.5	47.8	40.3	41.5	40.0	43.8	44.8
2-year	16.1	18.7	16.5	15.7	13.8	25.1	16.9	21.2	18.6	14.6	24.0	16.1	22.3	16.6	19.6	28.4	18.7	21.1	26.1	16.5	26.8	24.7	25.8	21.5	22.1	
3-year	10.9	11.7	10.8	11.0	11.8	17.5	10.3	14.7	12.8	10.5	13.8	7.4	16.8	10.4	12.7	19.6	11.5	14.9	16.3	8.5	17.9	14.9	19.0	14.0		
4-year	9.2	8.4	8.0	8.2	8.9	11.4	7.8	12.6	10.5	7.0	11.4	5.2	13.5	8.5	10.2	14.8	7.1	12.5	14.3	6.0	15.8	10.5	17.3			
5-year	7.9	7.0	6.6	7.1	7.1	10.2	5.7	12.6	8.2	3.1	10.9	4.8	9.0	6.0	9.2	12.6	4.8	10.4	12.9	4.9	13.6	9.5				
6-year	6.7	5.7	5.2	6.0	6.1	10.0	4.6	11.3	7.8	1.3	9.9	4.3	8.0	6.0	9.2	11.3	4.3	8.5	11.2	3.9	10.2					
7-year	6.2	5.1	4.5	5.6	5.1	8.6	3.0	10.6	7.4	1.3	7.7	3.8	7.0	4.6	9.1	10.3	3.7	7.3	10.8	3.9						
8-year	5.6	4.8	4.0	5.0	5.1	6.5	3.0	8.5	7.4	0.7	7.7	3.8	5.7	4.1	8.2	9.2	3.7	7.3	10.8							
9-year	5.3	4.6	3.8	4.7	3.0	4.9	2.5	8.5	6.0	0.7	7.7	3.8	5.7	3.5	8.2	6.4	3.7	7.0								
10-year	5.2	4.3	3.4	4.3	3.0	4.4	2.5	7.7	6.0	0.7	6.8	3.7	4.0	3.5	8.2	6.4	3.7									
11-year	5.0	4.3	3.4	4.1	3.0	4.0	1.9	7.4	5.6	0.7	5.0	3.7	4.0	3.5	8.2	6.4										
12-year	4.9	4.1	3.4	3.8	2.5	3.5	1.9	5.6	5.6	0.7	5.0	3.3	4.0	3.1	7.2											
13-year	4.7	4.0	3.3	3.6	2.5	3.0	1.9	5.6	5.6	0.7	5.0	2.8	3.5	2.5												
14-year	4.7	4.0	3.3	3.6	2.5	3.0	1.9	5.6	5.2	0.7	4.0	2.8	3.5													
15-year	4.3	3.5	3.2	3.6	2.5	3.0	1.9	5.6	5.2	0.0	3.2	1.9														
16-year	4.1	3.5	3.2	3.6	2.5	3.0	1.9	5.6	4.7	0.0	3.2															
17-year	4.0	3.5	3.2	3.6	2.5	2.9	1.9	5.4	4.7	0.0																
18-year	4.0	3.5	3.2	3.2	2.5	2.0	1.9	4.8	4.7																	
19-year	4.0	3.5	3.2	2.5	2.5	2.0	1.9	4.1																		
20-year	4.0	3.5	3.1	2.5	2.5	2.0	1.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 17.6
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	33.1	36.1	33.8	32.1	35.5	44.4	32.7	32.8	28.8	40.6	36.5	31.0	45.7	37.5	40.2	39.9	42.1	42.2	46.2	30.9	46.5	39.9	41.1	40.0	40.7	42.8
2-year	11.4	15.9	14.2	11.7	11.1	24.5	14.3	17.6	13.7	13.5	20.3	15.1	18.5	14.7	17.2	22.0	15.2	19.8	25.6	16.0	23.3	22.4	24.2	19.4	18.7	
3-year	6.6	9.5	8.4	7.5	8.9	16.2	8.7	11.7	6.8	9.0	10.8	6.7	13.9	8.9	8.9	12.8	10.5	14.2	13.9	6.2	14.9	11.6	14.8	11.9		
4-year	5.2	6.6	5.8	4.7	5.9	11.3	6.2	10.1	4.4	6.7	8.8	3.8	12.7	6.6	6.3	9.4	5.9	11.8	11.1	3.5	11.9	6.6	13.9			
5-year	4.0	5.1	4.6	3.6	4.2	10.3	3.6	10.1	3.8	4.0	8.2	3.8	7.1	3.9	6.3	7.6	4.3	9.2	8.9	2.9	10.9	6.6				
6-year	3.3	4.3	3.2	3.1	3.5	10.3	2.2	8.2	3.8	1.7	7.5	3.2	6.5	3.9	6.3	6.3	4.3	7.6	7.4	2.2	9.9					
7-year	2.9	3.4	2.5	2.7	2.2	9.5	1.5	7.8	3.8	1.7	5.1	2.4	5.7	2.5	6.3	6.3	3.5	7.6	6.7	2.2						
8-year	2.3	3.3	2.5	2.1	2.2	6.7	1.5	4.8	3.8	0.9	5.1	2.4	5.7	2.5	6.1	5.1	3.5	7.6	6.7							
9-year	2.1	3.1	2.2	1.8	0.0	5.2	0.8	4.8	2.8	0.9	5.1	2.4	5.7	1.8	6.1	4.4	3.5	7.5								
10-year	2.1	3.1	1.9	1.7	0.0	4.5	0.8	4.1	2.8	0.9	5.1	2.4	4.5	1.8	6.1	4.4	3.5									
11-year	2.1	3.1	1.9	1.6	0.0	3.9	0.0	4.1	2.2	0.9	2.8	2.4	4.5	1.8	5.7	4.4										
12-year	1.9	2.9	1.9	1.4	0.0	3.2	0.0	3.6	2.2	0.9	2.8	1.9	4.5	1.8	5.0											
13-year	1.7	2.8	1.8	1.4	0.0	2.3	0.0	3.6	2.2	0.9	2.8	1.9	3.8	1.0												
14-year	1.4	2.8	1.7	1.4	0.0	2.3	0.0	3.6	1.6	0.9	1.9	1.9	3.8													
15-year	1.1	2.5	1.6	1.4	0.0	2.3	0.0	3.6	1.6	0.0	1.9	1.0														
16-year	1.1	2.5	1.6	1.4	0.0	2.3	0.0	3.6	1.6	0.0	1.9															
17-year	1.1	2.4	1.6	1.2	0.0	1.8	0.0	3.6	1.6	0.0																
18-year	1.1	2.4	1.6	1.2	0.0	1.8	0.0	3.6	1.6																	
19-year	1.1	2.4	1.6	0.9	0.0	1.8	0.0	3.6																		
20-year	1.1	2.4	1.6	0.9	0.0	1.8	0.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 17.7
MesotheliomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	40.8	42.6	41.8	48.0	38.8	32.3	46.3	54.5	43.7	37.1	52.3	35.7	51.2	46.0	41.0	57.0	48.5	57.8	43.1	42.4	52.3	41.9	42.7	39.9	54.8	52.0
2-year	31.1	28.3	24.5	31.2	23.1	26.8	27.9	34.8	39.3	19.0	34.8	19.4	32.9	22.3	27.9	45.8	27.9	26.1	27.4	18.4	38.4	31.8	30.4	27.0	34.6	
3-year	24.4	19.4	18.9	24.4	21.2	21.3	17.2	26.0	37.4	16.7	22.8	9.8	24.8	14.3	25.9	37.8	14.3	17.8	23.1	16.4	28.1	25.4	30.4	19.5		
4-year	21.5	14.4	15.4	21.3	19.3	11.8	14.4	21.6	35.4	8.6	18.8	9.8	16.0	14.3	23.7	29.3	10.3	15.1	23.1	14.2	28.1	23.0	26.9			
5-year	20.1	13.3	13.4	20.2	17.6	9.8	14.4	21.6	26.2	0.0	18.8	7.8	14.2	12.4	19.3	26.1	6.3	15.1	23.1	12.0	22.5	18.2				
6-year	17.1	10.5	11.9	16.9	15.2	7.9	14.4	21.6	23.8	0.0	16.9	7.8	12.2	12.4	19.3	24.9	4.3	12.2	22.3	9.7	11.6					
7-year	16.3	10.5	10.8	16.5	15.2	6.0	8.9	20.5	21.4	0.0	15.0	7.8	10.2	10.4	17.6	21.1	4.3	4.3	22.3	9.7						
8-year	15.7	9.9	8.7	15.8	15.2	6.0	8.9	20.5	21.4	0.0	15.0	7.8	5.2	8.4	15.5	20.5	4.3	4.3	22.3							
9-year	15.2	9.5	8.7	15.3	12.9	4.0	8.9	20.5	19.0	0.0	15.0	7.8	5.2	8.4	15.5	11.7	4.3	4.3								
10-year	14.4	7.7	8.4	14.0	12.9	4.0	8.9	20.2	19.0	0.0	11.1	6.8	2.9	8.4	15.5	11.7	4.3									
11-year	13.7	7.7	8.4	13.7	12.9	4.0	8.9	18.1	19.0	0.0	11.1	6.8	2.9	8.4	15.5	11.7										
12-year	13.7	7.7	8.1	12.7	10.5	4.0	8.9	12.6	19.0	0.0	11.1	6.8	2.9	6.6	14.8											
13-year	13.7	7.7	8.1	11.8	10.5	4.0	8.9	12.6	19.0	0.0	11.1	5.1	2.9	6.6												
14-year	13.7	7.7	8.1	11.8	10.5	4.0	8.9	12.6	19.0	0.0	9.6	5.1	2.9													
15-year	13.7	6.8	8.1	11.6	10.5	4.0	8.9	12.6	19.0	0.0	7.3	5.1														
16-year	12.9	6.8	8.1	11.6	10.5	4.0	8.9	12.6	17.3	0.0	7.3															
17-year	12.3	6.8	8.1	11.6	10.5	4.0	8.9	9.7	17.3	0.0																
18-year	12.3	6.8	7.9	10.5	10.5	2.2	8.9	6.5	17.3																	
19-year	12.3	6.8	7.9	8.7	10.5	2.2	8.9	6.5																		
20-year	12.3	6.5	7.4	8.7	10.5	2.2	8.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 18.9
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																									
	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	67.9	69.4	72.2	71.9	74.1	70.6	74.3	74.9	73.7	72.5	72.7	73.5	77.4	74.6	76.1	73.9	77.3	77.9	81.9	81.4	80.1	81.7	81.4	83.2	82.3	83.7
2-year	52.1	54.4	57.6	57.5	58.7	57.4	60.3	62.4	60.1	60.0	59.6	63.0	65.1	66.5	64.6	66.5	67.1	69.1	72.8	72.2	71.9	74.0	72.6	75.4	74.1	
3-year	39.9	43.0	45.4	46.3	47.8	44.8	48.6	51.2	48.0	49.8	49.7	53.6	56.0	58.7	57.6	58.3	58.5	61.9	65.6	65.7	65.2	67.1	66.7	69.1		
4-year	31.5	34.3	35.8	37.0	40.2	38.2	38.7	40.2	39.1	41.3	44.4	47.2	47.9	50.9	51.0	51.4	50.8	56.2	60.0	59.1	58.2	60.8	61.3			
5-year	25.1	27.0	27.4	29.9	33.3	31.7	31.3	34.3	31.8	34.3	37.5	40.2	42.3	45.4	46.7	46.3	46.6	50.6	53.2	53.3	51.3	56.3				
6-year	20.1	21.0	22.1	24.6	28.4	27.2	26.3	30.0	27.3	29.3	32.9	35.6	37.8	40.1	41.6	41.3	41.7	44.9	49.0	49.7	46.4					
7-year	16.3	16.7	17.6	20.5	23.7	23.7	22.4	24.7	23.7	25.4	29.5	32.1	33.3	35.8	37.1	36.7	37.5	42.0	44.2	45.9						
8-year	13.6	13.4	15.0	17.4	20.4	21.4	19.8	22.5	20.3	22.8	26.0	28.9	30.2	32.1	33.8	33.0	34.3	38.7	40.8							
9-year	11.5	11.3	11.9	15.4	18.2	19.5	17.1	19.8	18.5	20.6	23.4	26.0	27.4	30.1	31.3	31.0	31.0	35.8								
10-year	10.0	9.6	10.4	13.8	17.1	17.2	15.5	18.1	16.7	19.5	21.4	23.4	24.4	27.1	28.6	28.9	28.6									
11-year	8.8	8.7	9.2	12.8	15.6	15.6	14.3	17.0	14.4	18.6	20.1	20.5	22.1	25.1	25.5	27.6										
12-year	7.8	7.7	8.4	11.2	14.3	14.9	13.0	15.4	13.9	17.0	19.1	19.3	20.8	23.2	23.9											
13-year	6.6	6.6	7.6	10.4	13.6	13.3	11.9	14.4	13.0	16.0	17.1	18.4	19.1	22.2												
14-year	6.1	6.1	7.0	9.6	12.4	12.7	11.2	14.1	11.7	15.0	16.3	17.3	17.8													
15-year	5.4	5.7	6.1	9.3	12.0	11.6	11.0	12.5	11.4	14.2	15.2	16.1														
16-year	4.9	5.2	5.7	8.7	10.9	10.0	10.5	11.5	10.7	13.4	14.4															
17-year	4.6	4.7	5.5	8.2	10.1	9.0	9.9	10.8	10.4	12.7																
18-year	4.1	4.3	5.1	7.6	9.9	8.5	8.7	10.1	9.5																	
19-year	3.8	3.8	5.0	7.0	9.0	7.9	8.1	9.5																		
20-year	3.4	3.5	4.8	6.9	7.8	7.3	7.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 18.10
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	67.8	69.5	73.1	72.6	74.4	71.1	74.0	76.2	75.8	73.7	74.3	74.4	77.2	76.9	76.4	73.4	77.7	77.0	83.5	83.2	80.5	80.6	82.0	84.9	82.6	84.3
2-year	52.4	54.5	59.2	58.5	59.8	58.0	60.8	63.7	61.5	62.0	61.5	61.8	65.4	68.3	64.2	67.0	68.4	68.0	74.0	73.9	72.8	71.7	73.2	77.1	75.1	
3-year	39.3	42.8	46.2	46.7	49.3	44.7	50.3	53.6	50.8	52.3	51.3	52.9	56.6	60.6	57.1	58.6	60.7	59.6	67.1	66.9	64.8	64.3	67.0	71.9		
4-year	30.9	33.7	36.7	37.2	41.3	39.7	40.8	44.0	42.9	43.4	46.7	47.2	47.9	51.9	51.2	52.2	52.4	53.2	61.7	59.4	57.8	59.3	62.7			
5-year	24.3	25.9	28.4	30.4	36.3	33.6	31.5	38.2	35.2	36.0	40.3	42.1	42.4	46.3	47.5	47.7	48.2	48.0	53.8	53.8	52.2	54.9				
6-year	19.3	20.8	23.5	25.3	31.8	29.6	26.3	34.3	30.0	30.2	35.4	36.9	37.9	40.8	41.6	43.6	43.4	41.6	48.9	50.3	48.5					
7-year	15.5	16.9	19.2	21.5	26.1	25.7	22.2	28.2	26.3	26.3	31.9	33.6	34.6	36.6	36.8	38.2	39.2	39.7	44.6	46.3						
8-year	13.1	13.8	16.3	18.8	22.2	23.9	20.0	25.8	21.7	23.6	27.8	30.4	31.0	33.4	34.5	34.5	36.8	36.0	41.7							
9-year	11.2	11.7	13.0	16.9	20.2	21.1	18.0	23.6	19.5	20.0	25.0	27.4	28.7	31.5	31.9	32.2	33.3	34.0								
10-year	9.4	9.8	11.3	15.0	19.2	18.2	16.3	22.0	17.7	18.8	22.5	24.9	24.2	29.0	28.6	29.7	30.8									
11-year	7.6	8.9	10.0	13.8	17.6	16.9	14.9	21.1	14.5	17.4	21.3	22.3	21.9	27.1	24.9	28.9										
12-year	7.0	7.7	9.5	12.2	15.7	16.5	14.3	19.3	13.8	15.8	20.1	21.6	20.8	25.9	23.9											
13-year	5.8	6.7	8.5	11.4	15.1	14.4	13.1	17.6	13.1	14.9	18.1	20.6	19.2	24.7												
14-year	5.3	6.1	8.1	10.7	13.5	14.0	12.7	17.2	11.5	14.2	17.7	19.5	18.3													
15-year	4.7	5.7	7.2	10.2	13.3	12.7	12.2	16.4	11.1	13.8	16.2	18.0														
16-year	4.4	5.2	6.8	9.9	11.6	9.8	11.7	15.0	10.2	13.8	15.4															
17-year	4.0	4.8	6.5	9.1	10.8	9.4	10.8	14.6	10.1	13.1																
18-year	3.6	4.4	6.2	8.4	10.5	9.4	9.1	13.7	9.2																	
19-year	3.4	4.2	6.2	7.5	9.5	9.0	9.1	13.0																		
20-year	2.8	3.9	6.1	7.4	8.2	7.7	8.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 18.11
MyelomaSEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	68.0	69.2	71.2	71.0	73.8	70.0	74.6	73.5	71.3	71.0	70.9	72.5	77.7	72.0	75.7	74.4	76.7	78.9	79.9	79.1	79.6	83.0	80.7	80.9	81.8	82.9
2-year	51.9	54.3	55.9	56.3	57.5	56.7	59.7	61.0	58.5	57.7	57.5	64.6	64.7	64.4	65.1	65.9	65.3	70.3	71.3	70.1	70.9	76.8	71.9	73.2	72.8	
3-year	40.6	43.1	44.5	45.7	46.0	45.0	46.9	48.6	44.9	46.9	47.8	54.5	55.3	56.4	58.3	58.1	55.5	64.3	63.7	64.3	65.6	70.5	66.2	65.5		
4-year	32.2	34.9	35.0	36.8	38.9	36.5	36.5	35.9	34.7	39.0	41.8	47.1	48.0	49.8	50.9	50.6	48.7	59.4	57.8	58.8	58.8	62.7	59.6			
5-year	26.0	28.1	26.5	29.4	30.0	29.7	31.1	30.0	27.8	32.3	34.3	37.8	42.1	44.2	45.7	44.7	44.3	53.3	52.5	52.8	50.2	58.0				
6-year	21.1	21.3	20.7	23.7	24.7	24.6	26.3	25.3	24.2	28.2	30.1	33.8	37.7	39.2	41.5	38.8	39.4	48.5	49.2	48.8	44.2					
7-year	17.2	16.5	15.9	19.4	21.0	21.7	22.6	20.8	20.7	24.4	26.8	30.2	32.0	34.9	37.4	35.0	35.3	44.4	43.7	45.5						
8-year	14.1	13.1	13.6	15.9	18.2	18.7	19.5	18.7	18.6	21.7	23.9	27.0	29.2	30.6	33.0	31.4	31.0	41.5	39.8							
9-year	11.9	10.8	10.9	13.9	16.1	17.8	16.2	15.6	17.2	21.3	21.5	24.3	25.9	28.4	30.4	29.6	27.8	37.7								
10-year	10.6	9.4	9.6	12.5	14.8	16.1	14.7	14.0	15.5	20.3	20.2	21.6	24.6	25.0	28.6	28.0	25.6									
11-year	10.0	8.4	8.4	11.6	13.4	14.3	13.7	12.5	14.2	19.8	18.8	18.1	22.2	22.9	26.2	26.1										
12-year	8.7	7.6	7.3	10.1	12.5	13.2	11.5	11.1	14.0	18.3	18.0	16.4	20.8	20.1	23.8											
13-year	7.5	6.6	6.7	9.2	11.7	12.1	10.4	10.9	12.7	17.3	16.1	15.5	18.9	19.2												
14-year	6.9	6.0	6.0	8.5	11.1	11.2	9.6	10.7	11.8	15.9	14.7	14.5	17.1													
15-year	6.1	5.7	5.1	8.3	10.5	10.4	9.6	8.3	11.5	14.4	14.0	13.7														
16-year	5.4	5.1	4.7	7.5	10.0	10.0	9.2	7.7	11.1	12.9	13.2															
17-year	5.2	4.6	4.4	7.2	9.2	8.5	8.8	6.8	10.5	12.2																
18-year	4.5	4.2	4.0	6.8	9.2	7.4	8.3	6.1	9.7																	
19-year	4.3	3.5	3.8	6.5	8.5	6.7	7.1	5.6																		
20-year	3.9	3.2	3.5	6.3	7.2	6.7	6.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 19.9
Non-Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	70.2	73.0	71.2	70.4	70.4	73.2	75.5	76.6	77.5	77.2	78.6	80.9	80.3	81.0	81.6	80.9	81.7	84.1	83.8	84.3	83.5	84.1	84.4	83.5	84.6	85.0
2-year	60.2	63.7	62.8	62.2	61.9	65.8	68.4	70.1	71.1	71.4	73.3	76.1	75.6	76.0	77.3	76.9	77.6	78.9	79.5	79.6	79.8	80.0	80.1	79.8	80.8	
3-year	54.6	58.5	58.2	57.7	57.6	62.0	64.9	66.4	67.9	70.1	73.2	73.1	73.5	74.8	74.5	75.1	76.2	77.3	77.5	77.6	77.9	78.1	78.2			
4-year	50.6	54.6	54.4	54.4	54.8	58.8	62.1	63.6	64.6	65.5	68.1	71.2	71.6	71.5	73.1	72.6	73.3	74.5	75.6	75.3	75.7	76.9	76.6			
5-year	46.8	50.7	51.0	51.6	52.1	56.0	59.3	61.0	62.1	63.8	66.0	69.3	70.0	70.1	71.6	70.5	71.4	72.8	74.1	74.3	74.3	75.2				
6-year	43.6	47.5	48.3	49.3	49.9	54.1	57.1	58.3	59.6	62.1	64.6	68.1	68.3	68.7	70.3	68.9	70.3	71.8	72.2	72.4	72.3					
7-year	40.8	45.0	45.9	47.0	47.9	52.2	55.2	56.5	57.8	60.5	63.0	66.3	67.2	67.2	68.5	67.3	68.9	70.2	71.1	71.4						
8-year	38.5	42.9	43.4	45.1	46.5	50.5	53.0	55.2	56.3	58.9	61.3	65.1	65.7	66.6	67.1	66.2	67.4	69.1	70.2							
9-year	36.6	40.8	41.5	43.5	45.3	49.3	51.5	53.4	54.9	57.5	60.0	64.0	64.8	64.9	65.2	65.2	66.2	68.0								
10-year	35.0	38.9	39.8	42.1	44.4	48.1	50.1	52.2	53.7	56.2	58.9	63.2	63.3	63.8	63.5	63.8	65.3									
11-year	33.5	37.4	38.5	40.6	43.4	46.5	48.8	51.1	52.6	54.8	58.1	61.8	62.2	63.0	63.2	63.1										
12-year	32.4	36.0	36.9	39.6	42.3	45.7	48.2	50.2	51.3	53.6	56.9	60.1	61.6	61.7	62.3											
13-year	31.4	34.8	35.8	38.3	41.6	44.8	46.9	49.2	50.0	52.0	56.1	59.4	60.3	60.8												
14-year	30.2	33.5	34.7	37.2	40.7	44.3	45.7	48.7	48.9	51.3	54.9	58.3	59.6													
15-year	28.9	32.2	33.5	36.6	39.9	43.7	44.7	47.5	47.9	50.5	54.1	58.2														
16-year	28.0	31.2	32.8	35.7	39.4	43.6	43.4	47.3	47.2	49.2	53.6															
17-year	27.4	30.4	32.1	34.9	38.9	42.8	42.8	46.9	45.9	48.1																
18-year	26.8	29.6	31.4	34.1	37.9	42.0	42.0	46.3	45.4																	
19-year	26.1	28.8	30.6	33.3	37.1	41.3	41.3	45.8																		
20-year	25.2	28.0	30.1	32.9	35.9	40.9	40.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 19.10
Non-Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	69.4	72.7	68.9	67.0	66.8	71.1	74.6	74.8	76.0	76.5	77.5	80.5	79.7	80.1	81.0	81.0	81.5	83.7	82.6	83.7	83.8	84.0	83.4	83.1	83.7	84.5
2-year	59.4	63.7	60.5	58.3	58.5	63.5	67.1	68.3	69.2	70.3	71.3	75.3	74.9	75.5	76.2	76.5	77.9	78.0	78.1	78.8	79.9	79.3	78.8	79.4	79.6	
3-year	53.4	58.0	55.7	53.7	54.5	59.7	63.8	64.6	64.3	66.8	68.0	72.0	72.6	72.6	73.2	73.6	75.5	75.2	75.8	76.6	77.6	76.9	76.7	77.6		
4-year	49.4	54.0	51.6	50.4	51.6	56.3	61.0	62.0	61.5	63.9	65.7	70.1	70.9	70.2	71.3	71.3	73.4	73.6	73.7	74.4	75.7	75.9	75.1			
5-year	45.4	50.1	48.2	47.6	48.6	53.6	58.1	59.2	59.0	61.8	63.6	68.0	68.9	68.9	69.9	68.9	71.2	72.3	72.4	73.5	74.3	74.5				
6-year	42.6	46.6	45.5	45.6	46.9	51.0	56.0	56.5	56.3	60.3	61.9	66.8	67.0	67.5	68.4	67.1	69.8	71.6	70.7	71.7	72.5					
7-year	39.9	44.2	43.2	43.7	44.9	49.1	54.3	54.5	54.6	58.7	60.1	65.2	65.8	65.5	66.9	65.1	68.3	69.7	69.8	70.7						
8-year	37.8	42.0	41.0	41.8	43.3	47.4	52.1	53.3	53.6	56.8	58.3	63.7	64.5	65.0	65.0	63.4	66.6	68.8	69.2							
9-year	35.7	40.1	39.1	40.2	42.4	46.5	50.4	51.4	52.3	55.3	57.2	62.8	63.4	63.2	63.6	61.9	65.3	67.7								
10-year	34.1	38.0	37.7	39.2	41.7	45.4	48.8	50.4	51.5	54.2	56.0	61.8	61.8	62.0	61.4	60.9	64.4									
11-year	32.8	36.4	36.4	38.1	40.8	43.6	47.7	49.5	50.8	52.7	54.7	60.3	60.6	61.2	61.3	60.0										
12-year	31.9	35.2	35.1	37.3	39.5	42.6	47.2	48.2	49.8	51.5	53.7	58.9	60.1	59.6	60.3											
13-year	30.8	34.2	34.2	36.1	39.0	41.8	45.6	46.8	49.1	50.2	53.1	58.3	59.2	58.8												
14-year	29.5	33.1	33.2	35.0	37.8	41.4	44.4	46.2	48.1	49.7	52.1	57.2	58.1													
15-year	28.2	32.0	32.4	34.4	36.9	40.8	43.5	45.4	47.3	48.9	51.3	57.1														
16-year	27.2	30.8	31.6	33.4	36.2	40.8	42.6	45.3	47.1	47.5	50.6															
17-year	26.6	30.0	30.8	32.6	35.9	40.3	42.0	44.9	46.3	46.8																
18-year	26.2	29.4	30.2	32.0	35.2	39.5	41.0	44.3	45.7																	
19-year	25.6	28.8	29.4	31.3	34.3	39.0	40.3	43.8																		
20-year	25.1	28.0	29.1	31.0	33.3	38.8	40.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 19.11
Non-Hodgkin Lymphoma

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	71.0	73.3	74.0	74.7	75.2	75.7	76.7	78.6	79.3	78.1	80.0	81.3	81.0	82.0	82.4	80.9	82.0	84.6	85.2	85.1	83.1	84.3	85.7	83.9	85.9	85.6
2-year	61.1	63.7	65.6	67.2	66.5	68.7	70.1	72.1	73.4	72.6	75.7	76.9	76.5	76.5	78.6	77.4	77.2	80.1	81.2	80.7	79.7	80.9	81.8	80.2	82.4	
3-year	55.9	59.0	61.2	62.8	61.6	64.9	66.2	68.4	70.1	69.2	72.7	74.4	73.6	74.5	76.8	75.6	74.7	77.4	79.2	78.6	77.6	79.2	79.8	79.0		
4-year	51.9	55.3	57.7	59.5	59.0	61.9	63.4	65.5	68.2	67.4	71.0	72.5	72.4	72.9	75.1	73.9	73.0	75.6	77.8	76.5	75.7	78.0	78.4			
5-year	48.3	51.4	54.4	56.7	56.7	59.1	60.7	62.9	65.9	66.3	68.8	70.7	71.2	71.5	73.5	72.3	71.6	73.5	76.3	75.2	74.3	76.1				
6-year	44.7	48.5	51.7	53.9	54.0	58.0	58.4	60.4	63.5	64.2	67.9	69.5	69.8	70.0	72.6	71.0	70.9	72.1	73.9	73.3	72.0					
7-year	41.8	45.9	49.1	51.2	52.0	56.0	56.3	58.6	61.5	62.6	66.4	67.4	68.7	69.2	70.4	69.9	69.6	70.8	72.7	72.2						
8-year	39.2	43.8	46.3	49.2	50.5	54.2	54.1	57.3	59.6	61.5	65.0	66.8	67.1	68.4	69.5	69.3	68.5	69.5	71.5							
9-year	37.5	41.6	44.3	47.6	49.2	52.9	52.8	55.8	57.9	60.3	63.3	65.4	66.4	66.8	67.1	68.9	67.3	68.5								
10-year	36.0	39.8	42.4	45.8	48.0	51.4	51.7	54.2	56.2	58.7	62.5	64.9	65.1	65.7	65.9	67.1	66.2									
11-year	34.2	38.5	40.9	43.8	46.9	50.2	50.0	53.0	54.8	57.2	62.2	63.6	64.0	65.1	65.4	66.5										
12-year	32.9	36.8	39.1	42.5	46.0	49.5	49.4	52.4	53.0	56.0	60.7	61.3	63.2	64.1	64.6											
13-year	32.1	35.3	37.5	41.1	45.0	48.5	48.5	51.8	51.1	54.2	59.8	60.7	61.6	63.1												
14-year	31.0	34.0	36.5	40.0	44.3	47.9	47.1	51.4	49.8	53.2	58.4	59.6	61.2													
15-year	29.7	32.5	35.0	39.3	43.7	47.2	46.2	49.9	48.6	52.4	57.6	59.2														
16-year	28.8	31.7	34.3	38.6	43.6	46.7	44.3	49.4	47.0	51.2	57.1															
17-year	28.2	30.6	33.6	37.8	42.8	45.9	43.9	49.1	45.4	49.7																
18-year	27.4	29.9	32.7	36.8	41.4	45.0	43.0	48.6	44.8																	
19-year	26.5	28.8	32.1	35.9	40.7	44.0	42.4	48.0																		
20-year	25.4	28.0	31.2	35.4	39.4	43.4	42.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 20.12
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	79.3	79.1	80.4	81.0	82.7	81.7	83.2	80.7	83.7	84.3	82.4	83.3	84.0	84.4	83.5	85.1	84.9	86.2	85.8	87.2	87.2	85.7	86.0	87.0	87.0	87.4
2-year	65.9	65.9	67.5	69.4	71.6	69.6	71.1	70.2	72.3	73.5	72.9	72.8	75.0	75.3	74.0	75.5	76.3	77.5	77.4	78.7	78.8	78.1	78.1	79.1	79.6	
3-year	59.8	59.5	61.1	63.2	65.4	63.5	66.2	64.3	65.6	68.7	67.1	67.5	70.4	70.7	68.8	70.6	71.3	71.9	73.1	74.4	74.5	73.1	72.7	74.9		
4-year	55.9	55.5	57.1	59.0	61.6	60.0	62.1	60.2	62.2	64.9	62.6	64.1	67.1	66.9	66.0	67.4	68.4	68.5	69.7	71.3	71.9	70.8	69.4			
5-year	52.8	52.3	54.0	56.1	58.5	57.6	58.5	57.1	58.9	62.0	60.0	61.7	65.2	64.9	64.0	65.1	66.5	66.5	67.3	69.0	69.9	67.6				
6-year	50.5	49.6	51.5	53.6	56.0	55.2	56.0	55.7	56.3	59.9	57.9	60.0	63.9	62.2	61.8	62.9	65.1	64.5	65.4	67.3	68.3					
7-year	48.4	47.5	49.0	51.6	53.5	53.3	54.3	54.0	53.9	57.9	56.1	58.5	62.4	60.5	60.1	61.5	62.9	62.7	64.0	65.6						
8-year	46.2	45.9	47.0	49.4	51.3	51.1	53.2	52.9	52.1	55.6	53.9	56.2	60.8	59.3	58.4	59.5	61.5	60.7	62.4							
9-year	44.3	43.9	45.1	47.1	49.4	49.8	51.7	52.2	50.3	54.0	51.6	54.5	58.4	57.2	56.8	57.9	59.4	59.9								
10-year	42.3	42.2	43.3	45.4	47.8	47.6	49.9	51.1	49.0	52.3	49.5	52.6	57.5	55.5	54.9	56.3	58.7									
11-year	40.4	40.7	42.1	43.6	46.0	46.3	47.8	49.0	48.0	50.6	48.4	50.9	55.5	54.0	54.4	53.7										
12-year	39.1	39.3	40.5	42.4	44.1	45.3	46.1	47.9	46.7	48.9	46.6	50.3	53.1	52.9	52.6											
13-year	37.7	37.6	39.0	40.8	42.5	43.1	44.6	46.2	46.0	47.9	46.1	49.9	51.7	51.4												
14-year	36.7	36.5	37.7	39.6	40.0	41.4	43.8	44.4	44.7	46.6	45.4	48.4	50.6													
15-year	35.3	35.0	36.6	38.6	38.6	40.0	42.5	43.8	43.1	45.8	44.3	47.9														
16-year	33.9	33.8	35.5	37.5	37.6	39.0	41.3	42.6	42.4	44.6	43.1															
17-year	32.9	32.8	34.5	36.4	36.6	38.1	40.4	41.0	40.9	43.4																
18-year	31.9	31.7	33.1	35.3	35.2	37.2	39.2	40.0	40.4																	
19-year	31.1	30.2	32.0	34.3	34.6	35.7	37.6	39.1																		
20-year	29.9	29.7	31.4	33.5	33.9	34.9	35.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 20.13
Cancer of the Oral Cavity and Pharynx For Selected Subsites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Survival Time																											
<u>Cancer of the Tongue</u>																											
1-year	74.2	73.4	76.7	79.4	79.3	79.2	81.6	79.9	80.9	84.0	81.0	80.9	84.3	86.8	84.8	84.3	86.0	86.6	87.2	86.5	88.0	87.8	87.2	87.7	87.1	88.0	
2-year	54.8	57.5	60.5	64.7	65.2	66.7	71.0	65.8	68.2	71.0	70.4	69.5	71.4	75.4	72.2	73.9	76.5	76.9	79.0	76.7	78.6	80.8	78.4	79.8	79.3		
3-year	47.4	50.8	54.4	57.4	58.6	60.4	65.1	59.1	60.3	65.8	63.8	64.0	66.9	70.9	65.1	69.3	70.6	72.0	73.8	72.0	74.7	74.9	71.9	75.0			
4-year	43.5	47.4	49.9	53.6	56.4	57.4	62.1	55.5	57.9	63.1	61.0	61.7	64.2	66.5	61.7	66.9	68.9	69.7	71.1	69.5	72.1	73.4	68.8				
5-year	40.1	44.6	47.4	50.8	54.0	55.5	58.0	52.8	54.7	61.8	58.4	59.9	61.5	64.8	60.2	65.2	66.9	67.4	68.7	67.5	71.5	70.8					
6-year	38.4	41.8	45.1	48.5	50.4	52.9	56.4	52.3	52.8	59.8	56.3	57.8	61.0	61.6	59.2	63.8	65.6	66.0	67.0	66.5	70.1						
7-year	37.2	38.9	42.4	46.7	48.4	51.3	55.3	49.6	50.5	57.6	54.3	56.8	59.5	59.1	57.8	63.0	63.4	65.0	66.0	65.0							
8-year	35.7	37.4	40.5	44.3	46.5	49.9	53.8	48.0	50.4	55.5	50.6	53.6	58.9	56.9	56.1	60.3	61.8	63.1	64.5								
9-year	34.1	34.8	38.9	42.0	44.6	48.7	52.9	47.4	48.5	54.5	48.1	52.3	57.0	55.0	55.0	58.6	59.9	62.7									
10-year	32.5	33.3	37.5	40.0	43.3	46.7	52.3	45.9	47.7	53.2	46.2	50.0	56.2	53.9	52.4	56.8	58.5										
<u>Cancer of the Lip</u>																											
1-year	99.1	97.5	99.2	100.0	99.9	98.1	100.0	99.3	97.6	98.1	95.1	99.3	99.3	99.6	98.2	96.9	96.1	100.0	93.7	100.0	98.5	95.6	99.8	100.0	100.0	98.9	
2-year	97.4	95.9	98.0	98.7	99.4	94.0	97.0	97.9	95.1	95.7	93.8	99.3	99.3	99.6	95.3	94.5	96.1	98.7	92.2	99.6	98.5	91.7	98.5	99.7	97.6		
3-year	96.9	93.5	97.2	98.4	98.7	92.8	95.8	95.1	93.6	95.1	93.8	97.1	98.2	98.7	95.2	93.2	95.7	98.2	91.7	94.8	98.5	89.7	95.6	99.1			
4-year	95.7	92.2	96.9	97.2	96.2	91.6	93.4	93.2	91.8	92.6	90.5	97.1	98.2	95.7	95.2	90.0	94.6	94.0	90.5	94.8	98.5	88.9	91.6				
5-year	94.1	90.0	95.9	95.8	95.0	88.1	91.2	90.8	90.5	90.7	90.5	93.8	98.2	94.2	95.2	89.0	94.6	94.0	87.8	93.7	98.5	87.0					
6-year	92.5	87.6	94.8	94.5	95.0	87.4	89.4	89.6	89.3	90.6	87.8	93.5	98.2	94.2	95.2	85.9	94.6	91.9	86.2	92.3	97.5						
7-year	91.1	86.4	93.4	93.3	95.0	86.6	88.2	89.6	88.8	90.4	84.7	93.5	97.3	93.5	95.2	85.9	94.6	89.8	83.5	90.3							
8-year	89.9	86.1	91.8	92.0	91.0	83.2	88.2	87.5	88.8	90.4	82.5	93.3	97.3	93.3	92.0	85.9	94.6	87.2	78.7								
9-year	87.4	85.0	91.2	89.8	86.0	82.5	88.0	86.8	88.5	89.1	79.0	90.6	96.1	93.3	89.5	85.9	92.0	84.4									
10-year	85.7	83.6	91.1	87.9	82.0	82.5	85.5	86.8	88.3	88.3	76.3	87.7	95.4	92.2	89.5	85.9	92.0										
<u>Cancer of the Oropharynx and Tonsil</u>																											
1-year	70.3	70.8	72.9	74.4	80.3	79.4	81.1	81.3	86.9	82.2	83.7	85.2	82.7	83.9	84.2	85.2	85.8	86.1	88.6	90.1	90.7	86.6	87.0	89.7	87.7	90.4	
2-year	53.4	52.6	55.8	61.2	69.7	66.3	67.1	67.4	74.8	67.3	72.0	76.0	75.4	74.9	74.4	75.3	77.6	78.6	81.2	83.6	84.0	80.1	81.7	83.1	81.1		
3-year	44.5	44.4	45.7	53.7	62.7	60.6	61.8	60.2	67.2	62.3	66.7	71.4	71.4	71.2	70.7	70.4	72.2	73.3	76.5	80.0	79.7	75.6	77.5	78.8			
4-year	38.9	39.4	41.5	49.1	58.4	54.1	56.3	55.2	63.0	57.4	60.6	67.3	68.2	67.2	67.0	68.4	70.5	70.0	74.2	76.8	77.4	73.6	74.5				
5-year	35.4	35.3	37.8	44.9	53.9	52.6	51.3	53.2	59.7	53.6	58.7	64.4	65.2	64.6	65.5	65.6	68.0	67.4	72.6	74.9	75.4	70.1					
6-year	32.1	31.8	34.6	42.6	50.1	47.7	48.3	51.3	57.5	52.2	54.7	63.6	63.7	62.5	64.2	65.0	66.1	65.8	71.3	72.3	74.5						
7-year	30.2	29.7	31.9	41.3	46.6	43.7	47.3	50.1	55.7	48.8	52.8	62.7	62.8	61.7	61.7	62.9	63.5	62.5	70.2	70.8							
8-year	26.1	28.0	30.1	39.0	45.4	42.3	46.1	49.3	52.1	46.6	52.8	59.6	61.1	60.9	59.5	60.2	61.0	61.8	68.8								
9-year	23.7	25.8	28.3	36.6	44.1	41.3	44.3	48.8	51.9	45.0	49.6	59.0	60.7	57.6	57.8	58.8	58.9	60.2									
10-year	22.4	23.2	26.1	35.4	42.9	40.1	43.6	46.5	49.8	43.1	48.5	57.2	58.3	56.1	57.5	58.8	58.8										

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 20.14
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	78.6	78.5	79.5	80.4	82.6	81.4	82.4	80.3	84.1	84.1	82.9	83.4	84.2	84.9	84.0	84.9	85.9	86.5	86.3	87.1	88.0	86.9	85.4	86.8	87.2	87.8
2-year	65.0	65.0	66.0	68.2	70.6	68.9	69.6	70.1	71.2	72.9	73.2	71.6	74.9	75.1	74.2	74.8	77.0	78.1	77.8	78.2	78.5	79.1	77.0	78.9	79.7	
3-year	59.1	58.2	59.0	61.4	64.8	62.6	64.7	63.8	64.1	67.4	67.4	65.5	70.0	69.8	68.6	69.3	71.2	72.4	72.3	73.6	74.0	73.9	71.4	74.6		
4-year	54.7	54.1	54.8	57.1	60.6	58.8	60.0	59.3	60.5	63.9	62.3	62.4	66.7	65.5	65.8	66.1	67.7	68.6	68.6	70.0	70.9	71.4	67.9			
5-year	51.6	50.6	51.4	54.1	57.5	56.3	56.3	56.1	57.3	60.8	59.5	60.4	65.2	63.7	63.8	64.3	65.9	66.2	66.4	67.6	68.9	68.0				
6-year	49.4	48.0	49.2	51.7	54.8	53.4	53.9	54.7	54.6	58.3	57.2	58.5	64.1	61.2	61.1	62.3	64.3	63.9	64.4	66.2	67.3					
7-year	47.2	45.9	47.0	49.5	51.9	51.0	52.5	52.9	52.6	56.1	55.3	57.5	62.9	59.9	59.5	61.0	62.0	61.9	62.9	64.6						
8-year	45.2	44.4	44.7	47.6	49.5	49.1	51.5	51.4	51.1	53.8	52.8	55.3	61.1	58.2	57.3	58.9	60.6	60.2	61.3							
9-year	43.2	42.5	42.8	45.3	47.6	48.1	49.8	50.5	49.5	51.9	50.7	53.2	58.6	55.8	55.5	57.5	58.3	59.3								
10-year	41.2	40.6	40.8	43.8	45.8	45.9	48.2	49.6	48.5	50.3	48.8	51.1	57.4	54.1	53.5	56.4	57.6									
11-year	39.2	39.1	39.6	42.2	43.8	44.1	46.4	47.3	47.3	48.5	47.7	49.7	55.6	53.2	53.3	53.7										
12-year	38.0	37.9	38.3	40.9	42.1	43.3	44.5	46.5	46.4	46.6	46.1	49.0	53.2	52.3	51.4											
13-year	36.7	36.3	36.7	39.3	40.3	41.1	42.9	44.9	45.6	45.9	46.1	48.3	51.6	50.6												
14-year	35.8	35.3	35.6	38.0	37.0	40.0	41.5	43.2	43.9	44.5	45.3	47.0	50.8													
15-year	34.5	33.9	34.6	36.9	35.0	38.3	40.2	42.6	42.3	43.7	44.2	46.3														
16-year	33.1	32.8	33.6	35.7	34.0	37.3	38.9	41.0	41.5	42.3	43.1															
17-year	32.0	31.7	32.6	34.7	33.0	36.8	38.0	39.2	39.4	40.8																
18-year	31.1	30.6	31.3	33.5	31.5	35.7	37.0	38.1	38.8																	
19-year	30.5	28.8	29.9	32.7	31.0	33.6	35.7	37.1																		
20-year	29.4	28.3	29.2	31.8	30.2	33.0	33.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 20.15
Cancer of the Oral Cavity and Pharynx (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	80.8	80.2	82.2	82.2	82.9	82.2	85.1	81.5	82.8	84.8	81.7	83.0	83.5	83.3	82.4	85.4	82.4	85.3	84.4	87.3	85.1	83.0	87.4	87.6	86.4	86.4
2-year	67.8	67.7	70.5	72.0	73.3	71.1	74.5	70.2	74.4	74.8	72.5	75.3	75.4	75.8	73.4	77.2	74.7	76.2	76.6	79.8	79.6	75.9	80.9	79.4	79.5	
3-year	61.6	62.3	65.4	67.0	66.5	65.3	69.5	65.4	68.6	71.4	66.4	71.7	71.0	72.7	69.2	73.8	71.5	70.8	74.8	76.3	75.7	71.4	76.1	75.9		
4-year	58.5	58.6	61.9	63.1	63.4	62.5	66.9	62.1	65.6	67.1	63.1	67.9	68.1	70.1	66.3	70.4	69.6	68.3	72.4	74.5	74.3	69.5	73.1			
5-year	55.4	56.0	59.2	60.3	60.4	60.2	63.4	59.2	62.1	64.7	60.9	64.6	65.0	67.6	64.3	67.0	67.8	67.1	69.5	72.4	72.4	66.5				
6-year	53.0	52.9	56.3	57.8	58.3	58.8	60.8	57.7	59.7	63.3	59.4	63.0	63.3	64.4	63.1	64.4	66.8	66.0	67.7	69.9	70.7					
7-year	51.0	50.8	53.2	56.0	56.6	57.9	58.1	56.4	56.7	61.6	57.9	60.5	61.3	61.8	61.5	62.4	65.1	64.6	66.8	68.3						
8-year	48.3	49.2	51.6	53.2	54.6	55.0	56.8	55.7	54.1	59.7	56.3	58.2	60.1	61.8	60.5	60.8	63.2	61.9	65.0							
9-year	46.8	46.9	49.7	51.0	52.7	53.1	55.9	55.7	52.1	58.6	53.5	57.1	58.1	60.3	59.4	58.8	61.6	61.4								
10-year	44.7	45.7	48.5	48.8	51.5	51.1	53.5	54.4	50.0	56.8	50.8	55.7	57.7	58.6	58.2	56.2	61.0									
11-year	42.9	44.2	47.0	46.6	50.0	50.5	50.9	52.3	49.3	55.2	49.8	53.4	55.4	55.8	56.8	53.7										
12-year	41.4	42.4	45.1	45.7	47.9	49.3	49.5	50.7	47.2	54.1	47.5	53.1	52.9	54.1	55.0											
13-year	39.9	40.2	43.5	44.1	46.6	47.1	48.4	49.0	46.8	52.4	46.3	53.1	51.9	53.0												
14-year	38.6	39.1	42.0	43.0	45.4	44.2	48.4	46.9	46.2	50.8	45.7	51.6	49.8													
15-year	37.0	37.2	40.7	42.0	44.9	43.5	47.5	46.2	44.5	50.4	44.4	51.1														
16-year	35.8	36.0	39.4	41.2	44.0	42.5	46.4	45.9	44.2	49.2	43.0															
17-year	34.9	35.1	38.4	39.9	42.9	40.9	45.5	44.5	43.8	48.9																
18-year	33.6	34.2	36.8	39.1	41.8	40.2	43.9	43.8	43.6																	
19-year	32.6	33.0	36.0	37.5	41.1	39.8	41.8	42.9																		
20-year	31.3	32.5	35.7	36.9	40.4	38.6	39.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 21.9
Cancer of the Ovary (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	64.0	69.4	69.4	71.9	73.1	73.7	76.6	75.0	73.9	76.8	76.1	75.3	76.0	75.3	75.1	75.3	75.4	75.8	76.2	76.9	78.4	78.4	78.1	81.1	78.2	79.4
2-year	48.9	53.2	54.1	58.7	60.2	61.4	65.2	64.2	61.6	63.6	65.2	63.6	66.0	64.8	64.6	64.1	64.2	64.9	67.3	67.1	69.4	67.2	68.0	69.6	69.0	
3-year	42.2	45.8	46.4	49.9	51.2	52.9	56.6	55.1	53.0	54.5	56.0	54.4	58.3	56.6	56.1	55.8	55.9	56.5	58.4	59.3	60.7	59.7	61.0	61.8		
4-year	38.6	41.5	41.5	44.5	46.2	46.2	48.7	50.3	47.6	47.7	50.4	48.0	50.0	49.9	50.1	49.8	49.3	50.7	52.8	52.5	54.2	53.3	55.6			
5-year	36.5	38.8	38.2	40.9	42.4	42.1	44.0	45.3	42.8	43.1	45.8	42.8	44.5	44.4	44.6	45.7	44.6	46.8	47.1	47.2	49.5	48.8				
6-year	35.2	37.0	36.0	38.3	39.0	38.8	40.4	42.4	40.0	40.1	42.9	39.5	41.2	41.5	41.3	42.6	42.0	44.1	44.1	43.9	46.6					
7-year	33.8	36.1	34.8	36.8	37.0	37.3	38.2	40.6	38.0	37.6	40.4	37.6	38.4	39.5	39.4	40.6	40.6	41.7	41.8	41.1						
8-year	33.0	34.6	33.8	35.7	35.2	36.2	36.7	38.6	36.4	36.0	38.1	35.8	37.1	37.7	38.2	38.9	38.8	39.4	39.9							
9-year	32.4	33.7	33.0	34.5	33.7	35.4	35.3	36.9	34.9	34.6	36.3	34.9	35.6	36.2	37.3	37.9	37.1	37.6								
10-year	31.9	32.9	32.4	33.7	32.8	34.4	34.2	35.9	34.1	33.3	35.2	34.1	34.3	35.2	36.2	37.4	36.0									
11-year	31.6	32.4	32.1	33.0	32.1	33.7	33.0	35.1	32.6	32.6	34.5	33.4	33.7	34.3	35.8	36.7										
12-year	30.9	31.8	31.4	32.4	32.0	33.2	32.2	34.6	31.8	32.0	33.3	33.0	33.0	33.5	34.7											
13-year	30.7	31.2	30.8	32.0	31.5	32.6	31.7	33.3	31.2	31.4	32.8	32.2	32.4	33.0												
14-year	30.2	30.8	30.4	31.8	31.5	31.8	31.4	32.6	30.2	31.1	32.6	31.5	32.3													
15-year	29.6	30.6	30.3	31.3	30.9	31.4	31.4	32.0	30.2	30.4	32.0	30.9														
16-year	29.3	30.4	30.3	31.1	30.7	30.8	30.6	31.4	29.3	30.4	32.0															
17-year	28.9	30.0	30.1	30.8	30.5	30.6	30.2	31.1	29.2	30.3																
18-year	28.8	29.8	29.9	30.5	30.1	30.6	30.0	30.7	28.5																	
19-year	28.5	29.5	29.9	30.1	29.6	30.4	29.9	30.5																		
20-year	28.3	29.2	29.6	29.9	29.0	29.9	29.8																			

^a Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
[Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 22.9
Cancer of the Pancreas (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Survival Time																											
1-year	14.3	15.8	17.0	18.5	17.8	19.0	19.9	19.3	22.3	21.7	23.8	25.0	22.8	22.8	26.7	25.8	28.0	29.8	29.8	30.3	31.5	34.8	35.6	35.0	36.2	37.3	
2-year	5.9	6.3	6.7	8.6	8.3	9.3	9.8	8.8	10.5	11.0	11.4	11.9	11.5	11.9	12.9	13.3	13.9	15.1	15.9	16.9	16.3	19.8	20.6	19.7	20.4		
3-year	3.6	4.1	4.7	5.8	5.3	6.2	7.1	5.6	6.6	7.3	7.4	8.5	7.6	8.3	9.1	9.5	9.8	10.5	11.9	12.0	11.7	14.8	15.6	15.2			
4-year	3.0	3.3	3.8	4.8	4.0	4.8	5.5	4.4	5.7	6.1	6.1	6.8	5.8	6.4	7.2	8.2	8.4	8.6	10.0	10.0	9.4	12.6	13.0				
5-year	2.5	2.8	3.3	4.2	3.6	4.2	4.9	3.8	5.1	5.2	5.1	6.2	5.1	5.5	6.4	7.6	7.7	7.8	8.7	8.4	8.2	11.5					
6-year	2.3	2.5	2.9	4.0	3.2	3.8	4.4	3.7	4.6	4.7	4.7	5.7	4.6	5.0	5.8	7.0	7.1	7.2	8.2	7.6	7.6						
7-year	2.2	2.3	2.7	3.6	3.1	3.5	4.3	3.4	4.3	4.3	4.2	5.0	4.3	4.7	5.5	6.2	6.5	6.6	7.5	7.1							
8-year	1.9	2.1	2.5	3.4	2.7	3.4	4.1	3.0	4.0	3.9	3.9	4.7	4.1	4.4	5.1	6.0	5.8	6.1	7.0								
9-year	1.8	2.0	2.2	3.2	2.7	3.3	4.0	2.7	3.8	3.8	3.8	4.6	3.8	4.2	4.8	5.9	5.3	6.0									
10-year	1.8	1.9	2.2	3.0	2.5	3.2	3.9	2.6	3.6	3.7	3.5	4.5	3.5	4.1	4.4	5.7	5.0										
11-year	1.7	1.7	2.1	2.8	2.5	2.9	3.8	2.5	3.3	3.6	3.3	4.1	3.3	4.1	4.0	5.5											
12-year	1.7	1.7	2.1	2.7	2.3	2.8	3.6	2.4	3.2	3.3	3.2	4.0	3.3	4.1	3.9												
13-year	1.6	1.6	2.0	2.6	2.2	2.7	3.3	2.2	3.1	3.1	3.2	3.8	3.1	3.9													
14-year	1.6	1.6	1.9	2.5	1.9	2.7	3.1	1.9	3.1	2.8	3.0	3.8	2.9														
15-year	1.5	1.5	1.8	2.4	1.8	2.7	3.0	1.9	3.0	2.7	2.7	3.5															
16-year	1.5	1.4	1.7	2.2	1.7	2.2	2.9	1.8	2.7	2.6	2.5																
17-year	1.4	1.4	1.7	2.2	1.5	1.9	2.7	1.5	2.7	2.5																	
18-year	1.3	1.3	1.7	2.2	1.4	1.9	2.7	1.3	2.4																		
19-year	1.3	1.3	1.6	2.1	1.4	1.9	2.7	1.2																			
20-year	1.2	1.3	1.6	2.0	1.3	1.9	2.7																				

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 22.10
Cancer of the Pancreas (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	14.3	14.9	16.7	17.8	16.9	19.8	21.2	18.8	22.8	21.9	23.2	24.8	24.3	25.2	25.7	27.0	28.2	31.4	29.3	29.9	32.7	34.5	36.0	36.1	35.6	37.0
2-year	5.8	5.8	6.2	8.4	7.3	10.9	9.9	9.1	9.8	11.4	12.0	11.1	11.6	12.6	11.9	13.7	13.6	16.2	16.6	16.6	17.8	19.7	20.8	20.3	20.2	
3-year	3.5	3.8	4.4	5.8	4.2	7.1	7.0	5.6	6.0	7.0	8.0	8.2	7.6	8.2	7.9	9.8	9.1	11.5	12.8	12.1	12.7	14.6	15.6	16.1		
4-year	3.0	3.0	3.3	4.6	3.5	5.8	5.5	4.5	5.5	5.9	6.4	6.9	5.6	5.9	5.6	8.2	7.8	9.9	10.9	10.1	10.5	12.6	13.1			
5-year	2.6	2.5	2.9	4.0	3.0	5.1	4.7	3.9	4.9	5.2	5.5	6.1	4.6	5.1	5.2	7.6	7.2	9.0	9.5	8.5	9.2	11.5				
6-year	2.4	2.2	2.4	3.8	2.7	4.7	3.9	3.7	4.0	4.9	5.0	5.3	4.2	4.5	4.7	7.1	6.4	8.5	8.9	7.5	8.6					
7-year	2.3	1.9	2.3	3.4	2.7	4.2	3.4	3.4	3.9	4.5	4.6	4.8	4.0	4.1	4.3	6.1	6.0	7.6	8.2	6.9						
8-year	2.0	1.7	2.1	3.1	2.4	4.2	3.4	3.2	3.8	3.9	4.2	4.4	3.8	3.8	4.0	6.0	5.2	6.9	7.7							
9-year	1.9	1.6	1.8	3.0	2.2	4.0	3.3	2.8	3.6	3.7	4.1	4.3	3.4	3.6	3.7	5.9	4.8	6.9								
10-year	1.8	1.5	1.7	2.7	2.1	3.8	3.2	2.8	3.6	3.5	3.9	4.2	2.8	3.5	3.4	5.6	4.4									
11-year	1.7	1.5	1.6	2.6	2.0	3.2	2.9	2.8	3.2	3.5	3.7	3.6	2.6	3.5	2.9	5.3										
12-year	1.6	1.5	1.6	2.6	1.8	3.0	2.8	2.6	3.1	2.9	3.4	3.5	2.6	3.4	2.7											
13-year	1.6	1.4	1.5	2.5	1.7	3.0	2.6	2.5	3.0	2.7	3.4	3.5	2.2	3.2												
14-year	1.6	1.4	1.5	2.4	1.2	2.9	2.4	2.3	3.0	2.4	3.4	3.5	2.1													
15-year	1.6	1.3	1.3	2.3	1.2	2.8	2.2	2.3	3.0	2.3	3.3	2.9														
16-year	1.6	1.3	1.3	2.1	1.2	2.0	1.9	2.3	2.6	2.3	3.0															
17-year	1.4	1.3	1.3	2.1	0.8	1.7	1.9	1.9	2.6	2.3																
18-year	1.3	1.3	1.2	2.1	0.7	1.7	1.9	1.4	2.2																	
19-year	1.2	1.3	1.2	2.1	0.7	1.6	1.8	1.4																		
20-year	1.1	1.2	1.2	1.9	0.2	1.6	1.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 22.11
Cancer of the Pancreas (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

Survival Time	1975-	1980-	1985-	1990-	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
	1979	1984	1989	1994																							
1-year	14.3	16.6	17.3	19.1	18.6	18.2	18.7	19.8	21.9	21.6	24.3	25.1	21.5	20.4	27.9	24.7	27.7	28.2	30.2	30.8	30.1	35.2	35.2	33.8	36.8	37.5	
2-year	5.9	6.7	7.2	8.7	9.2	7.8	9.8	8.5	11.1	10.5	10.8	12.7	11.3	11.2	14.0	12.9	14.2	14.0	15.2	17.2	14.6	20.0	20.4	19.2	20.6		
3-year	3.8	4.4	5.1	5.8	6.2	5.4	7.2	5.7	7.1	7.6	6.7	8.7	7.6	8.4	10.3	9.2	10.5	9.5	11.0	11.9	10.6	15.0	15.6	14.4			
4-year	3.0	3.6	4.3	4.9	4.4	3.9	5.5	4.3	5.8	6.3	5.8	6.7	6.0	7.0	8.8	8.1	8.9	7.3	9.1	10.0	8.2	12.5	12.9				
5-year	2.4	3.2	3.7	4.4	4.2	3.4	5.2	3.7	5.3	5.2	4.6	6.3	5.5	6.0	7.6	7.6	8.3	6.5	7.9	8.3	7.1	11.6					
6-year	2.3	2.8	3.4	4.1	3.7	3.0	4.9	3.7	5.1	4.5	4.3	6.0	5.0	5.6	7.0	7.0	7.8	6.0	7.4	7.6	6.6						
7-year	2.0	2.7	3.2	3.8	3.4	2.8	4.9	3.4	4.7	4.1	3.7	5.2	4.6	5.3	6.7	6.3	7.0	5.6	6.8	7.3							
8-year	1.8	2.5	2.8	3.7	3.0	2.6	4.8	2.8	4.1	3.9	3.5	5.0	4.4	5.0	6.2	6.0	6.4	5.2	6.3								
9-year	1.8	2.3	2.7	3.4	3.0	2.5	4.6	2.5	3.9	3.8	3.4	4.8	4.2	4.8	5.9	5.9	5.9	5.0									
10-year	1.7	2.2	2.6	3.2	2.9	2.5	4.6	2.4	3.7	3.7	3.1	4.7	4.1	4.7	5.4	5.8	5.7										
11-year	1.7	2.0	2.5	2.9	2.9	2.5	4.6	2.1	3.4	3.7	3.0	4.5	3.9	4.7	5.1	5.7											
12-year	1.7	1.9	2.5	2.7	2.8	2.5	4.3	2.1	3.2	3.7	2.9	4.4	3.8	4.7	5.0												
13-year	1.6	1.8	2.4	2.7	2.6	2.3	3.8	1.9	3.2	3.5	2.8	4.1	3.8	4.7													
14-year	1.5	1.7	2.4	2.6	2.4	2.3	3.8	1.4	3.1	3.2	2.4	4.1	3.7														
15-year	1.3	1.6	2.3	2.3	2.2	2.3	3.7	1.4	3.0	3.0	2.1	4.0															
16-year	1.3	1.5	2.0	2.3	2.1	2.3	3.7	1.0	2.8	2.8	2.1																
17-year	1.3	1.5	2.0	2.3	2.1	2.1	3.5	1.0	2.7	2.7																	
18-year	1.3	1.4	2.0	2.2	2.0	2.1	3.5	1.0	2.6																		
19-year	1.3	1.4	2.0	2.2	2.0	2.1	3.5	0.8																			
20-year	1.3	1.4	2.0	2.1	2.0	2.1	3.5																				

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 23.9
Cancer of the Prostate (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	91.4	93.4	95.5	98.8	98.7	99.1	99.2	99.5	99.6	99.7	99.9	99.8	99.7	99.8	99.8	99.8	99.8	99.7	99.6	99.6	99.7	99.3	99.4	98.7	99.2	99.3
2-year	84.6	87.4	91.0	97.3	97.7	98.0	98.8	98.9	99.6	99.5	99.9	99.8	99.6	99.7	99.6	99.8	99.8	99.6	99.4	99.5	99.4	98.7	98.8	98.0	98.2	
3-year	78.4	81.5	86.9	96.0	97.0	97.4	98.4	98.5	99.6	99.1	99.9	99.7	99.4	99.7	99.5	99.7	99.8	99.3	99.3	99.5	99.3	98.4	98.3	97.5		
4-year	73.3	76.5	83.6	94.9	96.4	96.9	97.9	98.2	99.2	98.9	99.8	99.7	99.4	99.7	99.5	99.6	99.8	99.2	99.3	99.5	99.3	98.2	97.9			
5-year	68.8	71.9	80.4	93.7	95.7	96.4	97.4	98.1	99.2	98.8	99.8	99.7	99.1	99.7	99.1	99.6	99.8	99.2	99.3	99.5	99.2	98.1				
6-year	65.3	68.5	77.9	92.7	95.4	96.0	97.1	97.9	99.2	98.5	99.8	99.7	99.1	99.7	99.1	99.6	99.8	99.2	99.3	99.5	99.2					
7-year	61.8	65.0	75.4	91.6	95.0	95.4	97.1	97.7	99.1	98.5	99.8	99.7	99.1	99.7	99.1	99.6	99.8	99.2	99.3	99.5						
8-year	58.6	62.0	73.4	90.6	94.6	95.0	97.0	97.7	98.7	98.2	99.8	99.7	99.0	99.7	99.1	99.6	99.8	99.2	99.3							
9-year	55.7	59.7	71.4	89.4	94.3	94.4	96.6	97.0	98.5	98.2	99.8	99.7	98.5	99.6	99.1	99.6	99.8	99.2								
10-year	53.2	57.7	69.6	88.5	93.5	94.3	96.5	96.6	98.1	98.2	99.8	99.7	98.5	99.4	99.1	99.6	99.8									
11-year	50.7	55.5	67.9	87.5	92.8	93.9	96.4	96.2	97.9	97.3	99.8	99.7	98.3	99.2	99.1	99.6										
12-year	48.5	53.6	66.1	86.6	92.0	93.5	96.1	95.5	97.4	97.2	99.8	99.7	97.9	98.8	99.1											
13-year	46.5	51.6	64.3	85.9	91.0	93.0	95.3	95.3	97.2	96.6	99.8	99.7	97.9	98.8												
14-year	44.8	50.0	63.0	85.1	90.1	92.8	94.7	95.0	96.6	96.1	99.8	99.7	97.8													
15-year	43.6	48.7	61.5	84.2	89.4	92.1	94.7	94.9	96.0	95.4	99.8	99.7														
16-year	41.9	46.9	60.0	83.3	88.9	91.2	94.1	94.4	95.6	95.4	99.3															
17-year	40.1	45.8	59.4	82.4	88.0	90.6	93.6	94.1	95.6	95.4																
18-year	38.8	44.5	58.3	81.8	87.7	90.4	92.9	94.1	95.2																	
19-year	38.0	43.1	57.9	80.8	87.3	89.7	92.6	94.1																		
20-year	37.0	41.7	56.9	80.4	85.8	88.8	92.6																			
21-year	35.4	40.8	56.1	79.5	84.9	88.8																				
22-year	34.0	39.5	54.8	78.7	84.3																					
23-year	33.2	38.3	53.6	77.7																						
24-year	32.4	37.4	52.5																							
25-year	31.0	35.9	51.4																							
26-year	30.4	34.7	50.4																							
27-year	29.7	34.4	49.5																							
28-year	29.1	34.0	48.8																							
29-year	28.5	33.3																								
30-year	27.0	32.4																								

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 24.9
Cancer of the Stomach (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																									
	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	38.6	41.5	43.8	44.8	45.9	45.4	45.3	47.6	46.7	47.0	48.6	49.6	50.8	53.0	49.9	52.2	52.1	55.6	55.6	54.4	58.2	57.5	59.4	60.5	58.5	60.0
2-year	24.9	26.7	29.1	31.3	31.5	31.9	32.4	33.6	34.0	32.6	33.3	35.3	38.4	38.4	36.4	40.0	39.5	41.8	40.6	40.4	43.5	45.2	44.8	45.9	44.3	
3-year	19.7	21.0	23.4	25.2	26.5	26.6	26.3	28.4	27.7	26.9	29.0	30.2	33.5	33.0	31.6	34.6	33.8	36.5	36.1	35.3	36.0	38.7	38.3	38.2		
4-year	17.3	18.3	20.6	22.5	23.3	23.8	23.4	25.2	24.8	24.7	27.5	27.1	30.5	30.1	28.0	33.3	31.2	34.0	33.7	32.1	32.9	35.9	34.4			
5-year	15.6	16.6	19.3	20.8	21.9	22.2	21.8	22.8	23.1	23.0	26.3	26.0	28.0	28.4	25.9	31.3	29.1	32.1	31.7	30.5	31.0	34.3				
6-year	14.6	15.8	18.3	19.3	20.7	20.5	20.7	21.9	21.1	22.2	25.3	24.5	26.6	26.7	24.8	29.9	27.2	31.0	30.6	29.6	29.6					
7-year	14.0	15.2	17.6	18.4	19.6	20.0	20.1	21.1	20.1	21.4	24.5	23.8	25.8	25.4	23.9	29.0	26.0	30.0	29.6	28.5						
8-year	13.4	14.5	17.0	17.5	19.4	19.7	19.4	20.6	19.0	20.7	22.8	22.5	24.6	24.5	23.4	28.4	25.1	29.3	29.0							
9-year	13.1	14.1	16.5	16.9	19.1	19.3	19.3	19.6	18.6	20.0	22.3	21.7	24.2	23.7	22.2	27.8	24.4	28.1								
10-year	12.7	13.9	15.8	16.1	18.2	18.8	18.3	18.7	18.0	19.5	21.6	21.0	23.6	22.5	21.5	26.4	23.9									
11-year	12.3	13.4	15.4	15.9	17.8	17.7	17.8	18.6	17.2	18.5	21.1	20.1	23.1	22.1	21.2	26.0										
12-year	12.0	12.9	15.0	15.6	17.2	17.1	17.7	17.8	17.0	18.0	20.5	19.4	22.5	21.1	20.3											
13-year	11.7	12.9	14.9	15.2	16.7	16.0	17.7	17.0	15.7	17.6	20.4	19.1	22.0	19.7												
14-year	11.2	12.8	14.6	14.9	16.4	15.7	17.3	17.0	15.6	17.5	20.0	18.1	21.5													
15-year	10.8	12.3	14.3	14.7	16.3	15.4	17.3	17.0	15.6	17.3	19.2	17.2														
16-year	10.5	11.9	14.0	14.1	16.3	14.7	16.6	16.6	14.9	17.2	18.4															
17-year	10.2	11.5	13.8	13.9	16.3	14.0	15.6	15.6	14.2	16.5																
18-year	9.8	11.5	13.6	13.3	16.1	13.3	15.2	15.4	13.8																	
19-year	9.6	11.3	13.2	13.1	15.7	13.0	15.2	14.2																		
20-year	9.2	10.8	13.0	12.8	14.8	12.7	15.2																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 24.10
Cancer of the Stomach (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	38.0	41.0	42.8	43.5	46.7	43.6	45.5	48.9	46.6	46.2	49.4	47.8	50.3	53.5	51.7	51.9	51.1	53.9	55.6	53.8	57.8	56.0	58.7	60.3	58.8	59.7
2-year	23.8	25.8	27.4	29.4	31.7	28.7	30.9	35.5	33.8	32.7	33.0	32.7	36.3	37.8	37.0	37.8	37.1	39.6	39.4	39.0	42.4	43.2	43.1	45.4	43.4	
3-year	18.5	20.0	21.4	23.4	25.9	23.6	24.9	29.5	27.4	27.2	28.6	27.1	31.6	31.1	31.5	33.1	31.7	33.2	34.2	33.9	34.9	36.6	36.9	36.6		
4-year	16.1	17.2	18.2	20.7	22.4	21.5	21.7	25.6	24.2	24.7	26.6	24.8	28.1	27.7	27.5	31.6	28.7	30.4	31.9	30.0	31.6	32.9	33.1			
5-year	14.4	15.4	16.7	18.8	20.8	20.2	19.9	23.2	22.2	22.4	25.6	23.4	25.5	26.4	25.3	29.8	26.8	29.0	29.4	27.7	29.3	30.6				
6-year	13.4	14.6	15.8	17.2	19.7	19.1	18.7	22.4	20.4	21.3	23.8	22.0	24.2	24.4	24.3	28.4	25.0	28.3	28.2	26.9	28.2					
7-year	12.7	13.9	15.0	16.2	18.7	18.8	17.8	21.3	18.8	20.3	23.3	21.1	23.8	23.5	23.7	27.5	23.4	27.5	27.2	26.1						
8-year	12.3	13.3	14.5	15.2	18.6	18.7	17.7	20.8	17.7	19.4	21.6	20.2	22.4	22.6	23.1	26.9	22.6	27.3	26.6							
9-year	12.2	13.0	13.9	14.6	18.5	17.9	17.4	20.1	16.8	18.2	21.4	19.9	22.2	21.7	21.8	26.2	22.1	25.8								
10-year	11.8	12.9	13.4	13.6	17.3	17.5	16.5	19.1	16.5	17.9	20.9	18.7	22.2	20.5	21.3	25.1	21.9									
11-year	11.4	12.5	12.9	13.4	17.1	15.7	15.9	19.1	15.9	16.5	20.6	17.7	22.2	19.9	20.5	24.9										
12-year	11.3	11.8	12.7	13.1	16.6	15.0	15.7	18.4	15.6	16.0	19.9	17.2	21.7	19.1	19.6											
13-year	11.1	11.8	12.7	12.7	16.0	14.8	15.7	17.7	14.6	15.5	19.6	16.5	21.2	17.9												
14-year	10.4	11.8	12.3	12.1	15.4	14.7	15.3	17.7	14.1	15.1	19.6	15.8	20.4													
15-year	10.0	11.3	12.0	11.5	15.4	14.7	15.2	17.7	14.1	14.6	18.6	14.2														
16-year	9.8	10.8	11.7	11.2	15.4	13.5	15.2	16.5	13.9	14.4	17.9															
17-year	9.5	10.5	11.4	11.1	15.3	12.2	13.6	15.7	13.1	14.4																
18-year	8.8	10.4	11.3	10.7	15.0	11.5	12.7	15.7	12.3																	
19-year	8.7	10.3	10.9	10.4	14.3	11.5	12.6	15.0																		
20-year	8.6	9.9	10.8	10.3	13.6	10.9	12.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 24.11
Cancer of the Stomach (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	39.7	42.4	45.4	46.9	44.6	48.5	45.0	45.7	46.9	48.4	47.2	52.3	51.7	52.3	46.9	52.7	53.8	58.5	55.7	55.1	58.7	59.9	60.5	60.9	58.1	60.4
2-year	26.7	28.2	31.9	34.4	31.4	37.3	34.8	30.9	34.3	32.5	33.8	39.3	41.5	39.5	35.2	43.5	43.5	45.6	42.4	42.5	45.3	48.2	47.4	46.8	45.9	
3-year	21.6	22.6	26.7	28.1	27.6	31.6	28.6	26.6	28.3	26.5	29.6	34.9	36.5	36.0	31.8	36.9	37.0	42.1	38.9	37.5	37.9	42.0	40.5	40.9		
4-year	19.2	20.1	24.6	25.5	24.7	27.4	26.3	24.6	25.7	24.9	28.9	30.5	34.1	34.0	29.0	35.6	35.2	40.0	36.3	35.3	35.0	40.4	36.3			
5-year	17.4	18.7	23.6	24.1	23.7	25.5	24.9	22.2	24.3	23.8	27.5	29.8	31.8	31.5	26.8	33.5	32.9	37.1	35.2	34.7	33.6	39.9				
6-year	16.6	18.0	22.5	22.8	22.4	22.8	24.0	21.0	22.1	23.6	27.2	28.0	30.3	30.1	25.5	32.3	30.6	35.6	34.2	33.5	32.0					
7-year	16.1	17.3	21.8	22.1	21.2	21.6	23.3	20.8	21.9	23.0	26.4	27.7	28.9	28.1	24.4	31.3	30.3	34.1	33.3	32.0						
8-year	15.2	16.3	21.3	21.3	20.7	21.0	22.1	20.2	20.9	22.7	24.7	25.9	27.9	27.4	23.9	30.5	29.0	32.4	32.3							
9-year	14.5	15.9	20.8	20.7	19.8	21.0	22.1	19.0	20.9	22.7	23.5	24.2	27.0	26.8	23.0	30.2	28.0	32.0								
10-year	14.3	15.5	19.8	20.2	19.6	20.9	21.1	17.8	19.9	22.0	22.8	24.1	25.2	25.9	21.5	28.3	27.0									
11-year	13.7	15.0	19.6	19.7	19.0	20.9	20.6	17.6	18.9	21.5	22.0	23.7	24.4	25.5	21.5	27.5										
12-year	13.2	14.6	19.0	19.7	18.2	20.5	20.5	16.9	18.9	21.1	21.4	22.6	23.6	24.2	21.3											
13-year	12.6	14.6	18.7	19.3	17.8	17.8	20.5	15.8	17.3	21.0	21.2	22.4	23.2	22.6												
14-year	12.4	14.4	18.5	19.3	17.8	17.3	20.1	15.7	17.3	21.0	20.7	21.4	23.1													
15-year	12.2	13.9	18.1	19.3	17.8	16.3	20.1	15.7	17.3	21.0	20.1	21.2														
16-year	11.7	13.6	17.8	18.9	17.8	16.2	18.7	15.7	16.1	21.0	18.9															
17-year	11.4	13.3	17.6	18.3	17.8	16.2	18.6	15.1	15.5	19.8																
18-year	11.3	13.2	17.5	17.6	17.8	16.1	18.6	14.8	15.5																	
19-year	10.9	12.8	17.2	17.4	17.8	15.5	18.6	12.7																		
20-year	10.2	12.4	16.4	16.9	17.0	15.4	18.6																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 25.9
Cancer of the Testis (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	93.2	95.8	97.9	97.7	98.3	97.6	97.3	97.9	96.9	97.9	98.5	98.2	98.4	97.9	98.1	98.9	98.9	98.7	98.5	98.1	97.6	99.1	97.8	97.7	98.6	98.3
2-year	88.5	92.6	96.2	96.4	97.4	96.4	95.7	96.6	96.3	96.5	97.5	97.1	96.9	96.5	97.3	98.4	97.5	97.2	97.4	97.4	97.4	97.9	97.2	97.0	97.1	
3-year	86.5	92.0	95.5	95.8	96.4	95.8	95.1	96.2	95.7	95.9	97.1	96.9	96.6	95.7	97.1	98.2	96.8	97.2	97.2	97.4	96.8	97.6	96.8	97.0		
4-year	85.6	91.5	95.0	95.7	96.3	95.8	94.9	96.1	95.6	95.9	96.8	96.7	96.5	95.4	96.8	98.0	96.6	96.7	97.2	97.2	96.7	97.4	96.4			
5-year	85.0	91.1	94.7	95.7	96.3	95.8	94.9	95.8	95.6	95.9	96.6	96.3	96.1	95.2	96.8	98.0	95.6	96.7	96.9	97.0	96.4	97.4				
6-year	84.7	90.9	94.5	95.6	96.3	95.8	94.7	95.6	95.6	95.9	96.2	96.1	96.1	95.2	96.4	97.8	95.0	96.6	96.7	96.8	96.3					
7-year	84.6	90.7	94.3	95.4	96.3	95.8	94.7	95.4	95.5	95.9	96.2	96.1	96.1	95.2	96.3	97.4	95.0	96.6	96.7	96.8						
8-year	84.5	90.5	94.1	95.0	96.3	95.8	94.7	95.4	95.4	95.9	96.2	96.0	96.1	95.2	96.2	97.3	95.0	96.6	96.7							
9-year	84.5	90.1	94.0	94.9	96.2	95.8	94.7	95.4	95.2	95.9	96.2	95.7	96.1	95.2	96.1	97.3	95.0	96.6								
10-year	83.7	90.0	93.8	94.9	96.1	95.8	94.7	94.7	95.2	95.9	96.2	95.5	96.1	95.2	95.9	97.3	95.0									
11-year	83.3	89.8	93.7	94.9	96.1	95.8	94.7	94.2	95.1	95.9	96.1	95.2	96.1	95.2	95.8	97.2										
12-year	83.0	89.8	93.5	94.8	96.1	95.8	94.7	94.1	95.1	95.9	96.0	95.2	96.1	95.2	95.3											
13-year	82.6	89.4	93.2	94.8	95.9	95.8	94.7	94.1	95.1	95.9	95.8	95.2	96.1	94.7												
14-year	82.1	88.9	93.1	94.8	95.7	95.8	94.7	94.0	95.1	95.9	95.6	95.0	96.1													
15-year	81.8	88.6	93.1	94.8	95.7	95.8	94.7	94.0	95.1	95.9	95.4	94.9														
16-year	81.3	88.2	92.9	94.6	95.7	95.8	94.7	93.8	95.1	95.9	95.4															
17-year	80.8	87.9	92.7	94.4	95.7	95.8	94.7	93.4	95.1	95.9																
18-year	80.1	87.7	92.6	94.3	95.7	95.8	94.5	93.2	95.1																	
19-year	79.6	87.4	92.5	94.2	95.7	95.8	94.4	93.2																		
20-year	79.5	87.1	92.3	94.1	95.7	95.8	94.2																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 26.9
Cancer of the Thyroid (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																									
	1975-1979	1980-1984	1985-1989	1990-1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
1-year	94.9	95.1	96.0	96.2	96.7	96.8	96.4	97.7	97.3	97.6	97.6	97.1	98.0	98.3	97.7	98.4	98.5	98.3	98.8	98.9	98.8	98.8	98.3	98.7	98.8	98.7
2-year	93.5	94.5	95.2	95.7	96.1	96.6	96.2	97.1	97.1	97.0	97.4	97.0	97.9	97.9	97.3	98.3	98.1	98.2	98.7	98.6	98.7	98.5	98.3	98.7	98.7	
3-year	92.9	93.7	94.8	95.3	96.0	96.0	96.2	96.9	97.1	96.9	97.0	97.0	97.7	97.9	97.3	98.2	98.1	98.2	98.7	98.6	98.6	98.4	98.2	98.7		
4-year	92.5	93.4	94.3	95.0	95.9	95.4	95.7	96.3	97.0	96.9	97.0	97.0	97.4	97.7	97.3	98.2	98.0	98.0	98.7	98.4	98.6	98.4	98.2			
5-year	92.0	93.3	94.1	94.6	95.8	95.0	95.3	96.2	96.8	96.3	96.2	97.0	97.3	97.4	97.3	98.2	97.9	97.8	98.7	98.1	98.6	98.4				
6-year	91.7	93.1	93.8	94.3	95.8	94.8	95.2	95.8	96.3	96.3	96.1	96.7	97.0	97.2	97.3	98.2	97.9	97.8	98.7	98.1	98.6					
7-year	91.5	92.7	93.5	94.2	95.6	94.5	95.2	95.8	96.2	96.1	96.1	96.5	96.8	96.9	97.3	98.1	97.9	97.7	98.7	98.1						
8-year	91.1	92.6	93.3	94.1	95.5	94.2	95.1	95.8	96.0	95.9	95.9	96.3	96.7	96.8	97.3	98.0	97.9	97.5	98.7							
9-year	90.7	92.3	93.1	93.9	95.5	93.8	95.0	95.8	95.7	95.9	95.4	96.3	96.7	96.8	97.3	98.0	97.9	97.5								
10-year	90.5	92.0	92.8	93.6	95.2	93.7	95.0	95.8	95.4	95.8	95.3	96.3	96.7	96.7	97.3	98.0	97.9									
11-year	90.3	91.7	92.7	93.3	95.1	93.3	95.0	95.8	95.0	95.8	95.3	96.1	96.7	96.5	97.2	98.0										
12-year	90.3	91.4	92.7	93.0	95.1	93.0	95.0	95.8	94.7	95.8	95.3	95.9	96.7	96.4	97.0											
13-year	90.0	91.3	92.5	93.0	95.1	92.8	95.0	95.1	94.6	95.8	95.3	95.9	96.7	96.4												
14-year	90.0	91.1	92.4	92.7	95.1	92.7	95.0	94.7	94.5	95.8	95.3	95.9	96.7													
15-year	89.7	90.7	92.4	92.4	94.3	92.4	95.0	94.2	94.4	95.8	95.2	95.9														
16-year	89.7	90.6	92.4	92.2	93.8	92.4	95.0	94.0	94.4	95.8	94.9															
17-year	89.6	90.6	92.3	92.2	93.8	92.4	95.0	93.8	94.4	95.7																
18-year	89.4	90.5	92.2	92.2	93.8	92.4	94.4	93.8	94.1																	
19-year	89.2	90.5	91.9	92.2	93.8	92.4	94.3	93.4																		
20-year	89.2	90.5	91.9	92.2	93.8	92.3	94.3																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 26.10
Cancer of the Thyroid (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	94.6	92.5	94.8	94.5	93.4	93.7	91.8	97.1	96.5	96.3	96.1	95.1	96.0	96.7	97.1	96.4	97.3	96.8	98.3	97.9	97.4	97.9	96.7	97.2	97.9	97.4
2-year	92.5	92.0	93.7	93.2	91.9	93.4	90.9	95.4	96.0	94.9	95.5	94.9	95.4	95.1	95.6	96.1	96.6	96.1	98.2	96.3	96.9	97.1	96.5	96.9	97.9	
3-year	91.4	90.8	93.5	92.4	91.6	92.2	90.9	94.7	96.0	94.9	94.2	94.9	94.2	94.8	95.6	96.1	96.6	96.1	98.2	96.1	96.7	96.8	96.4	96.5		
4-year	90.5	90.4	92.1	91.9	91.3	91.8	89.5	93.7	96.0	94.9	93.6	94.7	93.4	94.3	95.6	96.1	96.6	95.8	98.2	95.7	96.7	96.7	95.8			
5-year	89.8	90.2	91.6	91.0	91.2	89.9	88.2	93.7	96.0	93.7	92.4	94.3	93.2	94.0	95.6	95.7	96.6	95.1	98.0	94.4	96.7	96.4				
6-year	89.6	89.8	91.2	90.6	90.4	89.9	88.0	93.5	95.6	93.7	92.0	93.8	91.7	94.0	95.6	95.7	96.6	95.0	97.8	94.3	96.7					
7-year	89.4	89.3	90.0	90.4	88.7	89.6	88.0	93.3	95.2	93.4	92.0	92.4	91.7	92.8	95.6	95.3	96.6	94.8	97.8	94.3						
8-year	89.3	89.3	89.7	90.1	88.7	88.6	87.8	93.3	94.9	93.4	91.1	91.5	91.7	92.5	95.4	95.3	96.6	94.6	97.8							
9-year	88.8	88.6	88.4	89.9	88.7	86.8	86.9	93.3	92.8	93.4	89.8	91.0	91.7	92.1	95.2	95.3	96.6	94.5								
10-year	88.3	87.9	88.4	89.7	87.4	86.7	86.9	93.2	92.4	93.4	89.6	91.0	91.7	91.5	94.7	95.3	96.6									
11-year	87.4	87.2	88.4	89.5	86.2	85.0	86.5	93.2	92.4	93.4	89.6	90.1	91.7	91.5	94.7	95.3										
12-year	87.4	86.4	88.4	88.7	86.0	84.3	86.5	93.1	92.4	93.4	89.3	89.9	91.7	91.5	94.2											
13-year	87.3	86.2	87.8	88.2	86.0	83.3	86.5	93.1	92.4	92.9	89.3	89.9	91.7	91.5												
14-year	87.3	86.2	87.7	87.8	85.8	83.0	86.5	93.1	92.4	92.9	89.3	89.5	91.7													
15-year	86.7	85.5	87.7	87.2	85.8	83.0	86.5	93.1	92.4	92.1	87.8	89.4														
16-year	86.6	85.4	87.7	87.0	85.8	83.0	86.5	92.5	92.4	91.8	87.5															
17-year	86.6	85.0	87.7	87.0	85.8	83.0	86.5	91.9	92.4	91.6																
18-year	86.6	84.6	87.7	86.6	85.8	83.0	86.1	91.9	92.4																	
19-year	86.6	84.6	87.7	86.5	85.8	82.6	85.1	91.9																		
20-year	86.6	84.6	87.7	86.5	85.8	81.3	85.1																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 26.11
Cancer of the Thyroid (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	95.0	96.0	96.3	96.8	97.8	97.7	97.7	97.9	97.6	98.0	98.0	97.7	98.6	98.8	97.9	99.0	98.9	98.7	98.9	99.2	99.1	99.1	98.8	99.1	99.0	99.1
2-year	93.8	95.3	95.7	96.6	97.3	97.5	97.7	97.6	97.4	97.7	98.0	97.7	98.5	98.7	97.9	99.0	98.5	98.7	98.8	99.1	99.1	98.8	98.7	99.1	98.9	
3-year	93.3	94.7	95.2	96.3	97.3	97.1	97.7	97.5	97.4	97.5	97.7	97.7	98.5	98.7	97.9	98.9	98.5	98.7	98.8	99.1	99.1	98.8	98.7	99.1		
4-year	93.1	94.4	95.0	96.1	97.3	96.6	97.5	97.1	97.2	97.5	97.7	97.7	98.5	98.7	97.9	98.9	98.3	98.6	98.7	99.1	99.1	98.8	98.7			
5-year	92.7	94.3	94.9	95.9	97.2	96.4	97.4	96.9	96.9	97.0	97.3	97.7	98.4	98.3	97.9	98.9	98.3	98.5	98.7	99.0	99.1	98.8				
6-year	92.3	94.2	94.7	95.6	97.2	96.2	97.3	96.5	96.5	97.0	97.3	97.6	98.3	98.1	97.9	98.9	98.3	98.5	98.7	99.0	99.1					
7-year	92.2	93.8	94.6	95.4	97.2	95.9	97.3	96.5	96.3	97.0	97.3	97.6	98.1	98.0	97.9	98.9	98.3	98.5	98.7	99.0						
8-year	91.7	93.7	94.5	95.4	97.2	95.9	97.3	96.5	96.2	96.6	97.3	97.6	98.0	98.0	97.9	98.7	98.3	98.3	98.7							
9-year	91.4	93.6	94.5	95.3	97.2	95.8	97.3	96.5	96.2	96.6	97.0	97.6	98.0	98.0	97.9	98.7	98.3	98.3								
10-year	91.2	93.4	94.2	94.9	97.2	95.7	97.3	96.5	96.0	96.4	97.0	97.6	97.8	98.0	97.9	98.7	98.3									
11-year	91.2	93.2	94.1	94.6	97.2	95.7	97.3	96.5	95.8	96.4	97.0	97.6	97.8	97.9	97.9	98.7										
12-year	91.1	93.1	94.0	94.5	97.2	95.5	97.3	96.5	95.3	96.4	97.0	97.6	97.8	97.8	97.7											
13-year	91.0	92.9	94.0	94.5	97.2	95.5	97.3	95.6	94.8	96.4	97.0	97.6	97.8	97.7												
14-year	90.8	92.8	94.0	94.3	97.2	95.5	97.3	95.0	94.6	96.4	97.0	97.6	97.8													
15-year	90.7	92.5	93.9	94.1	96.7	95.1	97.3	94.4	94.4	96.4	97.0	97.6														
16-year	90.7	92.3	93.8	94.0	95.9	95.1	97.3	94.3	94.4	96.4	96.9															
17-year	90.4	92.3	93.6	94.0	95.9	95.0	97.2	94.2	94.3	96.4																
18-year	90.0	92.3	93.5	93.9	95.9	95.0	96.9	94.1	94.1																	
19-year	89.8	92.3	93.3	93.9	95.9	95.0	96.9	93.8																		
20-year	89.8	92.3	93.3	93.9	95.9	95.0	96.9																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 27.9
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	86.9	88.6	89.9	90.0	90.7	89.5	91.2	90.9	90.0	90.9	90.9	89.7	90.4	90.7	90.6	91.1	90.2	89.2	89.5	89.1	90.5	89.0	89.2	89.9	89.9	88.3
2-year	80.8	83.5	85.1	85.4	86.6	85.4	86.2	86.6	85.0	86.5	86.4	85.7	85.7	86.0	85.5	86.6	85.1	84.8	83.8	85.1	86.1	84.3	84.9	85.5	85.8	
3-year	77.7	80.4	82.2	82.8	84.0	83.2	83.1	83.4	82.4	83.5	84.6	83.6	83.5	83.5	82.7	83.9	82.1	81.9	81.6	81.7	83.0	81.4	82.2	82.8		
4-year	74.9	78.0	80.0	81.2	81.8	81.1	80.7	81.5	80.4	81.6	82.4	81.2	81.5	81.6	81.2	82.4	80.5	79.5	79.5	79.7	80.2	79.6	80.1			
5-year	73.0	76.0	78.1	79.5	81.1	79.5	78.5	79.2	78.1	80.3	81.4	79.9	79.8	80.1	80.1	80.8	79.1	77.5	78.2	77.3	79.2	77.6				
6-year	71.1	74.5	76.4	78.0	79.3	78.3	76.8	77.6	76.7	79.3	80.1	78.5	78.3	78.2	78.6	79.8	77.1	75.4	75.6	75.8	77.2					
7-year	69.0	73.0	74.9	76.3	77.4	77.3	75.8	76.5	75.6	77.9	78.3	77.4	77.2	76.0	77.8	77.9	75.2	73.3	74.8	74.9						
8-year	67.5	71.8	73.4	75.0	76.2	75.9	74.8	74.8	74.3	76.4	77.2	76.2	75.9	74.4	75.8	76.3	74.0	72.5	73.3							
9-year	65.8	70.1	72.2	73.5	74.6	74.1	73.2	73.5	73.2	75.5	76.0	74.6	75.0	73.2	74.4	75.6	73.2	72.1								
10-year	63.8	68.9	70.8	72.1	73.7	72.7	72.1	71.8	72.6	73.6	74.5	73.7	74.0	72.1	73.6	74.5	72.2									
11-year	62.1	67.5	69.7	70.9	73.0	71.6	71.3	71.0	71.5	71.6	73.9	72.4	72.7	70.9	73.0	73.7										
12-year	60.7	66.2	68.2	69.9	71.6	70.0	70.2	68.9	69.7	70.6	72.7	71.2	72.0	69.9	71.5											
13-year	59.7	65.2	66.8	69.0	69.3	68.7	69.9	68.0	68.3	69.2	70.2	70.8	71.3	69.3												
14-year	58.5	63.6	65.4	68.0	67.6	67.2	68.7	67.2	67.0	67.8	68.6	70.1	70.1													
15-year	57.2	62.2	64.1	66.3	67.1	67.1	67.3	66.8	65.7	66.9	67.7	68.4														
16-year	56.2	60.7	62.8	65.1	65.7	66.8	65.6	66.0	65.4	66.1	66.3															
17-year	55.1	59.7	61.8	63.8	64.2	65.4	64.6	64.5	64.0	65.5																
18-year	54.3	58.6	60.8	62.7	63.4	64.0	63.3	64.2	62.7																	
19-year	53.7	57.8	60.2	61.6	62.1	63.0	62.9	63.3																		
20-year	52.8	57.3	59.6	60.6	60.0	61.3	61.4																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 27.10
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	88.9	89.9	91.6	91.7	92.1	91.0	92.9	92.6	91.7	92.0	91.5	91.8	91.9	91.8	92.1	92.4	91.7	90.8	90.6	90.8	91.9	90.8	90.9	91.3	91.1	90.0
2-year	82.8	85.1	87.0	87.5	88.1	87.1	88.2	88.3	86.8	87.8	86.9	88.4	87.3	87.6	87.2	88.1	86.6	86.7	84.7	87.0	87.6	86.3	86.5	87.1	87.1	
3-year	79.5	82.0	84.2	85.2	85.6	84.8	85.2	84.8	84.2	84.6	85.3	86.2	85.0	85.2	84.4	85.4	83.4	83.9	82.4	83.7	84.2	83.1	83.5	84.2		
4-year	76.6	79.5	81.9	83.4	83.1	82.6	82.4	82.6	81.9	82.5	83.1	83.5	82.9	83.3	82.8	84.0	81.5	81.1	80.4	81.6	81.0	80.8	81.0			
5-year	74.3	77.4	80.0	81.7	82.8	80.9	79.9	80.3	79.3	81.0	82.3	82.0	81.0	82.0	82.0	82.6	80.1	79.0	78.8	78.8	79.7	78.8				
6-year	72.3	75.8	78.3	80.1	80.9	79.6	77.9	78.9	77.7	80.1	80.8	80.7	79.2	80.2	80.8	81.6	78.0	76.8	76.2	77.1	77.8					
7-year	70.0	74.4	76.5	78.3	79.2	78.4	76.7	77.9	76.6	78.4	78.6	79.8	77.7	77.7	79.9	79.2	75.6	74.6	75.2	76.0						
8-year	68.3	73.3	75.0	76.8	77.8	76.9	75.9	76.0	75.2	76.6	77.7	78.5	76.3	75.8	77.9	77.5	74.2	73.7	73.8							
9-year	66.4	71.4	73.7	75.3	75.6	75.5	74.3	74.9	74.5	75.2	77.1	76.6	75.2	74.7	76.2	76.4	73.3	73.2								
10-year	64.2	70.3	72.3	74.0	74.8	73.9	73.2	73.1	74.2	73.0	75.6	75.7	74.0	73.4	75.2	75.9	72.0									
11-year	62.2	69.0	71.2	72.7	73.9	72.5	72.4	72.2	73.4	70.9	74.9	74.0	72.3	71.9	74.7	75.4										
12-year	60.9	67.5	69.7	71.7	72.5	70.7	70.9	69.9	72.2	69.5	73.9	72.6	71.5	70.8	73.1											
13-year	59.6	66.5	68.3	70.8	70.7	69.4	70.0	68.9	70.8	67.7	71.4	72.1	71.0	70.3												
14-year	58.4	64.7	66.8	70.1	68.8	67.6	68.3	68.4	69.6	66.3	70.1	71.7	69.6													
15-year	57.2	63.1	65.3	68.5	68.0	67.4	66.5	67.9	68.3	65.3	68.8	70.1														
16-year	55.8	61.4	63.9	67.2	66.2	67.1	64.5	66.6	67.9	64.1	67.1															
17-year	54.7	60.4	63.1	65.9	64.9	66.0	63.5	65.4	66.8	63.6																
18-year	53.7	59.4	61.9	64.9	64.2	64.3	62.4	65.0	64.8																	
19-year	52.9	58.8	61.3	63.6	62.6	64.2	61.9	63.6																		
20-year	52.1	58.3	60.8	62.7	59.8	62.3	59.8																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 27.11
Cancer of the Urinary Bladder (Invasive and *In Situ*)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Survival Time																										
1-year	81.4	84.9	84.9	85.0	86.7	85.1	86.3	85.9	85.0	87.4	89.2	83.7	86.0	87.2	86.1	87.1	85.1	84.1	85.9	84.0	86.2	83.3	83.8	85.6	85.5	82.6
2-year	75.3	78.9	79.7	79.2	82.7	80.6	80.5	81.3	80.0	82.4	84.8	77.9	81.2	81.2	80.7	81.9	80.0	79.1	81.1	79.1	81.1	78.2	79.8	80.8	81.3	
3-year	72.6	75.9	76.5	76.1	79.7	78.7	77.2	79.5	77.2	80.4	82.2	76.1	79.4	78.7	77.6	79.2	77.9	75.8	79.2	75.6	79.1	76.2	78.2	78.7		
4-year	70.3	73.7	74.4	74.7	78.1	76.7	75.6	77.8	75.7	78.9	80.3	74.6	77.6	76.3	76.3	77.2	77.0	74.3	76.6	73.8	77.6	75.5	76.7			
5-year	69.2	72.2	72.4	73.0	76.1	75.6	74.5	76.1	74.5	78.3	78.8	73.6	76.2	74.1	74.3	75.3	75.8	73.1	76.2	72.5	77.2	73.8				
6-year	67.7	70.5	70.8	71.6	74.6	74.5	73.6	73.9	73.8	76.5	77.8	72.0	75.4	72.1	72.0	74.3	74.1	71.2	73.9	71.7	75.3					
7-year	66.1	68.8	70.0	70.7	72.5	73.9	72.9	72.4	72.8	76.3	77.4	70.1	75.2	71.0	71.3	74.0	73.5	69.7	73.3	71.5						
8-year	65.3	67.7	68.7	69.4	71.8	72.9	71.2	71.4	71.8	75.4	75.3	69.7	74.5	70.0	69.5	72.8	72.8	68.6	71.9							
9-year	64.2	66.1	67.6	68.2	71.2	69.9	70.0	69.5	69.2	75.4	72.7	68.9	74.2	68.6	69.1	72.8	72.5	67.9								
10-year	62.6	64.8	66.2	66.8	70.5	69.2	68.7	68.1	68.1	74.6	71.4	68.2	73.5	68.3	68.6	70.2	71.9									
11-year	61.5	63.3	65.0	65.6	69.8	68.7	68.0	67.5	65.7	73.8	71.0	67.4	73.5	67.9	68.0	68.5										
12-year	60.1	62.6	64.0	64.6	69.0	67.7	68.0	65.8	62.9	73.4	69.3	66.8	73.3	66.8	66.9											
13-year	59.6	61.4	62.6	63.6	65.5	66.5	68.0	65.0	61.3	73.3	66.6	66.8	71.6	65.6												
14-year	58.2	60.4	61.3	61.9	64.3	65.8	68.0	63.6	59.9	72.0	64.1	65.0	70.7													
15-year	56.8	59.5	60.4	60.2	63.9	65.8	68.0	63.5	58.5	71.6	64.1	63.3														
16-year	56.8	58.5	59.3	59.0	63.5	65.1	68.0	63.5	58.2	71.5	63.8															
17-year	55.8	57.5	57.9	57.8	62.1	63.5	67.5	61.8	56.7	71.0																
18-year	55.4	56.2	57.5	56.5	61.2	62.5	65.7	61.1	56.7																	
19-year	55.3	54.9	56.7	55.7	60.7	59.4	65.0	61.1																		
20-year	54.3	54.1	55.9	54.4	60.3	58.4	65.0																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 28.10
All Cancer Sites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Ages 0-14

Year of Diagnosis

Survival Time	1975-	1980-	1985-	1990-	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
	1979	1984	1989	1994																						
1-year	80.6	84.5	86.6	89.5	90.4	92.6	90.7	90.7	91.1	92.5	92.0	91.3	93.4	92.6	92.4	93.9	92.3	92.6	92.2	93.3	95.2	93.8	93.7	92.3	93.3	95.1
2-year	70.9	75.6	78.4	82.9	84.6	86.4	84.7	84.4	85.4	86.1	86.9	87.5	89.3	88.3	87.8	89.3	87.8	88.1	86.6	89.9	91.1	89.3	89.3	88.5	87.9	
3-year	65.4	70.7	74.4	79.7	81.0	83.6	82.3	81.6	83.1	83.3	84.1	84.2	86.3	85.5	84.9	87.9	86.2	86.1	84.0	87.4	89.7	87.7	88.0	86.3		
4-year	62.1	67.6	72.0	77.6	78.9	80.8	80.5	78.9	81.5	81.8	82.6	82.4	85.3	84.0	83.4	86.5	84.9	84.3	83.2	86.4	88.3	86.8	86.7			
5-year	59.9	65.9	70.7	76.4	77.4	80.4	79.3	78.1	79.7	80.8	81.9	81.4	84.3	83.1	82.1	85.8	83.8	83.5	82.7	85.7	87.7	86.1				
6-year	58.3	64.8	69.7	75.8	76.4	79.7	78.4	76.7	79.2	79.7	81.2	80.6	83.9	82.8	81.1	85.2	82.4	83.2	82.0	85.5	87.1					
7-year	57.2	63.9	68.8	75.2	75.9	79.1	78.0	76.3	78.5	79.1	80.3	80.2	83.3	82.2	80.8	84.8	81.9	83.1	81.6	85.1						
8-year	56.6	63.2	68.3	74.8	75.0	78.7	76.9	75.8	78.2	78.4	80.2	79.6	82.7	81.5	80.1	83.9	81.1	82.6	80.9							
9-year	56.2	62.7	67.9	74.3	74.6	78.3	76.7	75.1	77.7	78.4	79.8	79.4	82.2	81.1	79.8	83.4	80.8	82.3								
10-year	55.9	62.1	67.5	74.2	74.6	78.0	76.2	74.8	77.3	78.1	79.6	79.0	82.0	80.9	79.6	83.3	80.3									
11-year	55.6	61.9	67.2	73.8	74.3	77.9	76.1	74.4	77.0	77.9	79.4	78.3	81.5	80.8	79.6	83.2										
12-year	55.3	61.4	66.9	73.6	74.1	77.8	75.7	74.1	76.9	77.8	79.3	77.7	81.1	80.6	79.2											
13-year	55.0	61.0	66.7	73.4	74.0	77.7	75.5	73.7	76.5	77.4	79.2	77.4	81.0	80.5												
14-year	54.7	60.7	66.3	73.1	74.0	77.5	75.4	73.6	76.5	77.3	79.1	77.3	80.8													
15-year	54.4	60.5	66.2	72.9	73.9	77.3	75.4	73.5	76.5	77.0	79.1	76.8														
16-year	54.1	60.2	65.9	72.6	73.2	77.3	75.0	73.2	76.4	76.8	78.9															
17-year	53.7	60.0	65.7	72.4	72.9	77.1	74.8	73.2	76.1	76.5																
18-year	53.5	59.7	65.5	72.1	72.8	76.8	74.7	73.2	76.0																	
19-year	53.2	59.5	65.4	72.0	72.6	76.6	74.4	73.2																		
20-year	53.1	59.3	65.1	71.9	72.5	76.6	74.2																			

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

Table 28.11
All Cancer Sites (Invasive)

SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males and Females

Ages 0-19

Year of Diagnosis

Survival Time	1975-	1980-	1985-	1990-	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
	1979	1984	1989	1994																							
1-year	82.4	85.7	88.2	90.1	90.1	93.0	91.2	91.7	91.0	91.3	92.3	92.1	93.5	92.2	92.9	94.0	93.2	93.9	93.6	93.2	95.4	94.8	94.2	93.5	93.8	95.0	
2-year	73.0	77.4	80.4	83.3	84.6	87.5	85.0	85.4	85.9	84.4	87.3	87.4	89.2	88.0	88.6	89.1	88.4	89.3	88.5	89.7	91.5	90.1	90.4	90.0	88.8		
3-year	68.0	72.5	76.5	80.0	81.3	85.0	82.0	82.5	83.3	82.0	84.4	84.3	86.3	84.8	86.1	87.4	86.6	87.2	86.1	87.2	89.3	87.6	88.9	87.8			
4-year	64.9	69.6	74.0	78.1	79.8	82.3	80.4	80.0	81.5	80.5	82.6	82.3	85.3	83.6	84.6	85.9	85.0	85.2	85.1	86.3	88.1	86.4	87.4				
5-year	62.9	67.9	72.9	76.8	78.4	81.8	78.8	78.9	80.0	79.3	81.9	81.2	84.3	82.4	83.4	85.1	83.7	84.0	84.5	85.5	87.2	85.7					
6-year	61.3	66.5	72.0	76.1	77.4	81.2	77.7	77.8	79.6	78.5	81.3	80.3	83.9	82.1	82.2	84.1	82.5	83.4	83.6	85.3	86.6						
7-year	60.3	65.8	71.2	75.4	76.9	80.7	77.5	77.4	79.1	77.8	80.4	79.9	83.2	81.5	81.8	83.6	82.0	82.9	83.3	84.8							
8-year	59.6	65.0	70.5	75.0	76.0	80.2	76.7	77.1	78.9	77.3	80.2	79.1	82.6	81.0	81.1	82.8	81.2	82.5	82.8								
9-year	59.1	64.4	70.1	74.5	75.7	79.8	76.5	76.5	78.5	77.1	79.8	78.8	82.3	80.5	80.7	82.5	80.8	82.2									
10-year	58.7	63.9	69.7	74.3	75.5	79.5	76.0	76.1	78.2	76.8	79.7	78.5	82.1	80.2	80.5	82.3	80.3										
11-year	58.4	63.6	69.4	74.0	75.4	79.3	75.8	75.8	77.9	76.7	79.5	77.9	81.8	80.1	80.5	82.2											
12-year	58.0	63.2	69.2	73.8	75.2	79.2	75.6	75.3	77.6	76.4	79.3	77.4	81.4	79.9	80.1												
13-year	57.8	62.8	68.9	73.5	75.1	79.1	75.3	74.9	77.4	76.1	79.1	77.2	81.2	79.8													
14-year	57.6	62.5	68.4	73.2	75.0	79.0	75.3	74.9	77.4	75.8	79.0	77.1	81.0														
15-year	57.3	62.2	68.2	73.1	74.9	78.8	75.1	74.8	77.3	75.6	79.0	76.9															
16-year	57.0	61.8	67.9	72.8	74.3	78.7	74.7	74.5	77.2	75.4	78.9																
17-year	56.7	61.6	67.7	72.5	74.1	78.4	74.6	74.5	77.0	75.1																	
18-year	56.4	61.4	67.4	72.3	74.0	78.0	74.5	74.4	76.8																		
19-year	56.2	61.1	67.2	72.1	74.0	77.9	74.2	74.4																			
20-year	55.9	60.8	66.9	72.0	73.7	77.7	73.9																				

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). [Expected survival rates](#) are derived from the U.S. Annual Life Tables.