

Table 1.15

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Males, 21 SEER Areas, 2015-2017

Site	All Races	Whites	Blacks
	Percent ( 95% C.I. )	Percent ( 95% C.I. )	Percent ( 95% C.I. )
All Sites	40.46 ( 40.36, 40.57 )	40.15 ( 40.04, 40.27 )	38.04 ( 37.74, 38.36 )
Invasive and In Situ	42.14 ( 42.04, 42.25 )	41.87 ( 41.75, 41.99 )	38.45 ( 38.14, 38.77 )
Oral Cavity and Pharynx	1.66 ( 1.64, 1.68 )	1.74 ( 1.72, 1.77 )	1.11 ( 1.06, 1.16 )
Esophagus	0.79 ( 0.78, 0.81 )	0.85 ( 0.83, 0.86 )	0.55 ( 0.51, 0.59 )
Stomach	1.04 ( 1.02, 1.06 )	0.93 ( 0.91, 0.95 )	1.27 ( 1.21, 1.34 )
Colon and Rectum	4.30 ( 4.27, 4.33 )	4.21 ( 4.18, 4.25 )	4.33 ( 4.23, 4.44 )
Invasive and In Situ	4.46 ( 4.43, 4.50 )	4.36 ( 4.33, 4.40 )	4.51 ( 4.40, 4.62 )
Liver and Intrahepatic Bile Duct	1.46 ( 1.44, 1.47 )	1.31 ( 1.29, 1.33 )	1.64 ( 1.59, 1.70 )
Pancreas	1.68 ( 1.66, 1.70 )	1.69 ( 1.66, 1.71 )	1.59 ( 1.53, 1.66 )
Larynx	0.52 ( 0.50, 0.53 )	0.52 ( 0.50, 0.53 )	0.64 ( 0.60, 0.68 )
Invasive and In Situ	0.55 ( 0.54, 0.57 )	0.56 ( 0.54, 0.57 )	0.67 ( 0.63, 0.71 )
Lung and Bronchus	6.57 ( 6.52, 6.61 )	6.58 ( 6.54, 6.63 )	6.34 ( 6.21, 6.47 )
Melanoma of the Skin	2.79 ( 2.76, 2.81 )	3.23 ( 3.20, 3.26 )	0.10 ( 0.08, 0.12 )
Invasive and In Situ	4.84 ( 4.80, 4.88 )	5.49 ( 5.45, 5.54 )	0.13 ( 0.11, 0.15 )
Breast	0.13 ( 0.13, 0.14 )	0.13 ( 0.13, 0.14 )	0.17 ( 0.15, 0.19 )
Invasive and In Situ	0.15 ( 0.14, 0.15 )	0.14 ( 0.14, 0.15 )	0.18 ( 0.16, 0.21 )
Prostate	12.12 ( 12.06, 12.17 )	11.28 ( 11.22, 11.34 )	16.29 ( 16.10, 16.49 )
Testis	0.41 ( 0.40, 0.41 )	0.48 ( 0.47, 0.49 )	0.11 ( 0.10, 0.13 )
Urinary Bladder (Invasive and In Situ)	3.85 ( 3.81, 3.88 )	4.16 ( 4.12, 4.20 )	1.84 ( 1.76, 1.92 )
Kidney and Renal Pelvis	2.18 ( 2.15, 2.20 )	2.24 ( 2.22, 2.27 )	1.98 ( 1.92, 2.05 )
Brain and Other Nervous System	0.69 ( 0.68, 0.70 )	0.76 ( 0.74, 0.77 )	0.38 ( 0.35, 0.41 )
Thyroid	0.69 ( 0.67, 0.70 )	0.73 ( 0.72, 0.74 )	0.32 ( 0.29, 0.35 )
Hodgkin Lymphoma	0.23 ( 0.22, 0.24 )	0.24 ( 0.24, 0.25 )	0.21 ( 0.19, 0.23 )
Non-Hodgkin Lymphoma	2.40 ( 2.38, 2.43 )	2.51 ( 2.48, 2.54 )	1.47 ( 1.42, 1.54 )
Myeloma	0.95 ( 0.94, 0.97 )	0.89 ( 0.87, 0.91 )	1.50 ( 1.44, 1.57 )
Leukemia	1.83 ( 1.81, 1.86 )	1.93 ( 1.91, 1.96 )	1.18 ( 1.12, 1.24 )
Acute Lymphocytic Leukemia	0.15 ( 0.14, 0.16 )	0.16 ( 0.16, 0.17 )	0.08 ( 0.07, 0.10 )
Chronic Lymphocytic Leukemia	0.74 ( 0.73, 0.75 )	0.79 ( 0.77, 0.81 )	0.42 ( 0.39, 0.46 )
Acute Myeloid Leukemia	0.56 ( 0.54, 0.57 )	0.58 ( 0.56, 0.59 )	0.38 ( 0.35, 0.42 )
Chronic Myeloid Leukemia	0.25 ( 0.24, 0.26 )	0.26 ( 0.25, 0.27 )	0.19 ( 0.17, 0.22 )
Kaposi Sarcoma	0.18 ( 0.17, 0.19 )	0.20 ( 0.19, 0.21 )	0.09 ( 0.07, 0.11 )
Mesothelioma	0.07 ( 0.07, 0.08 )	0.06 ( 0.05, 0.06 )	0.14 ( 0.13, 0.16 )

Devcan Version 6.7.8, April 2020, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Note: Invasive cancer only unless specified otherwise.  
A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table 1.15 - continued

Lifetime Risk (Percent) of Being Diagnosed with Cancer by Site and Race/Ethnicity

Males, 21 SEER Areas, 2015-2017

Site	Asian/Pacific Islanders	American Indian/ Alaska Natives <sup>a</sup>	Hispanics <sup>b</sup>
	Percent ( 95% C.I. )	Percent ( 95% C.I. )	Percent ( 95% C.I. )
All Sites	34.92 ( 34.50, 35.34 )	26.69 ( 25.37, 28.21 )	37.16 ( 36.79, 37.54 )
Invasive and In Situ	35.23 ( 34.81, 35.66 )	26.96 ( 25.64, 28.49 )	37.71 ( 37.34, 38.09 )
Oral Cavity and Pharynx	1.29 ( 1.22, 1.36 )	1.13 ( 0.92, 1.64 )	1.10 ( 1.04, 1.17 )
Esophagus	0.53 ( 0.48, 0.59 )	0.59 ( 0.44, 1.08 )	0.64 ( 0.59, 0.71 )
Stomach	1.89 ( 1.79, 1.99 )	0.96 ( 0.74, 1.48 )	1.57 ( 1.50, 1.66 )
Colon and Rectum	4.61 ( 4.46, 4.76 )	3.78 ( 3.28, 4.53 )	4.53 ( 4.41, 4.67 )
Invasive and In Situ	4.78 ( 4.63, 4.94 )	3.90 ( 3.39, 4.66 )	4.72 ( 4.59, 4.85 )
Liver and Intrahepatic Bile Duct	2.54 ( 2.44, 2.65 )	2.28 ( 1.96, 2.86 )	2.36 ( 2.28, 2.45 )
Pancreas	1.64 ( 1.55, 1.75 )	1.70 ( 1.30, 2.38 )	1.65 ( 1.57, 1.74 )
Larynx	0.28 ( 0.24, 0.33 )	0.40 ( 0.27, 0.89 )	0.48 ( 0.44, 0.53 )
Invasive and In Situ	0.29 ( 0.26, 0.34 )	0.43 ( 0.29, 0.93 )	0.52 ( 0.48, 0.58 )
Lung and Bronchus	6.71 ( 6.51, 6.91 )	4.33 ( 3.73, 5.18 )	4.49 ( 4.35, 4.63 )
Melanoma of the Skin	0.20 ( 0.16, 0.24 )	0.53 ( 0.37, 1.03 )	0.55 ( 0.51, 0.60 )
Invasive and In Situ	0.28 ( 0.24, 0.33 )	0.86 ( 0.66, 1.38 )	0.87 ( 0.82, 0.94 )
Breast	0.09 ( 0.07, 0.12 )	0.03 ( 0.00, 0.54 )	0.09 ( 0.07, 0.11 )
Invasive and In Situ	0.10 ( 0.08, 0.13 )	0.04 ( 0.01, 0.55 )	0.09 ( 0.07, 0.12 )
Prostate	7.94 ( 7.75, 8.13 )	5.62 ( 5.03, 6.44 )	11.03 ( 10.84, 11.22 )
Testis	0.16 ( 0.15, 0.19 )	0.30 ( 0.23, 0.76 )	0.39 ( 0.37, 0.41 )
Urinary Bladder (Invasive and In Situ)	2.42 ( 2.30, 2.56 )	1.84 ( 1.51, 2.45 )	2.56 ( 2.45, 2.68 )
Kidney and Renal Pelvis	1.50 ( 1.43, 1.59 )	2.09 ( 1.80, 2.66 )	2.34 ( 2.26, 2.43 )
Brain and Other Nervous System	0.46 ( 0.42, 0.51 )	0.41 ( 0.28, 0.90 )	0.59 ( 0.55, 0.64 )
Thyroid	0.74 ( 0.70, 0.79 )	0.31 ( 0.21, 0.78 )	0.57 ( 0.53, 0.61 )
Hodgkin Lymphoma	0.13 ( 0.11, 0.16 )	0.09 ( 0.04, 0.58 )	0.22 ( 0.20, 0.25 )
Non-Hodgkin Lymphoma	2.15 ( 2.04, 2.26 )	1.05 ( 0.83, 1.58 )	2.39 ( 2.30, 2.49 )
Myeloma	0.68 ( 0.62, 0.74 )	0.74 ( 0.55, 1.26 )	0.97 ( 0.92, 1.04 )
Leukemia	1.21 ( 1.13, 1.29 )	1.04 ( 0.81, 1.57 )	1.40 ( 1.34, 1.48 )
Acute Lymphocytic Leukemia	0.13 ( 0.11, 0.17 )	0.14 ( 0.09, 0.61 )	0.22 ( 0.21, 0.25 )
Chronic Lymphocytic Leukemia	0.23 ( 0.20, 0.27 )	0.31 ( 0.20, 0.79 )	0.35 ( 0.32, 0.40 )
Acute Myeloid Leukemia	0.55 ( 0.50, 0.61 )	0.26 ( 0.17, 0.74 )	0.48 ( 0.45, 0.53 )
Chronic Myeloid Leukemia	0.20 ( 0.17, 0.24 )	0.18 ( 0.09, 0.67 )	0.20 ( 0.17, 0.24 )
Kaposi Sarcoma	0.09 ( 0.07, 0.12 )	0.10 ( 0.03, 0.60 )	0.16 ( 0.13, 0.19 )
Mesothelioma	0.04 ( 0.02, 0.06 )	0.06 ( 0.02, 0.56 )	0.12 ( 0.10, 0.15 )

Devcan Version 6.7.8, April 2020, National Cancer Institute (<https://surveillance.cancer.gov/devcan/>).

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

Note: Invasive cancer only unless specified otherwise.

<sup>a</sup> Underlying incidence data for American Indian/Alaska Native are based on the PRCD(A/Purchased/Referred Care Delivery Areas) counties.<sup>b</sup> Hispanic is not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. A percent or confidence interval value of 0.00 represents a value that is below 0.005.