

Table 2.4
All Cancer Sites (Invasive)

Delay-adjusted SEER Incidence^a Rates by Year, Race and Sex

Year of Diagnosis:	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
1975-1979	407.23	484.12	365.02	409.26	485.32	369.85	437.62	564.41	350.93
1980	418.09	505.83	368.15	419.52	505.23	373.41	467.51	615.07	366.63
1981	425.28	510.80	377.17	427.70	510.84	383.89	472.90	634.34	362.89
1982	424.49	511.10	376.18	427.78	512.52	383.41	465.28	622.36	359.24
1983	431.11	519.94	380.71	434.35	520.95	387.93	487.26	646.27	382.24
1984	440.01	525.48	392.46	443.66	527.77	399.53	488.26	643.87	386.15
1985	448.94	531.30	403.24	453.60	533.72	411.55	489.39	643.82	388.62
1986	451.36	538.15	402.34	456.09	542.94	409.26	493.79	643.41	396.80
1987	468.14	563.36	413.34	475.07	569.08	423.42	501.15	666.26	394.20
1988	464.04	558.53	409.48	470.38	565.08	418.02	504.65	658.20	405.13
1989	468.01	567.21	408.62	475.42	573.81	418.82	493.01	661.19	385.30
1990	482.13	592.90	415.66	488.99	599.49	424.82	522.41	699.83	409.93
1991	503.15	637.52	419.04	510.68	644.72	428.64	549.60	765.72	412.80
1992	510.81	658.50	414.95	517.14	663.96	423.57	570.51	811.80	414.26
1993	493.57	622.41	408.35	497.08	620.68	416.97	570.41	822.06	404.91
1994	483.72	589.99	413.71	487.79	588.05	423.34	559.02	785.28	409.71
1995	477.36	572.13	415.22	482.62	570.87	426.41	544.75	752.47	405.49
1996	479.61	573.03	418.49	485.43	574.80	428.46	543.26	730.94	418.76
1997	486.88	576.21	428.72	493.95	577.27	441.21	543.56	744.48	412.36
1998	488.82	573.44	434.38	496.99	577.03	446.80	541.58	728.72	421.12
1999	491.06	581.27	431.07	499.15	583.63	443.94	544.66	729.92	422.97
2000	487.04	581.18	422.77	497.23	585.15	438.21	532.73	719.39	407.79
2001	490.80	582.48	427.68	501.92	588.90	442.87	525.14	705.80	403.37
2002	488.64	575.91	428.89	498.47	582.40	441.42	539.22	704.83	428.65
2003	476.49	561.45	418.13	487.15	566.57	433.19	523.83	687.84	414.66
2004	477.74	559.60	421.11	487.69	565.33	434.14	528.39	691.22	421.15
2005	473.79	550.59	420.50	485.53	559.18	434.75	514.02	650.95	423.49
2006	477.61	557.91	420.33	489.57	565.91	435.26	514.15	653.78	418.31
2007	483.68	569.11	422.00	494.70	575.76	436.32	522.63	675.79	417.33
2008	477.30	550.56	425.18	487.67	556.14	439.23	521.08	670.28	419.70
2009	476.57	542.00	430.21	487.25	548.80	443.69	515.34	647.80	425.64
2010	467.37	531.87	421.05	479.45	540.30	435.92	497.77	621.58	412.37
2011	462.55	520.59	421.66	474.44	529.04	436.27	492.42	598.35	419.89
2012	450.56	492.91	422.32	461.29	500.25	436.17	489.75	582.86	425.59
2013	445.61	485.58	419.49	457.13	493.20	434.67	476.57	570.26	411.52
2014	444.73	479.27	423.12	458.87	490.12	440.57	458.81	536.14	406.95
2015	445.98	480.78	424.15	460.63	492.48	441.71	461.34	536.29	410.18
2016	445.66	482.56	421.44	458.51	491.62	437.79	470.84	546.76	418.53
2017	444.78	482.14	419.97	456.73	489.32	436.32	464.29	551.48	403.21

Reference: [Cancer Incidence Rates Adjusted for Reporting Delay](#).

Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#).

SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Delay-adjusted rate is not shown for observed rates based on less than 16 cases for the time interval.

^a

-