

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2013-2017</u>						
Age at Diagnosis/Death:						
All ages	7.4	7.2	8.7	2.3	2.2	3.4
Under 65	6.9	7.0	7.2	1.8	1.8	2.4
65 and over	10.4	9.0	18.9	5.5	5.0	10.9
All ages (WHO world std) ^c	6.4	6.3	7.2	1.7	1.6	2.5
<u>Age-Specific Rates, 2013-2017</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	0.8	0.8	0.9	0.1	0.1	-
25-29	4.9	5.2	4.2	0.6	0.6	0.6
30-34	9.7	10.6	7.7	1.4	1.4	1.6
35-39	13.2	14.1	11.4	2.3	2.4	2.8
40-44	14.3	14.5	14.4	3.1	3.2	3.5
45-49	13.3	13.1	13.8	4.0	4.0	4.8
50-54	12.1	11.5	14.2	4.4	4.2	5.8
55-59	12.1	11.2	16.2	4.7	4.4	7.4
60-64	11.7	10.6	15.8	4.9	4.5	8.4
65-69	11.0	10.0	16.9	5.1	4.6	9.5
70-74	10.5	9.2	19.6	5.2	4.7	10.5
75-79	10.9	9.3	19.0	5.6	5.0	11.0
80-84	10.1	8.2	22.0	6.0	5.3	12.6
85+	8.3	7.0	18.0	6.4	5.7	12.5

^a SEER 21 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.