



Table 5.14
Cancer of the Cervix Uteri (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2013-2017

Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	2.27	0.02		
High Five States				
Alabama	3.50	0.16	01(01,06)	54.36 ^c
Mississippi	3.49	0.21	02(01,07)	53.91 ^c
Arkansas	3.47	0.21	03(01,08)	53.08 ^c
Louisiana	3.13	0.16	04(01,12)	38.14 ^c
Oklahoma	3.00	0.18	05(01,14)	32.19 ^c
Low Five States				
Utah	1.31	0.14	47(37,51)	-42.11 ^c
North Dakota	1.31	0.26	48(24,51)	-42.39 ^c
New Hampshire	1.26	0.18	49(35,51)	-44.43 ^c
Massachusetts	1.11	0.08	50(45,51)	-50.99 ^c
Vermont	1.10	0.26	51(31,51)	-51.44 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.50	0.16	01(01,06)	54.36 ^c	Montana	1.96	0.28	31(07,49)	-13.67
Alaska	1.63	0.30	38(11,51)	-28.21 ^c	Nebraska	2.14	0.20	24(08,41)	-5.38
Arizona	2.20	0.11	23(12,34)	-3.00	Nevada	2.51	0.18	13(04,31)	10.92
Arkansas	3.47	0.21	03(01,08)	53.08 ^c	New Hampshire	1.26	0.18	49(35,51)	-44.43 ^c
California	2.28	0.05	20(14,26)	0.78	New Jersey	2.04	0.09	28(18,37)	-9.95
Colorado	1.70	0.11	37(27,46)	-24.81 ^c	New Mexico	2.20	0.21	22(07,41)	-2.93
Connecticut	1.35	0.11	46(37,51)	-40.36 ^c	New York	2.23	0.06	21(14,29)	-1.64
Delaware	2.51	0.30	15(02,39)	10.53	North Carolina	2.04	0.09	29(19,37)	-10.04
D.C.	2.78	0.41	09(01,39)	22.85 ^c	North Dakota	1.31	0.26	48(24,51)	-42.39 ^c
Florida	2.65	0.07	11(06,16)	16.79 ^c	Ohio	2.34	0.09	17(11,28)	3.04
Georgia	2.29	0.09	19(12,30)	1.12	Oklahoma	3.00	0.18	05(01,14)	32.19 ^c
Hawaii	1.75	0.22	36(15,50)	-22.67 ^c	Oregon	2.07	0.14	27(13,39)	-8.53
Idaho	1.76	0.21	35(16,49)	-22.23 ^c	Pennsylvania	2.09	0.08	26(17,35)	-7.95
Illinois	2.31	0.08	18(12,28)	1.75	Rhode Island	1.46	0.23	44(22,51)	-35.37 ^c
Indiana	2.45	0.12	16(07,27)	8.11	South Carolina	2.56	0.14	12(05,25)	13.02
Iowa	1.94	0.16	32(15,43)	-14.60	South Dakota	1.53	0.27	43(16,51)	-32.38 ^c
Kansas	2.11	0.17	25(10,41)	-6.84	Tennessee	2.81	0.12	08(04,16)	23.89 ^c
Kentucky	2.70	0.15	10(04,22)	18.97 ^c	Texas	2.89	0.07	06(04,11)	27.64 ^c
Louisiana	3.13	0.16	04(01,12)	38.14 ^c	Utah	1.31	0.14	47(37,51)	-42.11 ^c
Maine	1.61	0.20	40(22,51)	-29.05 ^c	Vermont	1.10	0.26	51(31,51)	-51.44 ^c
Maryland	1.89	0.10	34(22,42)	-16.59 ^c	Virginia	1.89	0.09	33(23,41)	-16.58 ^c
Massachusetts	1.11	0.08	50(45,51)	-50.99 ^c	Washington	1.58	0.09	42(33,48)	-30.41 ^c
Michigan	2.01	0.09	30(19,38)	-11.39	West Virginia	2.84	0.24	07(01,25)	25.30 ^c
Minnesota	1.40	0.10	45(37,50)	-38.35 ^c	Wisconsin	1.58	0.10	41(33,48)	-30.33 ^c
Mississippi	3.49	0.21	02(01,07)	53.91 ^c	Wyoming	1.61	0.34	39(09,51)	-28.95 ^c
Missouri	2.51	0.12	14(06,25)	10.70					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.