

Table 6.23
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2013-2017</u>									
California	35.12	39.90	31.04	35.21	39.74	31.29	39.47	43.60	36.09
Greater Bay Area	33.83	38.19	29.96	34.39	38.20	30.94	41.59	43.51	39.46
San Francisco-Oakland	34.41	38.82	30.54	34.50	38.41	30.96	42.92	44.71	40.89
San Jose-Monterey	32.66	36.97	28.79	34.17	37.84	30.85	32.74	36.07	29.31
Los Angeles	35.68	41.20	31.24	35.43	40.72	31.06	39.74	44.98	35.88
Greater California	35.34	39.95	31.36	35.40	39.84	31.52	38.19	42.58	34.49
Connecticut	35.27	40.22	31.11	34.54	39.29	30.49	37.24	44.58	32.21
Detroit	41.88	47.70	37.17	39.86	44.67	35.83	50.85	62.09	43.03
Georgia	41.32	48.46	35.48	39.75	46.61	33.82	46.63	55.22	40.62
Atlanta	39.94	47.26	34.25	37.25	44.21	31.40	45.25	53.92	39.50
Rural Georgia	49.46	57.86	42.21	43.38	48.78	37.99	62.90	78.61	51.62
Greater Georgia	41.92	48.94	36.06	40.66	47.52	34.69	47.38	55.56	41.44
Hawaii	41.27	47.23	35.69	39.50	41.54	37.15	35.88	43.93	-
Idaho	35.29	39.41	31.61	35.20	39.28	31.57	-	-	-
Iowa	43.72	49.31	38.72	43.67	49.28	38.64	45.66	54.62	38.69
Kentucky	48.27	56.43	41.51	47.98	56.05	41.21	50.36	59.48	44.38
Louisiana	45.11	53.14	38.66	42.43	49.96	36.14	53.01	63.18	45.75
Massachusetts	35.15	39.90	31.35	34.98	39.51	31.35	36.71	44.90	30.83
New Jersey	40.78	46.90	35.86	40.99	47.22	35.92	43.30	49.79	38.81
New Mexico	33.13	37.50	29.41	32.64	36.92	28.99	26.01	24.97	27.73
New York	38.74	44.68	34.03	38.51	44.04	34.00	39.96	48.42	34.53
Seattle-Puget Sound	36.92	41.55	32.79	36.68	40.75	33.01	42.53	48.63	36.05
Utah	29.76	32.86	27.08	29.49	32.61	26.81	40.02	49.26	-
SEER 9 Areas ^b	37.49	42.48	33.20	36.64	41.11	32.69	45.09	52.78	39.43
SEER 13 Areas ^b	36.93	41.98	32.63	36.23	40.82	32.23	43.91	51.00	38.67
SEER 18 Areas ^b	38.39	43.89	33.75	37.91	43.14	33.42	45.23	52.54	39.87
SEER 21 Areas ^b	38.22	43.73	33.63	37.78	42.99	33.35	43.61	51.30	38.16

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia

excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.

- Statistic not shown. Rate based on less than 16 cases for the time interval.