


Table 6.27  
Cancer of the Colon and Rectum (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2013-2017

Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	11.75	0.03		
High Five States				
West Virginia	15.66	0.49	01(01,05)	33.25 <sup>c</sup>
Mississippi	15.00	0.40	02(01,07)	27.60 <sup>c</sup>
Alaska	14.29	1.01	03(01,33)	21.57 <sup>c</sup>
Louisiana	14.15	0.32	04(02,11)	20.40 <sup>c</sup>
Nevada	14.08	0.42	05(01,15)	19.77 <sup>c</sup>
Low Five States				
Massachusetts	10.47	0.22	47(34,50)	-10.97
Hawaii	10.29	0.48	48(26,51)	-12.45
Utah	9.70	0.39	49(38,51)	-17.52 <sup>c</sup>
Wyoming	9.66	0.76	50(24,51)	-17.83 <sup>c</sup>
Connecticut	9.43	0.27	51(46,51)	-19.79 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	12.97	0.29	14(05,23)	10.33	Montana	10.78	0.57	41(18,51)	-8.30
Alaska	14.29	1.01	03(01,33)	21.57 <sup>c</sup>	Nebraska	12.56	0.47	18(06,35)	6.87
Arizona	10.60	0.22	44(31,49)	-9.78	Nevada	14.08	0.42	05(01,15)	19.77 <sup>c</sup>
Arkansas	13.60	0.38	08(02,18)	15.70 <sup>c</sup>	New Hampshire	11.56	0.52	27(11,48)	-1.67
California	10.95	0.10	38(30,45)	-6.82	New Jersey	11.94	0.20	24(17,32)	1.57
Colorado	10.51	0.27	46(30,50)	-10.62	New Mexico	11.25	0.42	31(17,49)	-4.32
Connecticut	9.43	0.27	51(46,51)	-19.79 <sup>c</sup>	New York	11.16	0.13	35(27,43)	-5.05
Delaware	10.73	0.58	43(17,51)	-8.74	North Carolina	11.24	0.19	32(25,44)	-4.39
D.C.	13.24	0.86	10(01,42)	12.68	North Dakota	11.04	0.70	37(10,51)	-6.07
Florida	10.92	0.12	39(31,46)	-7.09	Ohio	13.15	0.18	11(07,18)	11.85
Georgia	12.12	0.21	22(14,30)	3.14	Oklahoma	13.86	0.34	07(02,15)	17.93 <sup>c</sup>
Hawaii	10.29	0.48	48(26,51)	-12.45	Oregon	11.39	0.30	28(19,45)	-3.11
Idaho	11.06	0.49	36(17,50)	-5.90	Pennsylvania	12.59	0.17	17(10,23)	7.09
Illinois	12.67	0.18	16(10,23)	7.75	Rhode Island	10.78	0.54	42(18,51)	-8.33
Indiana	13.01	0.25	13(06,21)	10.71	South Carolina	11.84	0.28	25(15,38)	0.73
Iowa	12.52	0.34	19(08,29)	6.51	South Dakota	13.02	0.70	12(02,38)	10.77
Kansas	12.20	0.37	21(09,36)	3.83	Tennessee	12.97	0.25	15(06,21)	10.30
Kentucky	13.92	0.32	06(02,14)	18.46 <sup>c</sup>	Texas	11.23	0.13	34(26,41)	-4.45
Louisiana	14.15	0.32	04(02,11)	20.40 <sup>c</sup>	Utah	9.70	0.39	49(38,51)	-17.52 <sup>c</sup>
Maine	11.29	0.47	30(15,49)	-3.94	Vermont	13.54	0.80	09(01,37)	15.19 <sup>c</sup>
Maryland	11.71	0.25	26(17,39)	-0.41	Virginia	11.31	0.21	29(22,44)	-3.77
Massachusetts	10.47	0.22	47(34,50)	-10.97	Washington	10.54	0.22	45(32,50)	-10.37
Michigan	12.05	0.19	23(15,30)	2.54	West Virginia	15.66	0.49	01(01,05)	33.25 <sup>c</sup>
Minnesota	10.84	0.25	40(27,49)	-7.80	Wisconsin	11.23	0.24	33(23,46)	-4.43
Mississippi	15.00	0.40	02(01,07)	27.60 <sup>c</sup>	Wyoming	9.66	0.76	50(24,51)	-17.83 <sup>c</sup>
Missouri	12.21	0.25	20(12,30)	3.87					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 10% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.