



Table 8.19  
Cancer of the Esophagus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2013-2017

Males

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	7.01	0.03		
High Five States				
Maine	9.92	0.48	01(01,09)	41.62 <sup>c</sup>
West Virginia	9.35	0.41	02(01,15)	33.42 <sup>c</sup>
New Hampshire	9.34	0.50	03(01,18)	33.22 <sup>c</sup>
Ohio	8.95	0.17	04(01,11)	27.72 <sup>c</sup>
Indiana	8.94	0.23	05(01,13)	27.52 <sup>c</sup>
Low Five States				
Texas	5.71	0.10	47(45,49)	-18.55 <sup>c</sup>
California	5.50	0.08	48(46,50)	-21.52 <sup>c</sup>
D.C.	5.26	0.64	49(27,51)	-24.91 <sup>c</sup>
Utah	5.09	0.30	50(46,51)	-27.39 <sup>c</sup>
Hawaii	4.61	0.34	51(47,51)	-34.22 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	6.86	0.23	33(21,45)	-2.06	Montana	7.73	0.50	19(03,44)	10.37
Alaska	8.79	0.85	06(01,43)	25.40 <sup>c</sup>	Nebraska	8.13	0.41	13(02,33)	16.05 <sup>c</sup>
Arizona	6.47	0.18	41(30,47)	-7.65	Nevada	6.81	0.30	34(19,46)	-2.79
Arkansas	7.49	0.31	24(09,38)	6.89	New Hampshire	9.34	0.50	03(01,18)	33.22 <sup>c</sup>
California	5.50	0.08	48(46,50)	-21.52 <sup>c</sup>	New Jersey	6.43	0.17	44(31,47)	-8.29
Colorado	7.17	0.24	31(17,42)	2.27	New Mexico	6.15	0.33	46(28,50)	-12.23
Connecticut	6.73	0.26	37(23,46)	-3.98	New York	6.49	0.11	40(33,46)	-7.41
Delaware	7.31	0.53	27(05,47)	4.25	North Carolina	6.46	0.16	42(31,47)	-7.78
D.C.	5.26	0.64	49(27,51)	-24.91 <sup>c</sup>	North Dakota	7.21	0.62	29(03,48)	2.96
Florida	6.38	0.10	45(35,47)	-8.94	Ohio	8.95	0.17	04(01,11)	27.72 <sup>c</sup>
Georgia	6.44	0.17	43(30,47)	-8.09	Oklahoma	7.21	0.27	30(15,42)	2.93
Hawaii	4.61	0.34	51(47,51)	-34.22 <sup>c</sup>	Oregon	7.99	0.27	16(05,29)	13.96
Idaho	7.87	0.43	18(03,39)	12.33	Pennsylvania	8.39	0.15	09(05,18)	19.74 <sup>c</sup>
Illinois	7.51	0.15	23(15,32)	7.15	Rhode Island	7.71	0.52	20(03,45)	9.99
Indiana	8.94	0.23	05(01,13)	27.52 <sup>c</sup>	South Carolina	6.69	0.23	38(25,46)	-4.52
Iowa	8.31	0.31	12(03,26)	18.64 <sup>c</sup>	South Dakota	8.10	0.59	15(01,43)	15.63 <sup>c</sup>
Kansas	7.69	0.32	21(07,37)	9.68	Tennessee	7.59	0.21	22(12,33)	8.37
Kentucky	7.96	0.26	17(06,29)	13.61	Texas	5.71	0.10	47(45,49)	-18.55 <sup>c</sup>
Louisiana	7.32	0.25	26(14,39)	4.40	Utah	5.09	0.30	50(46,51)	-27.39 <sup>c</sup>
Maine	9.92	0.48	01(01,09)	41.62 <sup>c</sup>	Vermont	8.73	0.69	07(01,35)	24.61 <sup>c</sup>
Maryland	6.51	0.21	39(28,47)	-7.11	Virginia	6.74	0.18	36(26,46)	-3.88
Massachusetts	8.13	0.22	14(06,25)	15.95 <sup>c</sup>	Washington	7.29	0.20	28(16,37)	4.01
Michigan	8.36	0.18	11(04,20)	19.28 <sup>c</sup>	West Virginia	9.35	0.41	02(01,15)	33.42 <sup>c</sup>
Minnesota	7.44	0.23	25(13,36)	6.22	Wisconsin	8.42	0.23	08(03,21)	20.20 <sup>c</sup>
Mississippi	6.81	0.30	35(18,46)	-2.87	Wyoming	6.93	0.67	32(04,50)	-1.15
Missouri	8.38	0.23	10(03,21)	19.59 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p < .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.