


Table 9.19  
Hodgkin Lymphoma  
Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2013-2017

Males					
State	Rate	SE	Rank(CI) <sup>d</sup>	PD	
TOTAL U.S.	0.38	0.01			
High Five States					
New Mexico	0.49	0.10	01(01,37)	28.72 <sup>c</sup>	
Wisconsin	0.48	0.06	02(01,31)	27.13 <sup>c</sup>	
Arkansas	0.48	0.08	03(01,36)	26.48 <sup>c</sup>	
Kentucky	0.47	0.07	04(01,35)	24.33 <sup>c</sup>	
Tennessee	0.46	0.05	05(01,32)	21.08 <sup>c</sup>	
Low Five States					
Pennsylvania	0.31	0.03	34(15,38)	-18.22 <sup>c</sup>	
Utah	0.30	0.07	35(02,38)	-20.40 <sup>c</sup>	
Massachusetts	0.30	0.04	36(11,38)	-20.80 <sup>c</sup>	
Virginia	0.27	0.04	37(17,38)	-27.51 <sup>c</sup>	
Nevada	0.27	0.06	38(07,38)	-28.70 <sup>c</sup>	

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	0.44	0.06	09(01,36)	16.54 <sup>c</sup>	Montana	-	-	-(-,-)	-
Alaska	-	-	-(-,-)	-	Nebraska	-	-	-(-,-)	-
Arizona	0.40	0.05	16(01,36)	7.15	Nevada	0.27	0.06	38(07,38)	-28.70 <sup>c</sup>
Arkansas	0.48	0.08	03(01,36)	26.48 <sup>c</sup>	New Hampshire	-	-	-(-,-)	-
California	0.41	0.02	13(04,27)	8.38	New Jersey	0.35	0.04	27(05,38)	-6.80
Colorado	0.34	0.05	30(03,38)	-10.60	New Mexico	0.49	0.10	01(01,37)	28.72 <sup>c</sup>
Connecticut	0.38	0.06	19(01,38)	1.54	New York	0.35	0.03	28(11,37)	-8.57
Delaware	-	-	-(-,-)	-	North Carolina	0.36	0.04	26(05,37)	-5.10
D.C.	-	-	-(-,-)	-	North Dakota	-	-	-(-,-)	-
Florida	0.34	0.02	31(13,37)	-11.05	Ohio	0.42	0.04	12(01,32)	11.28
Georgia	0.37	0.04	23(03,37)	-1.17	Oklahoma	0.42	0.07	11(01,37)	12.51
Hawaii	-	-	-(-,-)	-	Oregon	0.34	0.05	29(03,38)	-10.09
Idaho	-	-	-(-,-)	-	Pennsylvania	0.31	0.03	34(15,38)	-18.22 <sup>c</sup>
Illinois	0.41	0.04	14(02,33)	8.06	Rhode Island	-	-	-(-,-)	-
Indiana	0.38	0.05	21(02,38)	0.03	South Carolina	0.40	0.06	17(01,38)	4.81
Iowa	0.36	0.07	25(01,38)	-4.03	South Dakota	-	-	-(-,-)	-
Kansas	0.37	0.07	24(01,38)	-2.00	Tennessee	0.46	0.05	05(01,32)	21.08 <sup>c</sup>
Kentucky	0.47	0.07	04(01,35)	24.33 <sup>c</sup>	Texas	0.45	0.03	06(01,22)	19.89 <sup>c</sup>
Louisiana	0.38	0.06	20(01,38)	0.56	Utah	0.30	0.07	35(02,38)	-20.40 <sup>c</sup>
Maine	0.44	0.11	08(01,38)	16.70 <sup>c</sup>	Vermont	-	-	-(-,-)	-
Maryland	0.32	0.05	33(06,38)	-16.32 <sup>c</sup>	Virginia	0.27	0.04	37(17,38)	-27.51 <sup>c</sup>
Massachusetts	0.30	0.04	36(11,38)	-20.80 <sup>c</sup>	Washington	0.33	0.04	32(06,38)	-13.93
Michigan	0.38	0.04	18(03,36)	1.91	West Virginia	0.37	0.09	22(01,38)	-1.02
Minnesota	0.41	0.06	15(01,37)	7.98	Wisconsin	0.48	0.06	02(01,31)	27.13 <sup>c</sup>
Mississippi	0.45	0.08	07(01,37)	19.86 <sup>c</sup>	Wyoming	-	-	-(-,-)	-
Missouri	0.43	0.05	10(01,35)	14.18					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 15% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.