

Table 13.24
LeukemiaAge-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

| | All Races | | | Whites | | | Blacks | | |
|--|-----------|-------|---------|--------|-------|---------|--------|-------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| <u>SEER Incidence Rates^a, 2013-2017</u> | | | | | | | | | |
| California | 12.36 | 15.72 | 9.56 | 13.38 | 17.01 | 10.30 | 10.10 | 12.98 | 7.94 |
| Greater Bay Area | 12.03 | 15.85 | 8.87 | 13.71 | 18.03 | 10.02 | 10.84 | 14.01 | 8.38 |
| San Francisco-Oakland | 11.79 | 15.66 | 8.65 | 13.42 | 17.81 | 9.73 | 10.81 | 13.87 | 8.43 |
| San Jose-Monterey | 12.43 | 16.15 | 9.22 | 14.08 | 18.31 | 10.38 | 11.36 | 14.78 | - |
| Los Angeles | 11.62 | 14.44 | 9.34 | 12.82 | 16.07 | 10.14 | 10.11 | 12.51 | 8.48 |
| Greater California | 12.80 | 16.23 | 9.91 | 13.48 | 17.04 | 10.43 | 9.81 | 12.97 | 7.22 |
| Connecticut | 14.18 | 17.46 | 11.56 | 14.41 | 17.58 | 11.84 | 9.87 | 11.89 | 8.50 |
| Detroit | 15.82 | 20.71 | 12.09 | 17.08 | 22.38 | 12.88 | 11.75 | 14.70 | 9.76 |
| Georgia | 14.44 | 18.49 | 11.33 | 15.58 | 19.83 | 12.11 | 11.61 | 15.12 | 9.45 |
| Atlanta | 14.17 | 18.71 | 10.84 | 16.04 | 20.77 | 12.26 | 11.82 | 16.29 | 9.08 |
| Rural Georgia | 15.43 | 17.81 | 13.14 | 17.62 | 20.94 | 14.16 | 11.73 | - | 11.90 |
| Greater Georgia | 14.58 | 18.48 | 11.55 | 15.43 | 19.56 | 12.04 | 11.48 | 14.50 | 9.65 |
| Hawaii | 10.82 | 13.61 | 8.42 | 13.34 | 14.97 | 11.53 | - | - | - |
| Idaho | 16.18 | 20.00 | 12.71 | 16.29 | 20.20 | 12.73 | - | - | - |
| Iowa | 16.23 | 21.40 | 12.03 | 16.33 | 21.52 | 12.09 | 13.66 | 18.34 | 9.68 |
| Kentucky | 15.64 | 19.70 | 12.35 | 15.66 | 19.78 | 12.30 | 12.07 | 14.19 | 10.37 |
| Louisiana | 13.99 | 17.82 | 10.99 | 15.13 | 19.07 | 11.96 | 11.08 | 14.04 | 8.81 |
| Massachusetts | 11.69 | 14.88 | 9.20 | 11.77 | 15.10 | 9.15 | 8.67 | 10.52 | 7.46 |
| New Jersey | 15.48 | 19.82 | 12.17 | 15.85 | 20.22 | 12.49 | 10.81 | 13.60 | 9.06 |
| New Mexico | 13.27 | 16.00 | 10.92 | 13.46 | 16.25 | 11.06 | 10.49 | - | - |
| New York | 16.25 | 21.16 | 12.47 | 17.60 | 22.94 | 13.36 | 10.62 | 13.16 | 9.01 |
| Seattle-Puget Sound | 15.64 | 20.35 | 11.68 | 16.40 | 21.15 | 12.35 | 13.50 | 16.88 | 10.33 |
| Utah | 13.90 | 18.07 | 10.25 | 13.99 | 18.26 | 10.24 | - | - | - |
| SEER 9 Areas ^b | 14.21 | 18.32 | 10.88 | 15.22 | 19.52 | 11.65 | 11.58 | 14.78 | 9.36 |
| SEER 13 Areas ^b | 13.55 | 17.37 | 10.45 | 14.72 | 18.82 | 11.31 | 11.19 | 14.17 | 9.12 |
| SEER 18 Areas ^b | 13.80 | 17.59 | 10.73 | 14.69 | 18.68 | 11.38 | 11.05 | 13.97 | 9.01 |
| SEER 21 Areas ^b | 14.14 | 18.09 | 10.97 | 15.02 | 19.20 | 11.60 | 10.83 | 13.63 | 8.93 |

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia. The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.

- Statistic not shown. Rate based on less than 16 cases for the time interval.