

Table 15.6
Cancer of the Lung and Bronchus (Invasive)

Age-adjusted SEER Incidence^a Rates by Year, Race and Sex

| Year of Diagnosis: | All Races | | | Whites | | | Blacks | | |
|--------------------|-----------|--------|---------|--------|--------|---------|--------|--------|---------|
| | Total | Males | Females | Total | Males | Females | Total | Males | Females |
| 1975-2017 | 61.21 | 81.45 | 46.27 | 61.64 | 80.94 | 47.48 | 75.91 | 112.05 | 50.79 |
| 1975-1979 | 56.22 | 94.32 | 28.33 | 55.60 | 93.59 | 28.39 | 72.01 | 125.85 | 30.72 |
| 1980 | 60.65 | 99.97 | 32.20 | 59.42 | 97.70 | 32.35 | 86.62 | 151.43 | 38.25 |
| 1981 | 62.03 | 99.41 | 35.11 | 61.53 | 98.44 | 35.52 | 82.31 | 142.96 | 37.45 |
| 1982 | 63.29 | 100.69 | 36.73 | 63.05 | 99.68 | 37.68 | 80.37 | 141.06 | 35.68 |
| 1983 | 63.45 | 99.46 | 37.88 | 63.03 | 97.63 | 39.09 | 86.62 | 151.34 | 39.49 |
| 1984 | 65.49 | 102.05 | 39.45 | 64.33 | 99.89 | 39.61 | 92.93 | 158.84 | 44.82 |
| 1985 | 64.61 | 98.62 | 40.23 | 63.97 | 96.87 | 40.85 | 89.56 | 149.68 | 46.04 |
| 1986 | 65.77 | 99.03 | 42.29 | 65.13 | 97.31 | 42.94 | 92.94 | 156.27 | 48.67 |
| 1987 | 67.90 | 101.51 | 44.16 | 68.08 | 100.66 | 45.49 | 84.76 | 143.59 | 43.28 |
| 1988 | 68.06 | 98.70 | 46.34 | 68.26 | 97.92 | 47.73 | 88.24 | 144.93 | 48.26 |
| 1989 | 67.56 | 97.96 | 46.13 | 67.57 | 97.06 | 47.26 | 89.11 | 142.77 | 51.49 |
| 1990 | 68.08 | 96.92 | 47.78 | 68.23 | 96.12 | 48.95 | 86.78 | 137.17 | 52.13 |
| 1991 | 69.19 | 97.21 | 49.57 | 68.96 | 95.89 | 50.61 | 93.12 | 144.76 | 57.02 |
| 1992 | 69.46 | 97.21 | 49.85 | 68.92 | 94.90 | 51.04 | 93.99 | 148.73 | 56.06 |
| 1993 | 67.75 | 93.95 | 49.15 | 67.80 | 92.51 | 50.63 | 86.36 | 135.35 | 52.54 |
| 1994 | 67.17 | 90.93 | 50.54 | 67.28 | 90.13 | 51.61 | 87.16 | 131.39 | 56.53 |
| 1995 | 66.85 | 89.82 | 50.44 | 67.10 | 87.95 | 52.45 | 85.87 | 138.94 | 50.12 |
| 1996 | 66.46 | 87.97 | 51.24 | 66.79 | 86.67 | 52.97 | 84.56 | 128.32 | 55.73 |
| 1997 | 66.63 | 86.31 | 52.59 | 67.19 | 84.81 | 54.92 | 82.27 | 129.54 | 50.49 |
| 1998 | 67.55 | 88.03 | 52.98 | 68.00 | 86.59 | 55.02 | 87.52 | 130.87 | 58.30 |
| 1999 | 65.83 | 84.57 | 52.50 | 66.29 | 83.25 | 54.42 | 84.56 | 122.46 | 58.85 |
| 2000 | 64.14 | 82.12 | 51.22 | 64.66 | 80.78 | 53.22 | 80.71 | 115.16 | 57.75 |
| 2001 | 64.16 | 81.41 | 51.67 | 65.06 | 81.09 | 53.50 | 80.63 | 114.51 | 57.82 |
| 2002 | 64.06 | 80.44 | 52.50 | 64.79 | 79.85 | 54.23 | 81.48 | 113.10 | 60.71 |
| 2003 | 64.77 | 81.05 | 53.10 | 65.51 | 79.56 | 55.65 | 81.20 | 119.40 | 55.81 |
| 2004 | 62.24 | 76.29 | 52.06 | 62.79 | 75.29 | 53.77 | 76.21 | 105.42 | 57.02 |
| 2005 | 63.03 | 75.82 | 53.84 | 64.00 | 75.53 | 55.79 | 74.73 | 97.97 | 59.55 |
| 2006 | 62.30 | 74.30 | 53.46 | 62.83 | 73.10 | 55.34 | 76.75 | 101.10 | 59.96 |
| 2007 | 62.10 | 73.77 | 53.57 | 63.12 | 72.89 | 56.03 | 74.19 | 100.68 | 56.64 |
| 2008 | 60.39 | 72.14 | 51.87 | 61.13 | 70.72 | 54.30 | 75.04 | 103.38 | 56.16 |
| 2009 | 59.99 | 70.67 | 52.27 | 60.69 | 69.76 | 54.13 | 74.27 | 98.01 | 58.63 |
| 2010 | 57.73 | 67.98 | 50.11 | 58.90 | 67.91 | 52.15 | 67.04 | 85.48 | 54.92 |
| 2011 | 56.25 | 65.67 | 49.33 | 57.26 | 65.34 | 51.32 | 65.67 | 86.19 | 52.35 |
| 2012 | 55.26 | 64.00 | 48.76 | 55.70 | 62.92 | 50.36 | 69.38 | 89.08 | 56.07 |
| 2013 | 54.11 | 61.72 | 48.46 | 54.79 | 60.82 | 50.34 | 63.72 | 81.63 | 51.74 |
| 2014 | 53.20 | 60.17 | 48.06 | 54.61 | 60.25 | 50.52 | 60.01 | 74.61 | 50.37 |
| 2015 | 51.86 | 58.81 | 46.64 | 52.76 | 58.08 | 48.78 | 60.01 | 76.06 | 49.43 |
| 2016 | 51.14 | 58.19 | 45.84 | 51.89 | 57.26 | 47.90 | 58.75 | 74.83 | 47.48 |
| 2017 | 49.45 | 55.43 | 45.05 | 50.24 | 54.71 | 47.00 | 56.12 | 71.20 | 46.33 |

Individual rates for diagnosis years 1975-1979 available in web browser version of [Cancer Statistics Review](#).

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.