

Table 15.14
Non-Small Cell Cancer of the Lung and Bronchus (Invasive)

5-Year Relative and Period Survival (Percent) by Race, Sex, Diagnosis Year, Stage and Age

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
5-Year Relative Survival (Percent)									
Year of Diagnosis:									
1975-1977 ^a	16.4	14.8	21.1	16.7	15.0	21.5	13.4	12.4	17.7
1978-1980 ^a	16.7	14.8	21.4	17.0	15.2	21.5	14.2	11.6	22.8
1981-1983 ^a	17.1	15.0	21.7	17.5	15.4	22.1	13.9	12.2	19.2
1984-1986 ^a	16.6	14.4	21.0	17.0	14.7	21.5	13.1	11.9	16.4
1987-1989 ^a	16.8	15.2	19.8	17.3	15.5	20.5	13.2	12.9	13.9
1990-1992 ^a	17.7	15.5	21.1	18.3	16.2	21.7	12.8	11.1	16.2
1993-1995 ^a	18.7	16.1	22.5	19.3	16.5	23.3	15.1	13.3	18.5
1996-1998 ^a	19.4	16.7	22.9	19.9	17.1	23.5	15.3	12.7	19.5
1999-2001 ^a	20.0	16.9	24.1	20.7	17.5	24.7	14.8	12.0	18.9
2002-2004 ^a	19.6	17.0	22.8	20.4	17.7	23.7	15.0	13.9	16.6
2005-2009 ^a	22.2	18.9	26.0	23.1	19.7	26.8	17.4	14.8	20.6
2010-2016 ^a	26.5 ^e	22.1 ^e	31.2 ^e	27.0 ^e	22.5 ^e	31.8 ^e	23.4 ^e	19.0 ^e	28.1 ^e
5-Year Period Survival (Percent) ^{bc}									
2016	26.3	21.8	31.2	26.8	22.1	31.7	22.4	18.4	27.1
Stage Distribution (%) 2010-2016 ^{bd}									
All Stages									
Number of cases	199,070	104,884	94,186	157,707	82,267	75,440	24,448	13,379	11,069
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	20	17	23	21	18	24	16	14	18
Regional	24	24	23	24	25	24	24	23	24
Distant	54	57	52	53	55	51	59	62	56
Unstaged/Unknown	2	2	2	2	2	2	2	2	2
5-Year Relative Survival (Percent), 2010-2016 ^b									
Age at Diagnosis:									
Ages <45	38.7	33.3	43.7	40.6	35.0	45.6	30.1	26.1	34.8
Ages 45-54	26.4	20.8	32.3	27.0	21.2	33.0	23.0	17.5	28.7
Ages 55-64	25.7	21.2	31.4	26.0	21.6	31.6	22.1	18.2	27.8
Ages 65-74	26.7	22.2	31.7	27.2	22.7	32.2	21.6	17.5	26.7
Ages 75+	20.8	17.0	24.5	21.3	17.4	24.9	15.8	11.7	19.7
Ages <65	26.5	21.6	32.4	27.0	22.0	32.8	22.7	18.4	28.4
Ages 65+	24.0	19.9	28.3	24.4	20.3	28.7	19.4	15.4	23.8
Stage ^d :									
All Stages	24.9	20.6	29.7	25.3	21.0	30.1	21.1	17.0	26.0
Localized	63.1	56.7	68.4	63.2	57.0	68.4	58.1	49.1	65.9
Regional	35.4	30.9	40.6	35.2	30.8	40.1	33.3	29.1	38.0
Distant	6.9	5.6	8.6	6.6	5.2	8.2	6.6	5.6	8.0
Unstaged/Unknown	14.8	14.2	15.5	13.7	12.9	14.6	11.5	9.4	13.5

Note: Non-Small Cell Cancer of the Lung and Bronchus includes histologies 8003-8004, 8012-8015, 8021-8022, 8030-8035, 8046, 8050-8052, 8070-8076, 8078, 8082-8084, 8090, 8094, 8120, 8123, 8140-8141, 8143-8145, 8147, 8190, 8200-8201, 8211, 8240-8241, 8243-8246, 8249-8255, 8260, 8290, 8310, 8320, 8323, 8333, 8401, 8430, 8440, 8470-8471, 8480-8481, 8490, 8503, 8507, 8525, 8550, 8560, 8562, 8570-8572, 8574-8576.

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

Based on follow-up of patients into 2017. [Expected survival rates](#) are derived from the U.S. Annual Life Tables.

^b SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG). Expected survival rates are derived from life tables by socio-economic status, geography and race developed by the SEER program.

Based on follow-up of patients into 2017.

^c Period survival provides a 2016 estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three year calendar blocks (2010-2012: 0-1 year survival), (2009-2011: 1-2 year survival), (2008-2010: 2-3 year survival), (2007-2009: 3-4 year survival), (2006-2008: 4-5 years survival).

^d Stage at diagnosis is classified using SEER Summary Stage 2000. Stage distribution percentages may not sum to 100 due to rounding.

^e The difference between 1975-1977 and 2010-2016 is statistically significant ($p < .05$).

^f The standard error is between 5 and 10 percentage points.

^g The standard error is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 25 cases during the time period.