

Table 15.22
Cancer of the Lung And Bronchus (Invasive)

Age-Adjusted SEER Incidence Rates^a
By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER Incidence Rates^a, 2013-2017</u>									
California	41.46	46.38	37.78	42.17	45.83	39.50	50.91	60.37	44.28
Greater Bay Area	39.73	44.47	36.22	39.66	41.73	38.32	54.62	65.57	46.83
San Francisco-Oakland	40.79	45.33	37.48	40.03	41.24	39.39	57.39	69.03	49.35
San Jose-Monterey	37.66	42.82	33.73	39.01	42.57	36.44	36.85	45.54	28.93
Los Angeles	36.31	43.15	31.29	35.32	40.94	31.17	51.02	61.41	44.21
Greater California	44.35	48.43	41.28	45.43	48.77	42.97	49.06	57.01	43.08
Connecticut	59.57	64.97	55.82	60.58	65.18	57.45	54.00	70.64	43.70
Detroit	68.94	78.00	62.55	69.20	76.22	64.40	72.34	90.78	60.04
Georgia	62.79	79.04	50.58	66.44	80.41	55.60	56.01	80.49	40.02
Atlanta	50.75	60.10	44.24	52.06	58.05	47.89	51.85	69.36	41.10
Rural Georgia	64.84	89.10	46.01	63.82	81.46	48.67	66.65	112.06	38.52
Greater Georgia	68.43	87.19	54.00	71.66	88.15	58.67	59.04	87.40	39.65
Hawaii	45.71	57.25	36.25	44.98	49.10	40.66	48.34	44.19	54.12
Idaho	49.49	54.25	45.73	49.46	54.22	45.73	-	-	-
Iowa	63.27	74.73	54.55	63.22	74.44	54.72	78.21	98.53	61.97
Kentucky	91.01	108.95	77.47	92.14	110.21	78.41	84.54	101.84	73.09
Louisiana	66.21	82.62	53.55	66.00	77.91	56.84	68.44	99.13	46.73
Massachusetts	61.21	64.99	58.90	62.83	65.89	61.16	43.75	52.14	38.15
New Jersey	55.32	60.75	51.70	58.19	62.79	55.23	51.82	62.86	45.18
New Mexico	38.52	43.90	34.34	40.22	45.38	36.19	47.06	60.51	35.68
New York	58.67	66.17	53.43	62.27	68.07	58.37	46.49	59.57	38.20
Seattle-Puget Sound	55.15	60.32	51.32	56.42	60.80	53.16	57.71	68.55	49.17
Utah	25.99	30.21	22.52	25.90	30.11	22.43	42.93	41.10	45.43
SEER 9 Areas ^b	51.89	58.77	46.76	52.80	58.14	48.86	59.60	75.48	49.00
SEER 13 Areas ^b	47.77	54.65	42.65	48.56	53.94	44.59	57.30	72.02	47.43
SEER 18 Areas ^b	52.63	60.57	46.63	54.13	60.74	49.15	58.43	76.21	46.23
SEER 21 Areas ^b	54.17	61.72	48.59	56.04	62.22	51.53	54.80	71.19	43.78

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 18 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia

excluding ATL/RG.

The SEER 21 areas comprise the SEER 18 areas plus Idaho, New York and Massachusetts.

- Statistic not shown. Rate based on less than 16 cases for the time interval.