


Table 15.24  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2013-2017

Males and Females

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	40.23	0.05		
High Five States				
Kentucky	63.64	0.50	01(01,01)	58.18 <sup>c</sup>
Arkansas	55.46	0.55	02(02,04)	37.85 <sup>c</sup>
West Virginia	54.90	0.66	03(02,05)	36.45 <sup>c</sup>
Mississippi	54.65	0.57	04(02,05)	35.83 <sup>c</sup>
Tennessee	53.57	0.37	05(03,06)	33.13 <sup>c</sup>
Low Five States				
Hawaii	30.49	0.59	47(46,49)	-24.21 <sup>c</sup>
California	29.55	0.12	48(47,49)	-26.56 <sup>c</sup>
New Mexico	28.48	0.48	49(48,50)	-29.21 <sup>c</sup>
Colorado	27.91	0.32	50(49,50)	-30.63 <sup>c</sup>
Utah	18.55	0.40	51(51,51)	-53.90 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	50.36	0.41	09(06,11)	25.17 <sup>c</sup>	Montana	37.27	0.76	36(30,41)	-7.38 <sup>c</sup>
Alaska	39.61	1.23	30(19,38)	-1.56	Nebraska	39.78	0.61	29(25,34)	-1.13
Arizona	32.88	0.28	45(43,46)	-18.27 <sup>c</sup>	Nevada	41.92	0.53	25(18,27)	4.18
Arkansas	55.46	0.55	02(02,04)	37.85 <sup>c</sup>	New Hampshire	42.49	0.72	23(17,27)	5.61 <sup>c</sup>
California	29.55	0.12	48(47,49)	-26.56 <sup>c</sup>	New Jersey	35.24	0.26	41(38,43)	-12.41 <sup>c</sup>
Colorado	27.91	0.32	50(49,50)	-30.63 <sup>c</sup>	New Mexico	28.48	0.48	49(48,50)	-29.21 <sup>c</sup>
Connecticut	34.68	0.40	42(39,44)	-13.80 <sup>c</sup>	New York	35.35	0.17	40(38,43)	-12.15 <sup>c</sup>
Delaware	45.02	0.87	16(13,22)	11.90 <sup>c</sup>	North Carolina	46.07	0.28	13(13,16)	14.49 <sup>c</sup>
D.C.	33.81	1.04	43(37,47)	-15.96 <sup>c</sup>	North Dakota	36.68	0.94	38(30,43)	-8.84 <sup>c</sup>
Florida	39.14	0.16	31(28,34)	-2.73	Ohio	48.45	0.26	12(10,12)	20.42 <sup>c</sup>
Georgia	42.64	0.29	22(18,25)	5.98 <sup>c</sup>	Oklahoma	52.02	0.48	06(05,08)	29.29 <sup>c</sup>
Hawaii	30.49	0.59	47(46,49)	-24.21 <sup>c</sup>	Oregon	38.58	0.40	34(29,36)	-4.11
Idaho	33.57	0.61	44(40,46)	-16.57 <sup>c</sup>	Pennsylvania	42.32	0.23	24(19,26)	5.17 <sup>c</sup>
Illinois	43.00	0.24	19(17,24)	6.86 <sup>c</sup>	Rhode Island	45.19	0.84	15(13,21)	12.31 <sup>c</sup>
Indiana	50.85	0.37	07(06,10)	26.38 <sup>c</sup>	South Carolina	44.83	0.39	17(13,18)	11.42 <sup>c</sup>
Iowa	43.20	0.48	18(17,25)	7.36 <sup>c</sup>	South Dakota	40.69	0.90	26(19,34)	1.13
Kansas	42.90	0.51	20(17,26)	6.61 <sup>c</sup>	Tennessee	53.57	0.37	05(03,06)	33.13 <sup>c</sup>
Kentucky	63.64	0.50	01(01,01)	58.18 <sup>c</sup>	Texas	35.82	0.17	39(37,41)	-10.96 <sup>c</sup>
Louisiana	49.65	0.44	10(07,12)	23.41 <sup>c</sup>	Utah	18.55	0.40	51(51,51)	-53.90 <sup>c</sup>
Maine	48.79	0.72	11(08,12)	21.26 <sup>c</sup>	Vermont	42.80	1.03	21(14,29)	6.38 <sup>c</sup>
Maryland	38.81	0.34	32(29,35)	-3.55	Virginia	40.40	0.30	27(25,30)	0.40
Massachusetts	38.69	0.31	33(29,35)	-3.84	Washington	37.29	0.31	35(34,38)	-7.33 <sup>c</sup>
Michigan	45.52	0.28	14(13,17)	13.14 <sup>c</sup>	West Virginia	54.90	0.66	03(02,05)	36.45 <sup>c</sup>
Minnesota	36.92	0.34	37(34,39)	-8.23 <sup>c</sup>	Wisconsin	40.14	0.34	28(26,31)	-0.23
Mississippi	54.65	0.57	04(02,05)	35.83 <sup>c</sup>	Wyoming	32.20	1.00	46(41,48)	-19.96 <sup>c</sup>
Missouri	50.81	0.37	08(06,10)	26.29 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.