


Table 15.25  
Cancer of the Lung and Bronchus (Invasive)

Age-Adjusted Cancer Death<sup>a</sup> Rates By State, All Races, 2013-2017

Males

State	Rate	SE	Rank(CI) <sup>d</sup>	PD
TOTAL U.S.	49.33	0.08		
High Five States				
Kentucky	80.77	0.85	01(01,01)	63.74 <sup>c</sup>
Mississippi	75.51	1.03	02(02,03)	53.07 <sup>c</sup>
Arkansas	71.85	0.95	03(02,05)	45.66 <sup>c</sup>
Tennessee	69.98	0.65	04(03,06)	41.86 <sup>c</sup>
West Virginia	69.40	1.11	05(03,06)	40.68 <sup>c</sup>
Low Five States				
Wyoming	36.43	1.58	47(40,49)	-26.16 <sup>c</sup>
California	34.99	0.20	48(47,49)	-29.07 <sup>c</sup>
New Mexico	33.42	0.78	49(47,50)	-32.26 <sup>c</sup>
Colorado	30.94	0.52	50(49,50)	-37.27 <sup>c</sup>
Utah	22.31	0.64	51(51,51)	-54.78 <sup>c</sup>

State	Rate	SE	Rank(CI) <sup>d</sup>	PD	State	Rate	SE	Rank(CI) <sup>d</sup>	PD
Alabama	68.39	0.74	06(04,06)	38.64 <sup>c</sup>	Montana	39.42	1.15	43(38,47)	-20.08 <sup>c</sup>
Alaska	45.79	1.98	32(22,42)	-7.17 <sup>c</sup>	Nebraska	48.38	1.01	26(22,32)	-1.92
Arizona	38.21	0.45	46(42,47)	-22.54 <sup>c</sup>	Nevada	45.96	0.81	31(26,36)	-6.83 <sup>c</sup>
Arkansas	71.85	0.95	03(02,05)	45.66 <sup>c</sup>	New Hampshire	48.31	1.15	27(22,34)	-2.07
California	34.99	0.20	48(47,49)	-29.07 <sup>c</sup>	New Jersey	41.04	0.43	40(39,43)	-16.79 <sup>c</sup>
Colorado	30.94	0.52	50(49,50)	-37.27 <sup>c</sup>	New Mexico	33.42	0.78	49(47,50)	-32.26 <sup>c</sup>
Connecticut	39.67	0.65	42(40,46)	-19.59 <sup>c</sup>	New York	42.54	0.29	39(35,40)	-13.75 <sup>c</sup>
Delaware	55.09	1.46	16(13,23)	11.67 <sup>c</sup>	North Carolina	60.03	0.50	12(11,14)	21.70 <sup>c</sup>
D.C.	40.25	1.76	41(32,47)	-18.41 <sup>c</sup>	North Dakota	45.56	1.56	33(24,40)	-7.65 <sup>c</sup>
Florida	47.47	0.27	29(25,31)	-3.77	Ohio	60.42	0.44	11(10,14)	22.48 <sup>c</sup>
Georgia	56.73	0.52	15(14,17)	15.00 <sup>c</sup>	Oklahoma	64.93	0.81	07(07,10)	31.63 <sup>c</sup>
Hawaii	38.97	0.99	45(40,47)	-20.99 <sup>c</sup>	Oregon	43.45	0.63	36(33,40)	-11.92 <sup>c</sup>
Idaho	39.05	0.97	44(40,47)	-20.84 <sup>c</sup>	Pennsylvania	52.50	0.38	20(18,23)	6.44 <sup>c</sup>
Illinois	52.37	0.41	21(18,24)	6.15 <sup>c</sup>	Rhode Island	53.19	1.39	19(15,25)	7.82 <sup>c</sup>
Indiana	64.06	0.63	09(07,10)	29.87 <sup>c</sup>	South Carolina	58.86	0.68	14(11,15)	19.32 <sup>c</sup>
Iowa	54.04	0.80	18(15,22)	9.55 <sup>c</sup>	South Dakota	49.95	1.48	24(17,32)	1.25
Kansas	51.91	0.84	22(17,25)	5.23 <sup>c</sup>	Tennessee	69.98	0.65	04(03,06)	41.86 <sup>c</sup>
Kentucky	80.77	0.85	01(01,01)	63.74 <sup>c</sup>	Texas	45.32	0.29	34(30,35)	-8.12 <sup>c</sup>
Louisiana	64.52	0.77	08(07,10)	30.80 <sup>c</sup>	Utah	22.31	0.64	51(51,51)	-54.78 <sup>c</sup>
Maine	59.20	1.19	13(10,16)	20.02 <sup>c</sup>	Vermont	49.65	1.66	25(17,34)	0.65
Maryland	46.29	0.58	30(27,35)	-6.17 <sup>c</sup>	Virginia	50.50	0.51	23(21,26)	2.38
Massachusetts	44.86	0.51	35(30,37)	-9.07 <sup>c</sup>	Washington	43.01	0.50	37(34,40)	-12.80 <sup>c</sup>
Michigan	54.32	0.45	17(16,20)	10.12 <sup>c</sup>	West Virginia	69.40	1.11	05(03,06)	40.68 <sup>c</sup>
Minnesota	42.62	0.55	38(35,41)	-13.61 <sup>c</sup>	Wisconsin	48.05	0.56	28(24,31)	-2.60
Mississippi	75.51	1.03	02(02,03)	53.07 <sup>c</sup>	Wyoming	36.43	1.58	47(40,49)	-26.16 <sup>c</sup>
Missouri	62.60	0.62	10(08,11)	26.89 <sup>c</sup>					



<sup>a</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>b</sup> Difference between state rate and total U.S. rate is statistically significant ( $p \leq .0002$ ).

<sup>c</sup> Absolute percent difference between state rate and total U.S. rate is 5% or more.

<sup>d</sup> Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.