

Table 22.21
Cancer of the Pancreas (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2017
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis		0 to 5	> 5 to 10	> 10 to 15	> 15 to 20	> 20 to 25	<=25 ^b		> 25 ^d	Complete ^e
Race	Sex									
All Races	Both Sexes	57,729	10,952	4,618	1,929	1,010	76,239		2,731	78,969
	Males	29,053	5,759	1,879	922	286	37,899		1,296	39,194
	Females	28,677	5,193	2,739	1,007	724	38,340		1,435	39,775
White	Both Sexes	47,797	8,888	3,811	1,535	804	62,836		2,220	65,055
	Males	24,315	4,813	1,594	812	246	31,779		1,120	32,898
	Females	23,482	4,075	2,218	724	558	31,057		1,100	32,157
Black	Both Sexes	6,894	1,388	510	247	114	9,153		861	10,015
	Males	3,194	619	177	70	9	4,069		404	4,473
	Females	3,700	770	333	177	105	5,085		457	5,541
Asian/ Pacific Islander	Both Sexes	2,378	526	211	85	52	3,252		+	+
	Males	1,159	241	85	20	13	1,518		+	+
	Females	1,219	285	127	64	38	1,734		+	+
Hispanic	Both Sexes	4,702	805	408	181	108	6,204		+	+
	Males	2,021	361	133	71	5	2,591		+	+
	Females	2,681	445	274	110	103	3,613		+	+

Estimated prevalence percent^a on January 1, 2017, of the SEER population diagnosed in the previous 25 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence		Age Specific (Crude)										Age-Adjusted ^c
		All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
Race	Sex											
All Races	Both Sexes	0.0235	0.0000	0.0004	0.0011	0.0033	0.0103	0.0283	0.0650	0.1048	0.0977	0.0196
	Males	0.0237	0.0001	0.0003	0.0004	0.0026	0.0087	0.0291	0.0723	0.1202	0.1108	0.0211
	Females	0.0233	-	0.0005	0.0018	0.0040	0.0118	0.0275	0.0585	0.0919	0.0896	0.0184
White	Both Sexes	0.0248	0.0000	0.0004	0.0012	0.0031	0.0097	0.0272	0.0643	0.1057	0.0980	0.0194
	Males	0.0253	0.0001	0.0003	0.0004	0.0026	0.0082	0.0281	0.0722	0.1202	0.1132	0.0210
	Females	0.0243	-	0.0004	0.0021	0.0037	0.0112	0.0263	0.0569	0.0933	0.0884	0.0181
Black	Both Sexes	0.0201	-	0.0003	0.0006	0.0039	0.0130	0.0366	0.0770	0.1050	0.0895	0.0216
	Males	0.0186	-	0.0003	0.0005	0.0027	0.0098	0.0382	0.0814	0.1264	0.0690	0.0219
	Females	0.0215	-	0.0003	0.0007	0.0050	0.0159	0.0352	0.0735	0.0900	0.0997	0.0213
Asian/ Pacific Islander	Both Sexes	0.0156	0.0003	0.0009	0.0007	0.0039	0.0101	0.0224	0.0497	0.0814	0.0921	0.0164
	Males	0.0152	0.0005	0.0005	0.0004	0.0028	0.0112	0.0218	0.0488	0.0987	0.1006	0.0176
	Females	0.0160	-	0.0014	0.0010	0.0048	0.0091	0.0230	0.0505	0.0675	0.0865	0.0156
Hispanic	Both Sexes	0.0107	-	0.0004	0.0014	0.0030	0.0086	0.0211	0.0516	0.0801	0.0840	0.0158
	Males	0.0088	-	0.0005	0.0004	0.0014	0.0065	0.0181	0.0490	0.0875	0.0799	0.0149
	Females	0.0126	-	0.0004	0.0025	0.0047	0.0108	0.0241	0.0540	0.0744	0.0866	0.0167

^a US 2017 cancer prevalence counts are based on 2017 cancer prevalence proportions from the SEER 13 Areas (excluding the Alaska Native Registry) and 1/1/2017 US population estimates based on the average of 2016 and 2017 population estimates from the US Bureau of the Census.

^b Prevalence was calculated using the first invasive tumor for each cancer site diagnosed during the previous 25 years. Maximum limited-duration prevalence is 25 years for 1992-2017 data.

^c Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^d Cases diagnosed more than 25 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000).

^e Complete prevalence is obtained by summing <=25 and > 25.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.