

Table 23.7  
Cancer of the Prostate (Invasive)

SEER Incidence<sup>a</sup> and U.S. Death<sup>b</sup> Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence <sup>a</sup>			U.S. Mortality <sup>b</sup>		
	All Races Males	White Males	Black Males	All Races Males	White Males	Black Males
<u>Age-Adjusted Rates, 2013-2017</u>						
Age at Diagnosis/Death:						
All ages	109.8	102.3	175.2	19.1	17.9	37.9
Under 65	43.9	39.4	82.0	1.7	1.4	4.0
65 and over	565.3	536.8	819.7	139.8	132.0	272.2
All ages (WHO world std) <sup>c</sup>	83.6	77.4	136.9	8.5	7.8	17.6
<u>Age-Specific Rates, 2013-2017</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	-	-	-	-	-	-
30-34	0.1	-	-	-	-	-
35-39	0.6	0.4	1.3	-	-	-
40-44	7.1	5.8	17.5	0.1	0.1	0.3
45-49	31.4	26.0	77.2	0.8	0.7	1.6
50-54	114.1	100.1	229.1	3.1	2.6	6.9
55-59	249.1	223.9	458.7	8.5	7.1	20.5
60-64	418.3	384.0	719.4	20.0	17.1	48.5
65-69	631.1	588.5	1,014.0	38.7	34.2	92.3
70-74	652.8	620.7	946.9	69.3	63.1	157.3
75-79	577.3	547.2	797.3	123.2	114.2	261.0
80-84	436.6	422.0	550.1	216.6	205.0	422.7
85+	368.1	364.3	478.7	447.8	435.9	751.9

<sup>a</sup> SEER 21 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

<sup>c</sup> Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.