


Table 24.18
Cancer of the Stomach (Invasive)

Age-Adjusted Cancer Death^a Rates By State, All Races, 2013-2017

Males and Females

State	Rate	SE	Rank(CI) ^d	PD
TOTAL U.S.	3.06	0.01		
High Five States				
D.C.	4.85	0.40	01(01,05)	58.69 ^c
Alaska	4.51	0.40	02(01,11)	47.45 ^c
Hawaii	4.25	0.22	03(01,06)	39.22 ^c
California	3.94	0.04	04(02,05)	29.04 ^c
Mississippi	3.60	0.15	05(03,16)	17.79 ^c
Low Five States				
Nebraska	2.03	0.14	47(36,51)	-33.43 ^c
Maine	2.00	0.15	48(37,51)	-34.40 ^c
New Hampshire	2.00	0.16	49(36,51)	-34.49 ^c
Iowa	1.99	0.10	50(41,51)	-34.76 ^c
Wyoming	1.49	0.22	51(44,51)	-51.37 ^c

State	Rate	SE	Rank(CI) ^d	PD	State	Rate	SE	Rank(CI) ^d	PD
Alabama	3.30	0.11	13(05,20)	7.95	Montana	2.12	0.18	46(27,51)	-30.51 ^c
Alaska	4.51	0.40	02(01,11)	47.45 ^c	Nebraska	2.03	0.14	47(36,51)	-33.43 ^c
Arizona	2.59	0.08	33(23,41)	-15.11 ^c	Nevada	2.74	0.14	26(15,42)	-10.48
Arkansas	2.81	0.13	23(14,38)	-8.10	New Hampshire	2.00	0.16	49(36,51)	-34.49 ^c
California	3.94	0.04	04(02,05)	29.04 ^c	New Jersey	3.41	0.08	11(05,16)	11.74
Colorado	2.45	0.10	37(25,46)	-19.69 ^c	New Mexico	3.47	0.17	10(03,20)	13.73
Connecticut	3.18	0.12	14(07,24)	3.92	New York	3.59	0.06	07(04,11)	17.42 ^c
Delaware	3.03	0.23	19(05,41)	-0.92	North Carolina	2.92	0.07	22(15,28)	-4.30
D.C.	4.85	0.40	01(01,05)	58.69 ^c	North Dakota	2.28	0.24	43(18,51)	-25.36 ^c
Florida	2.72	0.04	28(22,34)	-10.97	Ohio	2.64	0.06	30(23,38)	-13.52
Georgia	3.15	0.08	15(10,22)	2.96	Oklahoma	2.69	0.11	29(18,41)	-11.93
Hawaii	4.25	0.22	03(01,06)	39.22 ^c	Oregon	2.46	0.10	36(24,46)	-19.47 ^c
Idaho	2.31	0.16	42(24,50)	-24.50 ^c	Pennsylvania	2.62	0.06	31(23,38)	-14.15
Illinois	3.31	0.07	12(08,17)	8.22	Rhode Island	3.13	0.22	16(05,36)	2.38
Indiana	2.50	0.08	34(25,44)	-18.11 ^c	South Carolina	3.60	0.11	06(04,14)	17.74 ^c
Iowa	1.99	0.10	50(41,51)	-34.76 ^c	South Dakota	2.31	0.22	41(19,50)	-24.36 ^c
Kansas	2.21	0.12	45(33,50)	-27.58 ^c	Tennessee	2.72	0.09	27(19,37)	-10.87
Kentucky	2.95	0.11	21(13,31)	-3.54	Texas	3.56	0.05	08(04,12)	16.43 ^c
Louisiana	3.49	0.12	09(04,17)	14.22	Utah	2.27	0.14	44(28,50)	-25.63 ^c
Maine	2.00	0.15	48(37,51)	-34.40 ^c	Vermont	2.39	0.25	39(15,50)	-21.67 ^c
Maryland	3.04	0.10	18(12,26)	-0.48	Virginia	3.02	0.08	20(13,26)	-1.27
Massachusetts	3.08	0.09	17(11,24)	0.78	Washington	2.75	0.08	25(19,37)	-10.07
Michigan	2.76	0.07	24(19,34)	-9.72	West Virginia	2.61	0.15	32(17,45)	-14.55
Minnesota	2.34	0.09	40(32,47)	-23.37 ^c	Wisconsin	2.48	0.09	35(26,45)	-19.00 ^c
Mississippi	3.60	0.15	05(03,16)	17.79 ^c	Wyoming	1.49	0.22	51(44,51)	-51.37 ^c
Missouri	2.40	0.08	38(30,46)	-21.58 ^c					



^a US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p < .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

^d Rank of state level death rate and the 95% confidence intervals (low, high) for the rank.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

- Statistic not shown. Rate based on less than 16 cases for the time interval.