Table 29.5

Age-Adjusted SEER Cancer Incidence Trends, 1975-2017^a By International Classification of Childhood Cancer(ICCC)^b Selected Group and Subgroup^c Excluding benign brain and myelodysplastic syndromes All Races, Females

	Ages 0-14				Ages 0-19			
ICCC Group and Subgroup	% Change 1975-2017	APC 1975-2017	APC 1975-1996	APC 1997-2017	% Change 1975-2017	APC 1975-2017	APC 1975-1996	APC 1997-2017
All ICCC Groups Combined	48.9	0.7*	0.5*	0.7*	43.6	0.7*	0.5*	0.9*
I Leukemia	88.1	0.6*	0.6	0.7*	76.1	0.7*	0.7	0.7*
I(a) Lymphoid leukemia	100.1	0.7*	0.8	0.6	86.4	0.8*	0.7	0.6*
I(b) Acute myeloid leukemia	99.5	0.7*	1.1	1.0	81.8	0.9*	1.5*	1.0
II Lymphomas and reticuloendothelial neo.	8.4	0.5*	-0.6	0.9	10.1	0.2	0.2	0.5
II(a) Hodgkin lymphoma	-7.8	-0.5	-0.9	0.0	-8.2	-0.5*	-0.2	-0.3
II(b,c,e) Non-Hodgkin lymphoma	41.9	1.7*	-0.1	1.8	58.4	1.7*	1.0	1.8*
III CNS & misc intracranial & intraspinal neo.	43.2	0.8*	1.2*	0.4	42.3	0.8*	1.1*	0.7
III(a) Ependymomas and choroid plexus tumors	248.2	1.2*	1.3	1.8	194.1	1.3*	1.2	2.1*
III(b) Astrocytoma	73.0	1.0*	1.8*	0.6	59.0	0.9*	1.6*	1.1
III(c) Intracranial and intraspinal embryonal tum.		0.9*	1.7*	0.1	108.4	1.0*	1.8*	0.0
III(d) Other gliomas	-47.2	0.2	-1.2	-0.3	-28.5	0.5	-0.7	-0.3
IV Neuroblastoma and other peripheral nervous cell tumors	-33.0	0.2	-0.3	-0.5	-33.0	0.2	-0.4	-0.4
IV(a) Neuroblastoma and ganglioneuroblastoma	-36.1	0.2	-0.3	-0.5	-36.1	0.2	-0.4	-0.4
V Retinoblastoma	31.8	0.1	0.6	-0.1	31.8	0.1	0.6	-0.1
VI Renal tumors	59.6	0.1	0.0	0.4	50.9	0.2	0.3	0.6
VI(a) Nephroblastoma and other nonepithelial renal tumors	53.7	0.1	0.0	0.5	53.7	0.0	0.1	0.5
VII Hepatic tumors	376.1	2.1*	0.0	1.7	616.5	2.0*	0.1	1.6
VII(a) Hepatoblastoma	-	-	-	1.6	-	-	-	1.7
VIII Malignant bone tumors	42.5	0.1	-0.2	1.2	-4.7	-0.1	-0.4	0.6
VIII(a) Osteosarcoma	120.5	0.7	1.1	1.8	9.4	0.3	0.3	1.3
VIII(c) Ewing tum. and related sarcomas of bone	-12.0	-	-2.1	-	26.7	-0.5	-1.2	0.5
IX Soft tissue and other extraosseous sarcomas	34.4	0.3	0.7	-0.3	36.1	0.3	0.3	-0.6
IX(a) Rhabdomyosarcoma	-3.9	0.0	0.4	-0.3	5.4	0.0	0.7	-0.9
X Germ cell & trophoblastic tum. & neo. of gonads	81.7	1.0*	1.5	1.3	-7.4	0.2	0.7	0.7
X(a) Intracranial and intraspinal germ-cell tum.	301.5	-	-	1.9	301.5	-	-	1.3
X(c) Malignant gonadal germ cell tum. (ovary)	50.3	0.4	-0.7	1.2	5.5	0.1	1.0	0.0
XI Other malignant epithelial neo. and melanomas $^{\mathrm{d}}$	105.9	1.8*	0.4	3.2*	120.4	1.7*	0.6	3.0*
XI(b) Thyroid carcinoma	24.7	1.7*	0.2	3.4*	182.8	2.5*	1.0	4.3*
XI(d) Malignant melanoma	115.3	-	-	-0.2	66.5	0.5	1.1	-1.4

The APC is the Annual Percent Change over the time interval.

а SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). b International Classification of Childhood Cancer is based on ICD-0-3. International Classification of Childhood Cancer is based on ICD-0-3. Steliarova-Foucher E, Colombet M, Ries LAG, Moreno F, Dolya A, Shin HY, Hesseling P, Stiller CA. International Incidence of Childhood Cancer, Volume III. Lyon: International Agency for Research on Cancer, In press. с

Classifications are shown for invasive cases only except as noted. For ICCC groups and subgroups that include in situ behavior, only invasive cases are shown except as noted. d

Includes in situ urinary bladder tumors.

- * The APC is significantly different from zero (p<.05)
- # The APC for 1997-2017 is significantly different from the APC for 1975-1996 (p<.05).
- @ The APC for 1997-2017 is significantly different from the APC for 1975-1996 (p<.10).
- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval. _ Neoplasms are abbreviated as neo. Tumors are abbreviated as tum.