All Races

Table 1.23 Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity Both Sexes

White

Black

	Rate ^b 2014-2018	APC° 2009-2018		Rate ^b 2014-2018	APC° 2009-2018		Rate ^b 2014-2018	APC° 2009-2018
All Sites	450.5	-0.9*	All Sites	460.9	-1.0*	All Sites	445.2	
Breast	69.1	0.1	Breast	69.7	-0.1	Prostate ^f	74.8	-3.5*
Lung and Bronchus	53.1	-2.2*	Lung and Bronchus	54.9	-2.2*	Breast	71.1	0.0
Prostate ^f	50.9	-3.2*	Prostate ^f	48.2	-3.6*	Lung and Bronchus	53.7	-2.8*
Colon and Rectum	37.8	-1.9*	Colon and Rectum	37.4	-1.8*	Colon and Rectum	42.6	-2.9*
Melanoma of the Skin	22.8	1.0*	Melanoma of the Skin	27.4	1.1*	Kidney and Renal Pelvis	17.4	-0.2
Urinary Bladder	19.7	-1.5*	Urinary Bladder	21.7	-1.5*	Corpus and Uterus, NOSf	16.3	1.5*
Non-Hodgkin Lymphoma	19.6	-0.7*	Non-Hodgkin Lymphoma	20.6	-0.8*	Pancreas	15.5	0.2
Kidney and Renal Pelvis		0.6*	Kidney and Renal Pelvis		0.6*	Non-Hodgkin Lymphoma	14.7	-0.7*
Thyroid	15.5	0.1	Thyroid	16.2	-0.1	Myeloma	13.8	-0.1
Corpus and Uterus, NOS ^f		0.8*	Leukemia	15.2	-0.7	Urinary Bladder	11.5	-1.2*
Leukemia	14.3	-0.6	Corpus and Uterus, NOSf	15.0	0.5*	Leukemia	11.0	-0.2
Pancreas	13.2	0.6*	Pancreas	13.3	0.7*	Liver & IBD ^g	10.6	0.0
Oral Cavity and Pharynx		0.5*	Oral Cavity and Pharynx		0.6*	Stomach	10.0	-2.3*
Liver & IBD ^g	9.0	1.0*	Liver & IBD ^g	8.2	1.6*	Thyroid	9.4	0.0
Stomach	7.2	-1.0*	Brain and ONS ^g	7.1	-0.5*	Oral Cavity and Pharynx		-1.3*
				–				
Asian/Pacific Islander			American Indian/Alaska Native ^d			Hispanic ^e		
	Rate ^b	APC ^c		Rate ^b	APCc		Rateb	APC ^c
	2014-2018			2014-2018			2014-2018	
All Sites	307.9	-0.7*	All Sites	315.1	-1.0*	All Sites	355.3	-1.0*
Breast	57.5	1.5*	Breast	43.8	-0.7	Breast	53.7	0.5
Lung and Bronchus	35.7	-1.1*	Colon and Rectum	38.5	-1.7*	Prostate ^f	40.4	-4.7*
Colon and Rectum	31.3	-2.4*	Lung and Bronchus	37.7	-1.5*	Colon and Rectum	33.7	-1.5*
Prostate ^f	25.5	-3.7*	Prostate ^f	23.6	-5.4*	Lung and Bronchus	28.6	-2.2*
Thyroid	15.8	0.7	Kidney and Renal Pelvis		0.7	Non-Hodgkin Lymphoma	18.1	-0.6*
Non-Hodgkin Lymphoma	13.6	-0.3	Liver & IBD ^g	14.6	0.9	Kidney and Renal Pelvis		1.3*
Liver & IBD ^g	12.7	-2.6*	Non-Hodgkin Lymphoma	10.8	-3.5*	Thyroid	14.3	1.6*
Corpus and Uterus, NOS ^f		2.0*	Corpus and Uterus, NOS ^f	10.6	-0.1	Liver & IBD ^g	13.8	0.1
Stomach	10.1	-3.1*	Pancreas	10.4	1.4	Corpus and Uterus, NOSf	13.4	1.9*
Pancreas	10.0	0.3	Urinary Bladder	9.8	2.2	Pancreas	11.8	0.5
Urinary Bladder	8.9	-0.9*	Thyroid	9.6	2.2*	Leukemia	11.0	-0.7
Kidney and Renal Pelvis	8.8	0.9	Leukemia	8.9	0.9	Urinary Bladder	11.0	-1.6*
Oral Cavity and Pharynx		0.7	Oral Cavity and Pharynx		-1.7	Stomach	10.4	-1.4*
Leukemia	8.2	-0.3	Stomach	7.8	-3.3	Oral Cavity and Pharynx		-0.6
Ovary ^{fh}	5.1	-0.8	Myeloma	5.8	3.2	Myeloma	6.9	-0.1

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

- Top 15 cancer sites selected based on 2014-2018 age-adjusted rates for the race/ethnic group.
- Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval.
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.
- Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.
- Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. f
- The rates for sex-specific cancer sites are calculated using the population for both sexes combined.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- The APC is significantly different from zero (p<.05).
- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.