

Table 1.25  
Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites<sup>a</sup> by Race/Ethnicity

## Females

	All Races		White		Black	
	Rate <sup>b</sup> 2014-2018	APC <sup>c</sup> 2009-2018	Rate <sup>b</sup> 2014-2018	APC <sup>c</sup> 2009-2018	Rate <sup>b</sup> 2014-2018	APC <sup>c</sup> 2009-2018
All Sites	425.6	-0.3*	439.5	-0.3*	396.4	-0.6*
Breast	129.1	0.3	131.8	0.1	124.7	0.2
Lung and Bronchus	47.9	-1.5*	50.8	-1.4*	43.3	-2.1*
Colon and Rectum	33.3	-1.9*	33.0	-1.7*	37.5	-2.9*
Corpus and Uterus, NOS	28.1	1.0*	28.6	0.7*	28.5	1.7*
Thyroid	22.8	0.0	23.9	-0.3	14.3	0.4
Melanoma of the Skin	18.0	1.1*	22.1	1.2*	14.0	0.1
Non-Hodgkin Lymphoma	16.2	-0.7*	17.0	-0.8*	12.4	-0.3
Pancreas	11.8	0.6*	11.8	-0.7	12.1	-0.1
Kidney and Renal Pelvis	11.2	0.6*	11.7	0.6*	12.1	0.0
Leukemia	11.1	-0.6	11.6	0.6*	9.1	0.0
Ovary <sup>g</sup>	10.9	-2.3*	11.3	-2.6*	9.0	-1.7*
Urinary Bladder	8.5	-1.5*	9.3	-1.5*	8.6	-1.5*
Cervix Uteri	7.5	-0.3	7.3	-0.3	7.7	-2.3*
Oral Cavity and Pharynx	6.5	0.2	6.7	0.3*	6.4	-1.1*
Myeloma	5.7	0.3	6.0	-0.6	5.3	0.1
	Asian/Pacific Islander		American Indian/Alaska Native <sup>d</sup>		Hispanic <sup>e</sup>	
	Rate <sup>b</sup> 2014-2018	APC <sup>c</sup> 2009-2018	Rate <sup>b</sup> 2014-2018	APC <sup>c</sup> 2009-2018	Rate <sup>b</sup> 2014-2018	APC <sup>c</sup> 2009-2018
All Sites	311.5	0.1	311.1	-0.5	348.4	0.2
Breast	105.1	1.5*	81.7	-0.7	100.3	0.8*
Lung and Bronchus	28.7	-0.3	37.5	-0.1	29.3	-1.0*
Colon and Rectum	26.5	-2.7*	35.5	0.5	25.3	2.2*
Thyroid	22.8	0.4	20.0	0.0	24.6	-1.4*
Corpus and Uterus, NOS	22.3	2.0*	14.7	2.6*	22.5	1.6*
Non-Hodgkin Lymphoma	11.3	-0.2	14.0	0.3	16.0	-0.4
Ovary <sup>g</sup>	9.4	-0.7	10.1	-3.6	12.1	1.3*
Pancreas	9.3	0.2	8.6	0.3	11.1	0.9*
Stomach	7.6	-3.3*	8.5	-1.8	10.3	-0.8
Liver & IBD <sup>f</sup>	7.1	-2.5*	8.2	-0.2	9.3	-0.9
Leukemia	6.6	-0.3	7.6	-0.6	9.3	-0.6
Cervix Uteri	6.4	-0.6	6.6	-1.4	8.4	-0.6
Kidney and Renal Pelvis	5.9	1.5*	5.8	-2.8	8.2	1.3
Oral Cavity and Pharynx	5.6	0.9	5.0	-0.6	6.0	0.5
Urinary Bladder	3.7	-1.6*	4.9	0.2	5.0	-1.4*

Source: SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

<sup>a</sup> Top 15 cancer sites selected based on 2014-2018 age-adjusted rates for the race/ethnic group.

<sup>b</sup> Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>c</sup> The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>d</sup> Rates for American Indian/Alaska Native are based on the Purchased/Referred Care Delivery Area (PRCDA) counties.

<sup>e</sup> Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives.

Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

<sup>f</sup> IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.

<sup>g</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

\* The APC is significantly different from zero (p<.05).

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.