

Table 3.21

SEER<sup>a</sup> 5-Year Relative Survival for the Top 5 Cancer Sites<sup>b</sup> by Age  
All Races, Males, 1975-2017

	5-year Relative Survival (Percent)					
	1975-2017	1975-1983	1984-1992	1993-2001	2002-2010	2011-2017
<u>Ages 15-19</u>						
All Malignant Cancers	75.5	63.7	72.0	75.5	81.7	84.0
Hodgkin Lymphoma	91.8	86.0	90.1	93.8	96.6	94.1
Testis	90.7	79.2	92.0	92.5	93.9	94.0
Brain and Other Nervous System	71.8	59.7	70.5	74.5	75.0	79.8
Non-Hodgkin Lymphoma	73.4	49.6	67.2	71.7	82.3	87.5
Bones and Joints	59.4	42.6	59.9	60.5	67.9	66.3
<u>Ages 20-24</u>						
All Malignant Cancers	78.1	69.5	73.2	79.3	82.8	86.0
Testis	93.2	84.5	92.0	95.7	95.9	97.2
Hodgkin Lymphoma	89.9	83.2	87.9	90.3	94.7	96.9
Non-Hodgkin Lymphoma	68.5	56.7	52.8	67.0	80.4	79.8
Melanoma of the Skin	87.5	76.1	85.3	91.9	91.4	96.2
Brain and Other Nervous System	67.0	54.7	63.0	72.6	72.6	74.4
<u>Ages 25-29</u>						
All Malignant Cancers	75.3	71.5	63.4	75.7	83.2	85.8
Testis	95.4	89.1	94.6	97.3	97.1	97.4
Melanoma of the Skin	89.2	82.5	87.7	90.4	93.0	93.8
Hodgkin Lymphoma	89.2	84.5	85.7	90.0	93.3	96.2
Non-Hodgkin Lymphoma	62.3	59.0	41.7	57.7	79.0	83.1
Kaposi sarcoma	18.0	29.7	7.5	20.2	56.7	63.1
<u>Ages 30-34</u>						
All Malignant Cancers	70.1	68.2	54.9	68.1	81.7	84.8
Testis	95.9	92.0	95.7	95.5	97.4	97.3
Melanoma of the Skin	89.1	79.3	87.0	91.2	93.7	94.4
Kaposi sarcoma	20.0	24.7	7.6	29.6	64.4	69.8
Non-Hodgkin Lymphoma	57.7	69.2	36.5	48.6	78.6	84.3
Hodgkin Lymphoma	88.0	79.7	83.9	88.3	93.5	95.1
<u>Ages 35-39</u>						
All Malignant Cancers	66.2	59.7	50.9	65.4	78.1	81.1
Melanoma of the Skin	89.1	82.1	85.7	90.2	92.6	95.1
Testis	96.3	91.9	95.7	97.1	97.1	96.6
Non-Hodgkin Lymphoma	59.3	62.2	41.7	51.6	74.7	83.0
Kaposi sarcoma	20.7	14.1	7.0	30.5	66.5	83.8
Colon and Rectum	63.1	54.1	54.5	62.5	69.5	68.4

<sup>a</sup> SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

<sup>b</sup> Top 5 cancer sites selected based on 1975-2017 SEER 9 Incidence rates for the age group.

- Statistic could not be calculated due to fewer than 25 cases during the time period.