

Table X-4
KAPOSI SARCOMA

AGE ADJUSTED SEER INCIDENCE^a RATES BY YEAR, RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
YEAR OF DIAGNOSIS:									
1975	0.2	0.6	-	0.2	0.6	-	-	-	-
1976	0.2	0.5	-	0.2	0.5	-	-	-	-
1977	0.3	0.5	-	0.3	0.6	-	-	-	-
1978	0.2	0.3	0.2	0.2	0.3	-	-	-	-
1979	0.2	0.3	-	0.2	0.3	-	-	-	-
1980	0.3	0.6	-	0.3	0.6	-	-	-	-
1981	0.3	0.5	-	0.3	0.6	-	-	-	-
1982	0.7	1.2	0.2	0.7	1.3	0.2	-	-	-
1983	1.1	2.2	-	1.2	2.4	-	-	-	-
1984	1.8	3.6	-	2.0	3.9	-	1.3	2.9	-
1985	2.5	4.9	0.2	2.8	5.5	0.2	1.1	2.2	-
1986	3.3	6.5	-	3.7	7.3	-	1.6	3.6	-
1987	3.8	7.6	-	4.3	8.5	-	2.1	4.3	-
1988	4.4	8.9	0.1	4.8	9.5	0.1	3.3	7.2	-
1989	4.8	9.5	0.1	5.1	10.2	-	4.6	9.7	-
1990	4.6	9.2	0.2	5.0	10.0	0.2	3.6	7.5	-
1991	4.7	9.4	0.2	5.0	10.0	0.2	4.6	9.5	-
1992	4.4	8.7	0.2	4.6	9.0	0.2	5.0	10.5	-
1993	3.9	7.6	0.2	4.0	8.0	0.1	4.3	8.6	-
1994	3.8	7.5	0.2	3.9	7.6	0.1	4.1	8.4	-
1995	3.3	6.5	0.2	3.3	6.4	0.2	4.9	10.2	-
1996	2.2	4.4	0.2	2.0	4.0	-	4.6	9.1	-
1997	1.1	2.1	0.1	0.9	1.8	-	2.7	5.3	-
1998	0.8	1.6	-	0.7	1.3	-	1.8	3.8	-
1999	0.8	1.5	0.1	0.7	1.2	-	1.9	3.7	-
2000	0.8	1.5	-	0.7	1.3	-	2.0	4.0	-
2001	0.8	1.5	0.2	0.7	1.2	0.2	1.9	3.7	-
2002	0.7	1.3	0.1	0.5	1.0	-	1.5	3.1	-
2003	0.6	1.2	-	0.6	1.0	-	1.5	3.1	-
2004	0.7	1.3	-	0.6	1.1	-	1.6	3.3	-
2005	0.7	1.4	0.1	0.6	1.1	-	1.9	3.7	-
1975-2005	1.9	3.8	0.1	2.0	3.9	0.1	2.3	4.7	0.3

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.