


Figure 1.21

SEER Observed Incidence and Delay Adjusted Incidence Rates^a All Cancer Sites, By Sex



Source: SEER 9 areas.


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 1.22

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Both Sexes



Source: SEER 9 areas.


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 1.23

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Males



Source: SEER 9 areas.


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 1.24

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Females



Source: SEER 9 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 2.1
All Cancer Sites (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-89	1.2*	1989-92	2.8	1992-95	-2.4	1995-98	1.1	1998-09	-0.3*	2009-15	-1.2*	-0.9*	-1.2*
White	1975-82	0.8*	1982-92	1.8*	1992-95	-2.1	1995-98	1.3	1998-08	-0.2	2008-15	-1.0*	-0.8*	-1.0*
Black	1975-84	1.6*	1984-89	0.4	1989-92	4.8*	1992-95	-1.7	1995-09	-0.4*	2009-15	-1.9*	-1.4*	-1.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-94	-3.1*	1994-99	0.4	1999-05	-0.7*	2005-08	0.4	2008-15	-1.4*			-1.0*	-1.4*
White	1992-94	-3.3*	1994-98	0.6	1998-09	-0.3*	2009-13	-1.8*	2013-15	0.3			-0.8*	-0.8
Black	1992-09	-0.5*	2009-15	-2.0*									-1.5*	-2.0*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-89	1.2*	1989-92	2.8	1992-95	-2.4	1995-98	1.1	1998-09	-0.3*	2009-15	-1.5*	-1.1*	-1.5*
White	1975-82	0.8*	1982-92	1.8*	1992-95	-2.1	1995-98	1.3	1998-08	-0.2	2008-15	-1.3*	-1.1*	-1.3*
Black	1975-84	1.6*	1984-89	0.4	1989-92	4.8*	1992-95	-1.6	1995-09	-0.5*	2009-15	-2.4*	-1.8*	-2.4*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-94	-3.1*	1994-99	0.4	1999-05	-0.7*	2005-08	0.3	2008-15	-1.7*			-1.2*	-1.7*
White	1992-94	-3.1*	1994-01	0.4	2001-04	-1.2	2004-08	0.3	2008-15	-1.6*			-1.2*	-1.6*
White NH ^{ef}	1992-94	-2.9*	1994-01	0.5*	2001-04	-1.4	2004-07	0.9	2007-15	-1.4*			-1.1*	-1.4*
Black	1992-09	-0.6*	2009-15	-2.6*									-1.9*	-2.6*
Black NH ^{ef}	1992-09	-0.5*	2009-15	-2.5*									-1.8*	-2.5*
API ^e	1992-09	-0.4*	2009-15	-1.6*									-1.2*	-1.6*
AI/AN ^{eg}	1992-15	0.0											0.0	0.0
Hispanic ^f	1992-94	-3.5	1994-02	0.4	2002-11	-0.4*	2011-15	-2.2*					-1.2*	-2.2*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-84	0.5*	1984-91	0.3*	1991-94	-0.5	1994-98	-1.3*	1998-01	-0.8*	2001-15	-1.5*	-1.5*	-1.5*
White	1975-90	0.4*	1990-94	-0.3	1994-98	-1.2*	1998-01	-0.6	2001-15	-1.4*			-1.4*	-1.4*
Black	1975-83	1.3*	1983-91	0.7*	1991-95	-0.8*	1995-01	-1.6*	2001-15	-2.1*			-2.1*	-2.1*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	-1.0*	2001-15	-1.5*									-1.5*	-1.5*
White	1992-94	-0.3	1994-98	-1.2*	1998-01	-0.6	2001-15	-1.4*					-1.4*	-1.4*
White NH ^{ef}	1992-94	0.2	1994-98	-1.2*	1998-01	-0.5	2001-15	-1.4*					-1.4*	-1.4*
Black	1992-99	-1.3*	1999-15	-2.1*									-2.1*	-2.1*
Black NH ^{ef}	1992-94	0.0	1994-01	-1.6*	2001-15	-2.0*							-2.0*	-2.0*
API ^e	1992-15	-1.3*											-1.3*	-1.3*
AI/AN ^{eg}	1992-04	0.4	2004-15	-1.1*									-1.1*	-1.1*
Hispanic ^f	1992-98	0.2	1998-15	-1.2*									-1.2*	-1.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 2.2
All Cancer Sites (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-89	1.3*	1989-92	5.2*	1992-95	-4.9*	1995-99	0.6	1999-08	-0.6*	2008-15	-2.3*	-1.9*	-2.3*
White	1975-89	1.4*	1989-92	5.0*	1992-95	-5.1*	1995-00	0.6	2000-08	-0.6*	2008-15	-2.1*	-1.8*	-2.1*
Black	1975-81	2.8*	1981-89	0.6	1989-92	6.8*	1992-05	-1.5*	2005-08	0.5	2008-15	-3.4*	-2.5*	-3.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-94	-5.8*	1994-08	-0.5*	2008-15	-2.5*							-2.1*	-2.5*
White	1992-94	-6.1*	1994-08	-0.3*	2008-15	-2.4*							-2.0*	-2.4*
Black	1992-09	-1.3*	2009-15	-3.5*									-2.8*	-3.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-89	1.3*	1989-92	5.2*	1992-95	-4.9*	1995-99	0.6	1999-08	-0.6*	2008-15	-2.5*	-2.1*	-2.5*
White	1975-89	1.4*	1989-92	5.0*	1992-95	-5.1*	1995-00	0.6	2000-08	-0.6*	2008-15	-2.4*	-2.0*	-2.4*
Black	1975-81	2.8*	1981-89	0.6	1989-92	7.6*	1992-96	-2.5*	1996-09	-1.1*	2009-15	-3.9*	-3.0*	-3.9*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-94	-5.8*	1994-08	-0.5*	2008-15	-2.8*							-2.3*	-2.8*
White	1992-94	-6.1*	1994-08	-0.4*	2008-15	-2.8*							-2.2*	-2.8*
White NH ^{ef}	1992-94	-6.4*	1994-07	-0.2	2007-15	-2.3*							-2.1*	-2.3*
Black	1992-09	-1.3*	2009-15	-4.1*									-3.2*	-4.1*
Black NH ^{ef}	1992-09	-1.3*	2009-15	-4.0*									-3.1*	-4.0*
API ^e	1992-07	-0.7*	2007-15	-2.7*									-2.5*	-2.7*
AI/AN ^{eg}	1992-15	-0.5*											-0.5*	-0.5*
Hispanic ^f	1992-09	-0.5*	2009-15	-3.7*									-2.6*	-3.7*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-79	1.0*	1979-90	0.3*	1990-93	-0.5	1993-01	-1.5*	2001-15	-1.8*			-1.8*	-1.8*
White	1975-79	0.9*	1979-91	0.2*	1991-94	-0.8	1994-97	-1.8*	1997-00	-1.0*	2000-15	-1.7*	-1.7*	-1.7*
Black	1975-82	1.8*	1982-90	1.0*	1990-93	-0.2	1993-01	-2.0*	2001-15	-2.7*			-2.7*	-2.7*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	-1.4*	2001-15	-1.8*									-1.8*	-1.8*
White	1992-01	-1.3*	2001-15	-1.7*									-1.7*	-1.7*
White NH ^{ef}	1992-94	-0.2	1994-97	-1.8*	1997-00	-0.9	2000-15	-1.6*					-1.6*	-1.6*
Black	1992-94	-0.5	1994-01	-2.1*	2001-15	-2.7*							-2.7*	-2.7*
Black NH ^{ef}	1992-94	-0.1	1994-02	-2.1*	2002-15	-2.6*							-2.6*	-2.6*
API ^e	1992-15	-1.6*											-1.6*	-1.6*
AI/AN ^{eg}	1992-15	-0.4*											-0.4*	-0.4*
Hispanic ^f	1992-98	0.0	1998-15	-1.6*									-1.6*	-1.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 2.3
All Cancer Sites (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2006-15	2011-15								
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-79	-0.3	1979-87	1.6*	1987-95	0.1	1995-98	1.5	1998-03	-0.6	2003-15	0.1	0.1	0.1
White	1975-79	-0.2	1979-87	1.7*	1987-95	0.1	1995-98	1.7	1998-04	-0.4	2004-15	0.1	0.1	0.1
Black	1975-91	1.2*	1991-15	0.1									0.1	0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-98	0.8*	1998-03	-0.6	2003-15	0.0							0.0	0.0
White	1992-98	1.0*	1998-03	-0.6	2003-15	0.0							0.0	0.0
Black	1992-15	0.0											0.0	0.0
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-79	-0.3	1979-87	1.6*	1987-95	0.1	1995-98	1.1	1998-15	-0.2*			-0.2*	-0.2*
White	1975-79	-0.2	1979-87	1.7*	1987-95	0.1	1995-98	1.4	1998-15	-0.2*			-0.2*	-0.2*
Black	1975-91	1.1*	1991-12	0.1	2012-15	-2.0*							-0.6	-1.5*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-94	-0.3	1994-98	1.2	1998-04	-0.7*	2004-08	0.4	2008-15	-0.5*			-0.3	-0.5*
White	1992-94	-0.3	1994-98	1.4*	1998-04	-0.7*	2004-08	0.3	2008-15	-0.4*			-0.3	-0.4*
White NH ^{ef}	1992-98	1.1*	1998-15	-0.2*									-0.2*	-0.2*
Black	1992-11	0.1	2011-15	-1.8*									-0.7*	-1.8*
Black NH ^{ef}	1992-11	0.1	2011-15	-1.6*									-0.7*	-1.6*
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-15	0.5*											0.5*	0.5*
Hispanic ^f	1992-15	0.2*											0.2*	0.2*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-90	0.6*	1990-94	-0.2	1994-02	-0.8*	2002-15	-1.4*					-1.4*	-1.4*
White	1975-90	0.6*	1990-95	-0.2	1995-98	-1.2*	1998-01	-0.2	2001-15	-1.3*			-1.3*	-1.3*
Black	1975-91	1.0*	1991-99	-0.6*	1999-15	-1.6*							-1.6*	-1.6*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-95	-0.2	1995-98	-1.2*	1998-01	-0.4	2001-15	-1.4*					-1.4*	-1.4*
White	1992-95	-0.2	1995-98	-1.2*	1998-01	-0.2	2001-15	-1.3*					-1.3*	-1.3*
White NH ^{ef}	1992-94	0.6	1994-98	-1.0*	1998-01	-0.2	2001-15	-1.3*					-1.3*	-1.3*
Black	1992-01	-0.8*	2001-15	-1.6*									-1.6*	-1.6*
Black NH ^{ef}	1992-99	-0.5*	1999-15	-1.5*									-1.5*	-1.5*
API ^e	1992-15	-1.0*											-1.0*	-1.0*
AI/AN ^{eg}	1992-04	0.6	2004-15	-1.4*									-1.4*	-1.4*
Hispanic ^f	1992-99	0.1	1999-15	-1.0*									-1.0*	-1.0*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 2.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a All Cancer Sites, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 2.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a All Cancer Sites, Under 65 Years of Age, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 2.3

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a All Cancer Sites, Ages 65 and Over, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 2.5

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas All Cancer Sites, Males, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 2.6

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas All Cancer Sites, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 3.1
Cancer of the Brain and Other Nervous System (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-87	1.4*	1987-15	-0.1*									-0.1*	-0.1*
White	1975-87	1.6*	1987-15	0.0									0.0	0.0
Black	1975-15	0.2											0.2	0.2
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.2*											-0.2*	-0.2*
White	1992-15	0.0											0.0	0.0
Black	1992-15	0.0											0.0	0.0
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-87	1.5*	1987-15	-0.2*									-0.2*	-0.2*
White	1975-87	1.6*	1987-15	-0.1									-0.1	-0.1
Black	1975-15	0.1											0.1	0.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.2*											-0.2*	-0.2*
White	1992-15	-0.1											-0.1	-0.1
White NH ^{ef}	1992-15	0.1											0.1	0.1
Black	1992-15	-0.2											-0.2	-0.2
Black NH ^{ef}	1992-15	-0.1											-0.1	-0.1
API ^e	1992-15	0.0											0.0	0.0
AI/AN ^{eg}	1992-05	6.1*	2005-15	-4.4									-4.4	-4.4
Hispanic ^f	1992-02	1.1	2002-15	-1.0*									-1.0*	-1.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-83	0.4	1983-91	1.3*	1991-07	-1.0*	2007-15	0.6*					0.4*	0.6*
White	1975-92	1.0*	1992-07	-0.8*	2007-15	0.7*							0.5*	0.7*
Black	1975-92	0.9*	1992-07	-1.1*	2007-15	1.1							0.9	1.1
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-07	-1.0*	2007-15	0.6*									0.4*	0.6*
White	1992-07	-0.8*	2007-15	0.7*									0.5*	0.7*
White NH ^{ef}	1992-07	-0.7*	2007-15	0.8*									0.6*	0.8*
Black	1992-07	-1.1*	2007-15	1.1									0.9	1.1
Black NH ^{ef}	1992-07	-1.1*	2007-15	1.3*									1.0*	1.3*
API ^e	1992-15	0.6*											0.6*	0.6*
AI/AN ^{eg}	1992-15	1.2*											1.2*	1.2*
Hispanic ^f	1992-15	0.2											0.2	0.2

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 3.2
Cancer of the Brain and Other Nervous System (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-89	1.1*	1989-15	-0.3*									-0.3*	-0.3*
White	1975-89	1.3*	1989-15	-0.1									-0.1	-0.1
Black	1975-15	-0.1											-0.1	-0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.2*											-0.2*	-0.2*
White	1992-15	-0.1											-0.1	-0.1
Black	1992-95	9.9	1995-09	-1.2	2009-12	9.8	2012-15	-10.6*					-1.0	-5.9
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-89	1.2*	1989-15	-0.3*									-0.3*	-0.3*
White	1975-89	1.3*	1989-15	-0.2*									-0.2*	-0.2*
Black	1975-15	-0.2											-0.2	-0.2
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.3*											-0.3*	-0.3*
White	1992-15	-0.2*											-0.2*	-0.2*
White NH ^{ef}	1992-15	-0.1											-0.1	-0.1
Black	1992-95	10.0	1995-09	-1.3	2009-12	9.3	2012-15	-11.4*					-1.5	-6.6
Black NH ^{ef}	1992-15	0.0											0.0	0.0
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-15	0.2											0.2	0.2
Hispanic ^f	1992-15	-0.1											-0.1	-0.1
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-77	4.4	1977-82	-0.4	1982-91	1.3*	1991-07	-1.0*	2007-15	0.5*			0.4*	0.5*
White	1975-77	5.0	1977-81	-0.7	1981-91	1.3*	1991-07	-0.8*	2007-15	0.6*			0.5*	0.6*
Black	1975-88	0.7	1988-15	-0.5*									-0.5*	-0.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-07	-0.9*	2007-15	0.5*									0.4*	0.5*
White	1992-07	-0.8*	2007-15	0.6*									0.5*	0.6*
White NH ^{ef}	1992-07	-0.7*	2007-15	0.7*									0.6*	0.7*
Black	1992-15	-0.4*											-0.4*	-0.4*
Black NH ^{ef}	1992-15	-0.3											-0.3	-0.3
API ^e	1992-15	0.5											0.5	0.5
AI/AN ^{eg}	1992-15	1.3											1.3	1.3
Hispanic ^f	1992-15	0.1											0.1	0.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 3.3
Cancer of the Brain and Other Nervous System (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-87	1.6*	1987-15	-0.1									-0.1	-0.1
White	1975-87	1.7*	1987-15	0.0									0.0	0.0
Black	1975-15	0.4											0.4	0.4
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.1											-0.1	-0.1
White	1992-15	0.0											0.0	0.0
Black	1992-15	-0.2											-0.2	-0.2
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-87	1.6*	1987-15	-0.1									-0.1	-0.1
White	1975-87	1.8*	1987-15	0.0									0.0	0.0
Black	1975-15	0.4											0.4	0.4
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.2*											-0.2*	-0.2*
White	1992-15	-0.1											-0.1	-0.1
White NH ^{ef}	1992-15	0.2											0.2	0.2
Black	1992-15	-0.3											-0.3	-0.3
Black NH ^{ef}	1992-15	-0.3											-0.3	-0.3
API ^e	1992-15	-0.1											-0.1	-0.1
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-0.4											-0.4	-0.4
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-92	1.0*	1992-07	-1.1*	2007-15	0.6*							0.4	0.6*
White	1975-92	1.0*	1992-07	-0.9*	2007-15	0.6							0.4	0.6
Black	1975-90	1.3*	1990-15	-0.5*									-0.5*	-0.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-07	-1.1*	2007-15	0.6*									0.4*	0.6*
White	1992-06	-1.0*	2006-15	0.5*									0.5*	0.5*
White NH ^{ef}	1992-07	-0.8*	2007-15	0.7*									0.5*	0.7*
Black	1992-15	-0.5*											-0.5*	-0.5*
Black NH ^{ef}	1992-15	-0.4											-0.4	-0.4
API ^e	1992-08	-1.3	2008-15	4.8*									3.4*	4.8*
AI/AN ^{eg}	1992-15	1.3											1.3	1.3
Hispanic ^f	1992-15	0.2											0.2	0.2

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 3.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Brain and Other Nervous System, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 3.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Brain and Other Nervous System, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 3.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Brain and Other Nervous System, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 4.1
Cancer of the Female Breast (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2006-15	2011-15								
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-80	-0.5	1980-87	4.0*	1987-94	-0.2	1994-99	1.8*	1999-04	-2.3*	2004-15	0.4*	0.4*	0.4*
White	1975-80	-0.5	1980-87	4.1*	1987-94	-0.2	1994-99	2.0*	1999-04	-2.4*	2004-15	0.3	0.3	0.3
Black	1975-79	-1.5	1979-86	4.4*	1986-15	0.5*							0.5*	0.5*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-99	1.3*	1999-04	-2.2*	2004-15	0.3							0.3	0.3
White	1992-99	1.5*	1999-04	-2.4*	2004-15	0.1							0.1	0.1
Black	1992-15	0.4*											0.4*	0.4*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-80	-0.5	1980-87	4.0*	1987-94	-0.2	1994-99	1.8*	1999-04	-2.3*	2004-15	0.3*	0.3*	0.3*
White	1975-80	-0.5	1980-87	4.1*	1987-94	-0.2	1994-99	2.0*	1999-04	-2.4*	2004-15	0.2	0.2	0.2
Black	1975-91	2.3*	1991-15	0.3*									0.3*	0.3*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-99	1.3*	1999-04	-2.2*	2004-15	0.2							0.2	0.2
White	1992-99	1.5*	1999-04	-2.4*	2004-15	0.1							0.1	0.1
White NH ^{ef}	1992-99	1.7*	1999-04	-2.3*	2004-15	0.2							0.2	0.2
Black	1992-15	0.2*											0.2*	0.2*
Black NH ^{ef}	1992-15	0.3*											0.3*	0.3*
API ^e	1992-94	-4.6	1994-97	6.8	1997-05	-0.6	2005-15	1.2*					1.2*	1.2*
AI/AN ^{eg}	1992-15	0.2											0.2	0.2
Hispanic ^f	1992-15	0.3*											0.3*	0.3*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-90	0.4*	1990-95	-1.8*	1995-98	-3.4*	1998-15	-1.8*					-1.8*	-1.8*
White	1975-90	0.3*	1990-95	-2.0*	1995-98	-3.6*	1998-15	-1.8*					-1.8*	-1.8*
Black	1975-93	1.4*	1993-15	-1.5*									-1.5*	-1.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-95	-1.2*	1995-98	-3.7*	1998-03	-1.6*	2003-07	-2.3*	2007-15	-1.6*			-1.7*	-1.6*
White	1992-95	-1.5*	1995-98	-3.6*	1998-10	-1.9*	2010-15	-1.5*					-1.7*	-1.5*
White NH ^{ef}	1992-95	-1.2*	1995-98	-3.7*	1998-15	-1.8*							-1.8*	-1.8*
Black	1992-95	0.4	1995-15	-1.5*									-1.5*	-1.5*
Black NH ^{ef}	1992-95	0.6	1995-15	-1.5*									-1.5*	-1.5*
API ^e	1992-15	-0.8*											-0.8*	-0.8*
AI/AN ^{eg}	1992-15	-0.8*											-0.8*	-0.8*
Hispanic ^f	1992-96	1.3	1996-00	-3.7	2000-15	-1.1*							-1.1*	-1.1*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 4.2
Cancer of the Female Breast (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages <50 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-80	-1.3	1980-87	2.7*	1987-94	-0.8	1994-15	0.3*					0.3*	0.3*
White	1975-80	-1.1	1980-87	2.7*	1987-94	-0.9*	1994-15	0.3*					0.3*	0.3*
Black	1975-90	1.3*	1990-15	-0.1									-0.1	-0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.3*											0.3*	0.3*
White	1992-15	0.2*											0.2*	0.2*
Black	1992-15	0.0											0.0	0.0
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-80	-1.3	1980-87	2.7*	1987-94	-0.8	1994-15	0.3*					0.3*	0.3*
White	1975-80	-1.1	1980-87	2.7*	1987-94	-0.8	1994-15	0.2*					0.2*	0.2*
Black	1975-91	1.2*	1991-15	-0.2									-0.2	-0.2
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	0.2*											0.2*	0.2*
White	1992-15	0.2*											0.2*	0.2*
White NH ^{ef}	1992-15	0.4*											0.4*	0.4*
Black	1992-15	-0.1											-0.1	-0.1
Black NH ^{ef}	1992-02	-1.3*	2002-05	4.0	2005-15	-0.5							-0.5	-0.5
API ^e	1992-15	0.9*											0.9*	0.9*
AI/AN ^{eg}	1992-15	-0.3											-0.3	-0.3
Hispanic ^f	1992-15	0.3*											0.3*	0.3*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-90	-0.4*	1990-07	-3.2*	2007-15	-1.4*							-1.6*	-1.4*
White	1975-90	-0.6*	1990-04	-3.7*	2004-15	-1.7*							-1.7*	-1.7*
Black	1975-87	1.8*	1987-95	-1.2*	1995-15	-2.4*							-2.4*	-2.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-07	-3.1*	2007-15	-1.4*									-1.6*	-1.4*
White	1992-04	-3.7*	2004-15	-1.7*									-1.7*	-1.7*
White NH ^{ef}	1992-04	-3.6*	2004-15	-1.7*									-1.7*	-1.7*
Black	1992-15	-2.3*											-2.3*	-2.3*
Black NH ^{ef}	1992-15	-2.2*											-2.2*	-2.2*
API ^e	1992-01	-0.8	2001-08	-5.1*	2008-15	1.3							-0.2	1.3
AI/AN ^{eg}	1992-15	-2.0*											-2.0*	-2.0*
Hispanic ^f	1992-07	-3.1*	2007-15	-0.6									-0.9	-0.6

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 4.3
Cancer of the Female Breast (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages 50+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2006-15	2011-15								
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-82	0.8	1982-87	5.4*	1987-93	-0.3	1993-99	1.9*	1999-05	-2.5*	2005-15	0.5*	0.5*	0.5*
White	1975-82	0.9*	1982-87	5.5*	1987-93	-0.3	1993-00	1.7*	2000-04	-3.9*	2004-15	0.3	0.3	0.3
Black	1975-88	3.2*	1988-15	0.6*									0.6*	0.6*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-99	1.7*	1999-04	-2.9*	2004-15	0.2							0.2	0.2
White	1992-00	1.4*	2000-04	-3.7*	2004-15	0.1							0.1	0.1
Black	1992-15	0.5*											0.5*	0.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-82	0.8	1982-87	5.4*	1987-93	-0.3	1993-99	1.9*	1999-05	-2.5*	2005-15	0.4*	0.4*	0.4*
White	1975-82	0.9*	1982-87	5.5*	1987-93	-0.3	1993-00	1.7*	2000-04	-3.8*	2004-15	0.2	0.2	0.2
Black	1975-92	2.6*	1992-15	0.4*									0.4*	0.4*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-99	1.7*	1999-04	-2.8*	2004-15	0.1							0.1	0.1
White	1992-00	1.4*	2000-04	-3.7*	2004-15	0.0							0.0	0.0
White NH ^{ef}	1992-01	1.3*	2001-04	-5.0*	2004-15	0.2							0.2	0.2
Black	1992-15	0.4*											0.4*	0.4*
Black NH ^{ef}	1992-15	0.4*											0.4*	0.4*
API ^e	1992-99	2.9*	1999-03	-2.5	2003-15	1.1*							1.1*	1.1*
AI/AN ^{eg}	1992-15	0.4											0.4	0.4
Hispanic ^f	1992-15	0.2*											0.2*	0.2*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-90	0.5*	1990-95	-1.6*	1995-98	-3.0*	1998-15	-1.7*					-1.7*	-1.7*
White	1975-89	0.5*	1989-95	-1.4*	1995-98	-3.3*	1998-15	-1.8*					-1.8*	-1.8*
Black	1975-93	1.6*	1993-15	-1.3*									-1.3*	-1.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-95	-1.2*	1995-98	-3.1*	1998-15	-1.7*							-1.7*	-1.7*
White	1992-94	-0.6	1994-97	-3.4*	1997-15	-1.8*							-1.8*	-1.8*
White NH ^{ef}	1992-94	-0.1	1994-97	-3.3*	1997-15	-1.8*							-1.8*	-1.8*
Black	1992-95	0.8	1995-15	-1.3*									-1.3*	-1.3*
Black NH ^{ef}	1992-94	2.4	1994-15	-1.2*									-1.2*	-1.2*
API ^e	1992-15	-0.5*											-0.5*	-0.5*
AI/AN ^{eg}	1992-15	-0.6											-0.6	-0.6
Hispanic ^f	1992-15	-1.2*											-1.2*	-1.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 4.4
Cancer of the Female Breast (*In Situ*)

Trends in SEER Incidence^{ab} Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
SEER 9 Observed Incidence^a, 1975-2015														
All Races	1975-81	-3.1	1981-87	25.2*	1987-94	3.8*	1994-98	9.8*	1998-09	1.0*	2009-15	-0.9	-0.3	-0.9
White	1975-81	-2.9	1981-87	25.4*	1987-95	3.9*	1995-98	11.1	1998-09	0.9*	2009-15	-1.6*	-0.8	-1.6*
Black	1975-83	6.2	1983-87	23.6*	1987-98	8.3*	1998-15	1.4*					1.4*	1.4*
SEER 13 Observed Incidence^b, 1992-2015														
All Races	1992-94	1.7	1994-99	8.8*	1999-05	-0.6	2005-08	3.3	2008-15	-0.9			0.0	-0.9
White	1992-94	-0.3	1994-98	10.2*	1998-09	0.7*	2009-15	-1.5*					-0.8	-1.5*
White NH ^{ef}	1992-94	0.8	1994-99	9.2*	1999-05	-0.8	2005-08	3.6	2008-15	-1.8*			-0.6	-1.8*
Black	1992-98	9.4*	1998-15	1.5*									1.5*	1.5*
Black NH ^{ef}	1992-98	9.2*	1998-15	1.6*									1.6*	1.6*
API ^e	1992-99	9.7*	1999-15	1.7*									1.7*	1.7*
AI/AN ^{eg}	1992-15	1.5*											1.5*	1.5*
Hispanic ^f	1992-98	8.0*	1998-15	1.8*									1.8*	1.8*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.


The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Figure 4.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Female Breast, by Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 4.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Female Breast, by Age and Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 4.3

SEER Observed Incidence Rates^a Cancer of the Female Breast (*In Situ*), by Race



Source: SEER 9 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).


Figure 4.8

Cancer of the Breast

Delay-Adjusted SEER Incidence & US Mortality

White Females vs Black Females


1975-2015



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

Figure 4.10

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Female Breast, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 5.1
Cancer of the Cervix Uteri (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-82	-4.6*	1982-90	-0.1	1990-07	-2.6*	2007-15	0.1					-0.2	0.1
White	1975-82	-4.8*	1982-90	0.4	1990-06	-2.5*	2006-15	0.5					0.5	0.5
Black	1975-15	-3.6*											-3.6*	-3.6*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-06	-2.7*	2006-15	-1.3*									-1.3*	-1.3*
White	1992-13	-2.0*	2013-15	3.7									-0.8	0.8
Black	1992-15	-3.5*											-3.5*	-3.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-82	-4.6*	1982-90	-0.1	1990-07	-2.6*	2007-15	-0.1					-0.4	-0.1
White	1975-82	-4.8*	1982-90	0.5	1990-05	-2.5*	2005-15	0.1					0.1	0.1
Black	1975-15	-3.7*											-3.7*	-3.7*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-06	-2.7*	2006-15	-1.5*									-1.5*	-1.5*
White	1992-15	-1.9*											-1.9*	-1.9*
White NH ^{ef}	1992-04	-2.4*	2004-15	-0.5									-0.5	-0.5
Black	1992-15	-3.6*											-3.6*	-3.6*
Black NH ^{ef}	1992-15	-3.5*											-3.5*	-3.5*
API ^e	1992-15	-3.7*											-3.7*	-3.7*
AI/AN ^{eg}	1992-15	-0.8											-0.8	-0.8
Hispanic ^f	1992-15	-4.0*											-4.0*	-4.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-82	-4.4*	1982-96	-1.6*	1996-03	-3.8*	2003-15	-0.7*					-0.7*	-0.7*
White	1975-83	-4.2*	1983-96	-1.3*	1996-04	-3.3*	2004-15	-0.2					-0.2	-0.2
Black	1975-81	-4.8*	1981-93	-2.4*	1993-02	-4.8*	2002-15	-2.4*					-2.4*	-2.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-04	-3.2*	2004-15	-0.7*									-0.7*	-0.7*
White	1992-96	-1.4	1996-04	-3.3*	2004-15	-0.2							-0.2	-0.2
White NH ^{ef}	1992-96	-1.0	1996-04	-3.5*	2004-15	0.0							0.0	0.0
Black	1992-01	-5.2*	2001-15	-2.5*									-2.5*	-2.5*
Black NH ^{ef}	1992-01	-5.1*	2001-15	-2.4*									-2.4*	-2.4*
API ^e	1992-15	-3.3*											-3.3*	-3.3*
AI/AN ^{eg}	1992-15	-2.6*											-2.6*	-2.6*
Hispanic ^f	1992-15	-2.6*											-2.6*	-2.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 5.2
Cancer of the Cervix Uteri (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages <50 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-81	-3.7*	1981-98	-0.5	1998-01	-7.5	2001-15	-0.2					-0.2	-0.2
White	1975-82	-4.0*	1982-89	2.1	1989-06	-1.9*	2006-15	1.0					1.0	1.0
Black	1975-15	-3.2*											-3.2*	-3.2*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	-2.4*	2004-15	-1.0*									-1.0*	-1.0*
White	1992-05	-1.9*	2005-15	-0.7									-0.7	-0.7
Black	1992-15	-3.2*											-3.2*	-3.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-81	-3.7*	1981-98	-0.5	1998-01	-7.4	2001-15	-0.2					-0.2	-0.2
White	1975-82	-4.0*	1982-89	2.1	1989-06	-1.9*	2006-15	0.8					0.8	0.8
Black	1975-15	-3.2*											-3.2*	-3.2*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-04	-2.4*	2004-15	-1.1*									-1.1*	-1.1*
White	1992-04	-1.9*	2004-15	-0.9*									-0.9*	-0.9*
White NH ^{ef}	1992-98	0.1	1998-01	-5.6	2001-15	-0.2							-0.2	-0.2
Black	1992-15	-3.2*											-3.2*	-3.2*
Black NH ^{ef}	1992-15	-3.0*											-3.0*	-3.0*
API ^e	1992-15	-2.5*											-2.5*	-2.5*
AI/AN ^{eg}	1992-15	0.6											0.6	0.6
Hispanic ^f	1992-15	-3.5*											-3.5*	-3.5*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-81	-3.8*	1981-96	-0.7*	1996-03	-3.9*	2003-15	-0.3					-0.3	-0.3
White	1975-80	-4.0*	1980-97	-0.3	1997-03	-4.0*	2003-15	0.3					0.3	0.3
Black	1975-15	-3.2*											-3.2*	-3.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-04	-3.0*	2004-15	-0.3									-0.3	-0.3
White	1992-97	-0.8	1997-03	-3.5*	2003-10	-0.5	2010-15	1.8*					0.7	1.8*
White NH ^{ef}	1992-97	0.0	1997-03	-4.1*	2003-15	0.5							0.5	0.5
Black	1992-00	-5.6*	2000-15	-2.3*									-2.3*	-2.3*
Black NH ^{ef}	1992-00	-5.6*	2000-15	-2.1*									-2.1*	-2.1*
API ^e	1992-15	-3.0*											-3.0*	-3.0*
AI/AN ^{eg}	1992-15	-1.5											-1.5	-1.5
Hispanic ^f	1992-11	-3.1*	2011-15	2.4									-0.7	2.4

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 5.3
Cancer of the Cervix Uteri (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages 50+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-83	-4.9*	1983-91	-0.9	1991-06	-3.3*	2006-15	-1.1*					-1.1*	-1.1*
White	1975-85	-4.7*	1985-92	0.3	1992-95	-6.9	1995-13	-2.3*	2013-15	9.0			0.1	3.2
Black	1975-15	-3.9*											-3.9*	-3.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-13	-3.0*	2013-15	3.2									-1.6	0.1
White	1992-15	-2.5*											-2.5*	-2.5*
Black	1992-15	-3.8*											-3.8*	-3.8*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-83	-4.9*	1983-91	-0.9	1991-06	-3.3*	2006-15	-1.2*					-1.2*	-1.2*
White	1975-85	-4.7*	1985-92	0.3	1992-95	-6.8	1995-13	-2.3*	2013-15	8.1			-0.1	2.8
Black	1975-15	-4.0*											-4.0*	-4.0*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-2.9*											-2.9*	-2.9*
White	1992-15	-2.5*											-2.5*	-2.5*
White NH ^{ef}	1992-13	-2.6*	2013-15	8.2									-0.3	2.7
Black	1992-94	8.8	1994-15	-4.2*									-4.2*	-4.2*
Black NH ^{ef}	1992-94	9.7	1994-15	-4.2*									-4.2*	-4.2*
API ^e	1992-06	-6.1*	2006-15	-2.0									-2.0	-2.0
AI/AN ^{eg}	1992-15	-2.6*											-2.6*	-2.6*
Hispanic ^f	1992-15	-4.5*											-4.5*	-4.5*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-84	-4.3*	1984-94	-1.8*	1994-04	-3.4*	2004-15	-0.9*					-0.9*	-0.9*
White	1975-84	-4.6*	1984-96	-1.8*	1996-03	-3.6*	2003-15	-0.7*					-0.7*	-0.7*
Black	1975-79	-5.5*	1979-93	-2.7*	1993-04	-4.6*	2004-15	-2.2*					-2.2*	-2.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-04	-3.3*	2004-15	-0.9*									-0.9*	-0.9*
White	1992-04	-3.0*	2004-15	-0.6*									-0.6*	-0.6*
White NH ^{ef}	1992-04	-3.1*	2004-15	-0.6*									-0.6*	-0.6*
Black	1992-04	-4.7*	2004-15	-2.1*									-2.1*	-2.1*
Black NH ^{ef}	1992-05	-4.5*	2005-15	-1.9*									-1.9*	-1.9*
API ^e	1992-15	-3.4*											-3.4*	-3.4*
AI/AN ^{eg}	1992-15	-3.4*											-3.4*	-3.4*
Hispanic ^f	1992-15	-2.6*											-2.6*	-2.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 5.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Cervix Uteri, by Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 5.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Cervix Uteri, by Age and Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 5.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Cervix Uteri, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero (p < 0.05).

Table 6.1
Cancer of the Colon and Rectum (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-85	0.8*	1985-95	-1.8*	1995-98	1.4	1998-08	-2.3*	2008-11	-4.5*	2011-15	-1.1	-2.5*	-1.1
White	1975-85	0.8*	1985-95	-2.0*	1995-98	1.6	1998-08	-2.5*	2008-11	-4.8*	2011-15	-0.8	-2.5*	-0.8
Black	1975-80	3.2*	1980-03	-0.1	2003-15	-2.9*							-2.9*	-2.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-95	-2.1*	1995-98	1.6	1998-08	-2.2*	2008-12	-3.9*	2012-15	-0.7			-2.5*	-1.5*
White	1992-95	-2.2*	1995-98	1.7	1998-08	-2.4*	2008-12	-4.1*	2012-15	-0.1			-2.4*	-1.1
Black	1992-04	-0.1	2004-15	-3.2*									-3.2*	-3.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-85	0.8*	1985-95	-1.8*	1995-98	1.5	1998-08	-2.3*	2008-11	-4.5*	2011-15	-1.4*	-2.6*	-1.4*
White	1975-85	0.8*	1985-95	-2.0*	1995-98	1.6	1998-08	-2.5*	2008-11	-4.8*	2011-15	-1.1	-2.7*	-1.1
Black	1975-80	3.2*	1980-03	-0.1	2003-15	-3.1*							-3.1*	-3.1*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-95	-2.1*	1995-98	1.6	1998-08	-2.2*	2008-12	-4.0*	2012-15	-1.1			-2.6*	-1.8*
White	1992-95	-2.2*	1995-98	1.7	1998-08	-2.4*	2008-11	-4.7*	2011-15	-1.3*			-2.7*	-1.3*
White NH ^{ef}	1992-95	-2.1*	1995-98	1.8	1998-08	-2.5*	2008-11	-4.9*	2011-15	-1.0			-2.6*	-1.0
Black	1992-04	0.0	2004-15	-3.5*									-3.5*	-3.5*
Black NH ^{ef}	1992-04	0.0	2004-15	-3.4*									-3.4*	-3.4*
API ^e	1992-02	-0.3	2002-15	-2.6*									-2.6*	-2.6*
AI/AN ^{eg}	1992-15	-0.1											-0.1	-0.1
Hispanic ^f	1992-07	0.0	2007-15	-2.5*									-2.3*	-2.5*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-84	-0.5*	1984-02	-1.8*	2002-05	-3.9*	2005-15	-2.4*					-2.4*	-2.4*
White	1975-78	0.2	1978-85	-1.0*	1985-02	-1.9*	2002-05	-3.8*	2005-12	-2.5*	2012-15	-1.6*	-2.2*	-1.8*
Black	1975-90	0.7*	1990-02	-0.9*	2002-15	-3.1*							-3.1*	-3.1*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-02	-1.8*	2002-05	-3.9*	2005-15	-2.4*							-2.4*	-2.4*
White	1992-02	-1.9*	2002-05	-3.8*	2005-12	-2.5*	2012-15	-1.6*					-2.2*	-1.8*
White NH ^{ef}	1992-02	-1.8*	2002-05	-3.9*	2005-12	-2.5*	2012-15	-1.5*					-2.2*	-1.8*
Black	1992-01	-0.6*	2001-15	-3.0*									-3.0*	-3.0*
Black NH ^{ef}	1992-01	-0.6*	2001-15	-2.9*									-2.9*	-2.9*
API ^e	1992-15	-1.8*											-1.8*	-1.8*
AI/AN ^{eg}	1992-95	12.3	1995-15	-0.5									-0.5	-0.5
Hispanic ^f	1992-01	0.3	2001-15	-1.8*									-1.8*	-1.8*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 6.2
Cancer of the Colon and Rectum (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-85	1.1*	1985-91	-1.2*	1991-95	-3.2*	1995-98	2.1	1998-15	-2.9*			-2.9*	-2.9*
White	1975-86	1.1*	1986-95	-2.3*	1995-98	1.4	1998-07	-2.9*	2007-12	-4.1*	2012-15	-0.5	-2.8*	-1.4
Black	1975-80	4.5	1980-02	0.0	2002-15	-2.7*							-2.7*	-2.7*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-95	-2.6*	1995-98	1.4	1998-08	-2.5*	2008-12	-3.9*	2012-15	-1.1			-2.6*	-1.8*
White	1992-95	-2.8*	1995-98	1.5	1998-08	-2.7*	2008-12	-4.3*	2012-15	-0.4			-2.7*	-1.4*
Black	1992-03	0.0	2003-15	-3.0*									-3.0*	-3.0*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-86	1.1*	1986-95	-2.2*	1995-98	1.5	1998-15	-2.9*					-2.9*	-2.9*
White	1975-86	1.1*	1986-95	-2.3*	1995-98	1.4	1998-08	-2.9*	2008-11	-4.9*	2011-15	-1.5*	-3.0*	-1.5*
Black	1975-80	4.4	1980-02	0.1	2002-15	-2.9*							-2.9*	-2.9*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-95	-2.6*	1995-98	1.4	1998-08	-2.5*	2008-12	-3.9*	2012-15	-1.4*			-2.8*	-2.1*
White	1992-95	-2.8*	1995-98	1.5	1998-08	-2.7*	2008-12	-4.3*	2012-15	-0.8			-2.8*	-1.7*
White NH ^{ef}	1992-95	-2.9*	1995-98	1.6	1998-08	-2.9*	2008-12	-4.5*	2012-15	-0.3			-2.8*	-1.4*
Black	1992-04	-0.2	2004-15	-3.4*									-3.4*	-3.4*
Black NH ^{ef}	1992-04	-0.1	2004-15	-3.3*									-3.3*	-3.3*
API ^e	1992-01	-0.3	2001-15	-2.5*									-2.5*	-2.5*
AI/AN ^{eg}	1992-15	-0.2											-0.2	-0.2
Hispanic ^f	1992-98	1.6	1998-08	-0.9	2008-15	-2.9*							-2.4*	-2.9*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-79	0.6	1979-87	-0.6*	1987-02	-1.9*	2002-05	-4.1*	2005-15	-2.5*			-2.5*	-2.5*
White	1975-79	0.6	1979-87	-0.8*	1987-02	-2.1*	2002-05	-4.3*	2005-15	-2.4*			-2.4*	-2.4*
Black	1975-90	1.2*	1990-02	-0.7*	2002-15	-2.8*							-2.8*	-2.8*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-02	-2.0*	2002-05	-4.0*	2005-15	-2.5*							-2.5*	-2.5*
White	1992-02	-2.1*	2002-05	-4.1*	2005-12	-2.6*	2012-15	-1.8*					-2.3*	-2.0*
White NH ^{ef}	1992-02	-2.1*	2002-05	-4.3*	2005-15	-2.5*							-2.5*	-2.5*
Black	1992-02	-0.7*	2002-15	-2.8*									-2.8*	-2.8*
Black NH ^{ef}	1992-02	-0.7*	2002-15	-2.7*									-2.7*	-2.7*
API ^e	1992-15	-1.9*											-1.9*	-1.9*
AI/AN ^{eg}	1992-15	0.3											0.3	0.3
Hispanic ^f	1992-99	0.5	1999-15	-1.6*									-1.6*	-1.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 6.3
Cancer of the Colon and Rectum (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2006-15	2011-15								
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-85	0.3	1985-95	-1.9*	1995-98	1.8	1998-08	-2.0*	2008-11	-4.6*	2011-15	-0.9	-2.4*	-0.9
White	1975-85	0.3	1985-95	-2.1*	1995-98	2.0	1998-08	-2.2*	2008-11	-4.6*	2011-15	-0.6	-2.3*	-0.6
Black	1975-04	-0.1	2004-15	-3.5*									-3.5*	-3.5*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-95	-1.8	1995-98	1.8	1998-08	-1.9*	2008-12	-4.0*	2012-15	-0.6			-2.4*	-1.4*
White	1992-95	-1.7	1995-98	1.7	1998-08	-2.1*	2008-11	-4.5*	2011-15	-0.8			-2.3*	-0.8
Black	1992-06	-0.3	2006-15	-4.1*									-4.1*	-4.1*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-85	0.3	1985-95	-1.9*	1995-98	1.8	1998-08	-2.0*	2008-11	-4.7*	2011-15	-1.2	-2.6*	-1.2
White	1975-85	0.3	1985-95	-2.1*	1995-98	2.0	1998-08	-2.2*	2008-11	-4.6*	2011-15	-0.9	-2.4*	-0.9
Black	1975-04	0.0	2004-15	-3.7*									-3.7*	-3.7*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-95	-1.8	1995-98	1.8	1998-08	-2.0*	2008-12	-4.1*	2012-15	-0.9			-2.6*	-1.7*
White	1992-95	-1.7	1995-98	1.7	1998-08	-2.1*	2008-11	-4.6	2011-15	-1.1			-2.5*	-1.1
White NH ^{ef}	1992-95	-1.5	1995-98	1.9	1998-08	-2.2*	2008-11	-4.7	2011-15	-0.9			-2.5*	-0.9
Black	1992-06	-0.3	2006-15	-4.4*									-4.4*	-4.4*
Black NH ^{ef}	1992-06	-0.3	2006-15	-4.3*									-4.3*	-4.3*
API ^e	1992-02	0.1	2002-15	-2.7*									-2.7*	-2.7*
AI/AN ^{eg}	1992-15	-0.1											-0.1	-0.1
Hispanic ^f	1992-07	0.1	2007-15	-2.1*									-1.9*	-2.1*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-84	-1.0*	1984-01	-1.8*	2001-12	-2.9*	2012-15	-1.4					-2.4*	-1.8*
White	1975-84	-1.2*	1984-01	-2.0*	2001-11	-2.9*	2011-15	-1.5*					-2.3*	-1.5*
Black	1975-85	0.7*	1985-01	-0.7*	2001-15	-3.3*							-3.3*	-3.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	-1.7*	2001-10	-3.0*	2010-15	-2.0*							-2.5*	-2.0*
White	1992-01	-1.8*	2001-10	-3.0*	2010-15	-1.7*							-2.3*	-1.7*
White NH ^{ef}	1992-00	-1.7*	2000-11	-2.9*	2011-15	-1.4*							-2.2*	-1.4*
Black	1992-01	-0.7*	2001-15	-3.3*									-3.3*	-3.3*
Black NH ^{ef}	1992-01	-0.6*	2001-15	-3.2*									-3.2*	-3.2*
API ^e	1992-15	-1.7*											-1.7*	-1.7*
AI/AN ^{eg}	1992-15	-0.4											-0.4	-0.4
Hispanic ^f	1992-02	0.4	2002-15	-2.3*									-2.3*	-2.3*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 6.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Colon and Rectum, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 6.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Colon and Rectum, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 6.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Colon and Rectum, Female, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 7.1
Cancer of the Corpus and Uterus, NOS (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-79	-6.0*	1979-88	-1.7*	1988-97	0.7*	1997-06	-0.4*	2006-09	3.5*	2009-15	0.3	1.3*	0.3
White	1975-79	-6.1*	1979-88	-1.7*	1988-98	0.7*	1998-03	-1.1	2003-15	0.9*			0.9*	0.9*
Black	1975-93	-0.5	1993-15	2.4*									2.4*	2.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-06	-0.2	2006-09	3.4	2009-15	0.5							1.4	0.5
White	1992-97	0.8	1997-03	-1.0	2003-15	1.1*							1.1*	1.1*
Black	1992-15	2.3*											2.3*	2.3*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-79	-6.0*	1979-88	-1.7*	1988-97	0.7*	1997-06	-0.4*	2006-09	3.5	2009-15	0.2	1.3*	0.2
White	1975-79	-6.1*	1979-88	-1.7*	1988-98	0.7*	1998-02	-1.4	2002-15	0.8*			0.8*	0.8*
Black	1975-93	-0.5	1993-15	2.3*									2.3*	2.3*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-06	-0.2	2006-09	3.3	2009-15	0.4							1.3	0.4
White	1992-97	0.7	1997-03	-1.0	2003-15	1.0*							1.0*	1.0*
White NH ^{ef}	1992-98	0.8	1998-04	-1.4*	2004-09	1.9*	2009-15	-0.2					0.5	-0.2
Black	1992-15	2.2*											2.2*	2.2*
Black NH ^{ef}	1992-15	2.3*											2.3*	2.3*
API ^e	1992-15	1.7*											1.7*	1.7*
AI/AN ^{eg}	1992-15	2.0*											2.0*	2.0*
Hispanic ^f	1992-00	-0.6	2000-04	4.8	2004-08	-0.6	2008-12	6.5*	2012-15	0.1			2.7*	1.6
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-89	-1.6*	1989-97	-0.7*	1997-08	0.3*	2008-15	1.9*					1.6*	1.9*
White	1975-91	-1.7*	1991-07	-0.1	2007-15	1.7*							1.5*	1.7*
Black	1975-95	-0.9*	1995-08	0.4*	2008-15	2.5*							2.1*	2.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	-0.7	1997-08	0.3*	2008-15	1.9*							1.6*	1.9*
White	1992-07	0.0	2007-15	1.7*									1.5*	1.7*
White NH ^{ef}	1992-08	0.1	2008-15	1.7*									1.4*	1.7*
Black	1992-08	0.3	2008-15	2.6*									2.1*	2.6*
Black NH ^{ef}	1992-08	0.4*	2008-15	2.8*									2.2*	2.8*
API ^e	1992-15	1.8*											1.8*	1.8*
AI/AN ^{eg}	1992-15	0.6											0.6	0.6
Hispanic ^f	1992-07	0.1	2007-15	2.8*									2.5*	2.8*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 7.2
Cancer of the Corpus and Uterus, NOS (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages <50 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-88	-2.6*	1988-15	1.3*									1.3*	1.3*
White	1975-88	-2.6*	1988-15	1.1*									1.1*	1.1*
Black	1975-85	-5.1*	1985-15	2.4*									2.4*	2.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-96	-0.8	1996-15	1.8*									1.8*	1.8*
White	1992-15	1.5*											1.5*	1.5*
Black	1992-15	2.2*											2.2*	2.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-88	-2.6*	1988-15	1.3*									1.3*	1.3*
White	1975-88	-2.6*	1988-15	1.1*									1.1*	1.1*
Black	1975-85	-5.1*	1985-15	2.4*									2.4*	2.4*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-96	-0.7	1996-15	1.8*									1.8*	1.8*
White	1992-15	1.5*											1.5*	1.5*
White NH ^{ef}	1992-15	0.9*											0.9*	0.9*
Black	1992-15	2.1*											2.1*	2.1*
Black NH ^{ef}	1992-15	2.3*											2.3*	2.3*
API ^e	1992-15	1.3*											1.3*	1.3*
AI/AN ^{eg}	1992-15	1.3											1.3	1.3
Hispanic ^f	1992-94	-11.0	1994-15	3.4*									3.4*	3.4*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-92	-3.2*	1992-15	1.8*									1.8*	1.8*
White	1975-91	-3.1*	1991-15	1.6*									1.6*	1.6*
Black	1975-95	-4.5*	1995-15	2.8*									2.8*	2.8*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	1.7*											1.7*	1.7*
White	1992-15	1.6*											1.6*	1.6*
White NH ^{ef}	1992-15	1.5*											1.5*	1.5*
Black	1992-15	2.3*											2.3*	2.3*
Black NH ^{ef}	1992-15	2.6*											2.6*	2.6*
API ^e	1992-15	1.0											1.0	1.0
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	1.5*											1.5*	1.5*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 7.3
Cancer of the Corpus and Uterus, NOS (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages 50+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-79	-6.2*	1979-88	-1.7*	1988-97	0.6*	1997-06	-0.7*	2006-09	3.6	2009-15	0.1	1.3	0.1
White	1975-79	-6.2*	1979-88	-1.7*	1988-98	0.7*	1998-03	-1.4	2003-15	0.8*			0.8*	0.8*
Black	1975-97	0.0	1997-15	2.7*									2.7*	2.7*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-97	0.8	1997-05	-1.1*	2005-10	2.6*	2010-15	0.0					1.1*	0.0
White	1992-97	0.7	1997-05	-1.1*	2005-10	2.0*	2010-15	0.0					0.9*	0.0
Black	1992-15	2.3*											2.3*	2.3*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-79	-6.2*	1979-88	-1.7*	1988-97	0.6*	1997-06	-0.7*	2006-09	3.6	2009-15	0.0	1.2	0.0
White	1975-79	-6.2*	1979-88	-1.7*	1988-98	0.7*	1998-03	-1.4	2003-15	0.7*			0.7*	0.7*
Black	1975-97	0.0	1997-15	2.6*									2.6*	2.6*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-97	0.8	1997-05	-1.1*	2005-10	2.5*	2010-15	-0.2					1.0*	-0.2
White	1992-97	0.7	1997-05	-1.1*	2005-10	2.0*	2010-15	-0.2					0.8	-0.2
White NH ^{ef}	1992-97	1.1	1997-04	-1.4*	2004-10	1.6*	2010-15	-0.4					0.5	-0.4
Black	1992-15	2.3*											2.3*	2.3*
Black NH ^{ef}	1992-15	2.3*											2.3*	2.3*
API ^e	1992-95	6.7	1995-06	0.4	2006-09	7.2	2009-15	0.3					2.5	0.3
AI/AN ^{eg}	1992-15	2.2*											2.2*	2.2*
Hispanic ^f	1992-00	-0.6	2000-15	2.7*									2.7*	2.7*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-93	-1.4*	1993-08	0.1	2008-15	2.0*							1.6*	2.0*
White	1975-92	-1.6*	1992-08	-0.1	2008-15	1.9*							1.4*	1.9*
Black	1975-04	-0.4*	2004-15	1.9*									1.9*	1.9*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	-0.7	1997-08	0.2	2008-15	1.9*							1.5*	1.9*
White	1992-07	-0.1	2007-15	1.6*									1.4*	1.6*
White NH ^{ef}	1992-08	0.0	2008-15	1.8*									1.4*	1.8*
Black	1992-08	0.3	2008-15	2.5*									2.0*	2.5*
Black NH ^{ef}	1992-08	0.3	2008-15	2.6*									2.1*	2.6*
API ^e	1992-15	1.9*											1.9*	1.9*
AI/AN ^{eg}	1992-15	0.6											0.6	0.6
Hispanic ^f	1992-07	0.0	2007-15	2.8*									2.5*	2.8*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 7.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Corpus and Uterus, NOS, by Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 7.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Corpus and Uterus, NOS, by Age and Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 7.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Corpus and Uterus, NOS, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 8.1
Cancer of the Esophagus (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-04	0.6*	2004-15	-1.2*									-1.2*	-1.2*
White	1975-04	1.4*	2004-15	-0.5									-0.5	-0.5
Black	1975-86	1.1	1986-15	-4.4*									-4.4*	-4.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.5*											-0.5*	-0.5*
White	1992-04	1.3*	2004-15	-0.7*									-0.7*	-0.7*
Black	1992-15	-4.5*											-4.5*	-4.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-04	0.6*	2004-15	-1.2*									-1.2*	-1.2*
White	1975-04	1.4*	2004-15	-0.6									-0.6	-0.6
Black	1975-86	1.1	1986-15	-4.4*									-4.4*	-4.4*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.6*											-0.6*	-0.6*
White	1992-04	1.3*	2004-15	-0.8*									-0.8*	-0.8*
White NH ^{ef}	1992-04	1.7*	2004-15	-0.8*									-0.8*	-0.8*
Black	1992-15	-4.5*											-4.5*	-4.5*
Black NH ^{ef}	1992-15	-4.5*											-4.5*	-4.5*
API ^e	1992-15	-1.9*											-1.9*	-1.9*
AI/AN ^{eg}	1992-15	-1.2											-1.2	-1.2
Hispanic ^f	1992-15	-0.8*											-0.8*	-0.8*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-02	0.7*	2002-15	-0.9*									-0.9*	-0.9*
White	1975-83	0.4*	1983-00	1.7*	2000-05	0.8*	2005-15	-0.5*					-0.5*	-0.5*
Black	1975-82	1.1*	1982-93	-1.6*	1993-15	-4.6*							-4.6*	-4.6*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-04	0.5*	2004-15	-1.0*									-1.0*	-1.0*
White	1992-04	1.4*	2004-15	-0.5*									-0.5*	-0.5*
White NH ^{ef}	1992-03	1.8*	2003-15	-0.2*									-0.2*	-0.2*
Black	1992-15	-4.5*											-4.5*	-4.5*
Black NH ^{ef}	1992-15	-4.5*											-4.5*	-4.5*
API ^e	1992-15	-1.8*											-1.8*	-1.8*
AI/AN ^{eg}	1992-01	5.0*	2001-15	-0.8									-0.8	-0.8
Hispanic ^f	1992-15	-1.1*											-1.1*	-1.1*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 8.2
Cancer of the Esophagus (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-04	0.8*	2004-15	-1.2*									-1.2*	-1.2*
White	1975-04	1.7*	2004-15	-0.6									-0.6	-0.6
Black	1975-86	1.0	1986-15	-4.7*									-4.7*	-4.7*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	0.4	2004-15	-1.3*									-1.3*	-1.3*
White	1992-04	1.5*	2004-15	-0.8*									-0.8*	-0.8*
Black	1992-15	-4.6*											-4.6*	-4.6*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-04	0.8*	2004-15	-1.2*									-1.2*	-1.2*
White	1975-04	1.7*	2004-15	-0.7									-0.7	-0.7
Black	1975-86	1.0	1986-15	-4.7*									-4.7*	-4.7*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-04	0.4	2004-15	-1.3*									-1.3*	-1.3*
White	1992-04	1.6*	2004-15	-0.9*									-0.9*	-0.9*
White NH ^{ef}	1992-04	1.9*	2004-15	-0.8*									-0.8*	-0.8*
Black	1992-15	-4.7*											-4.7*	-4.7*
Black NH ^{ef}	1992-15	-4.6*											-4.6*	-4.6*
API ^e	1992-15	-2.0*											-2.0*	-2.0*
AI/AN ^{eg}	1992-15	-2.0											-2.0	-2.0
Hispanic ^f	1992-15	-1.0*											-1.0*	-1.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-85	0.7*	1985-94	1.2*	1994-05	0.4*	2005-15	-1.1*					-1.1*	-1.1*
White	1975-82	0.3	1982-99	1.9*	1999-05	1.1*	2005-15	-0.6*					-0.6*	-0.6*
Black	1975-82	1.4*	1982-93	-1.6*	1993-15	-4.8*							-4.8*	-4.8*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-05	0.5*	2005-15	-1.1*									-1.1*	-1.1*
White	1992-05	1.4*	2005-15	-0.7*									-0.7*	-0.7*
White NH ^{ef}	1992-04	1.7*	2004-15	-0.4*									-0.4*	-0.4*
Black	1992-15	-4.7*											-4.7*	-4.7*
Black NH ^{ef}	1992-15	-4.6*											-4.6*	-4.6*
API ^e	1992-15	-1.8*											-1.8*	-1.8*
AI/AN ^{eg}	1992-15	0.5											0.5	0.5
Hispanic ^f	1992-15	-1.1*											-1.1*	-1.1*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 8.3
Cancer of the Esophagus (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-15	-0.6*											-0.6*	-0.6*
White	1975-15	-0.1											-0.1	-0.1
Black	1975-81	8.5*	1981-15	-3.2*									-3.2*	-3.2*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.3*											-1.3*	-1.3*
White	1992-15	-0.6*											-0.6*	-0.6*
Black	1992-15	-4.1*											-4.1*	-4.1*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-97	0.0	1997-15	-1.3*									-1.3*	-1.3*
White	1975-15	-0.1											-0.1	-0.1
Black	1975-81	8.6*	1981-15	-3.3*									-3.3*	-3.3*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.3*											-1.3*	-1.3*
White	1992-15	-0.6*											-0.6*	-0.6*
White NH ^{ef}	1992-15	-0.4*											-0.4*	-0.4*
Black	1992-15	-4.2*											-4.2*	-4.2*
Black NH ^{ef}	1992-15	-4.1*											-4.1*	-4.1*
API ^e	1992-15	-1.2											-1.2	-1.2
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-0.8											-0.8	-0.8
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-01	0.0	2001-15	-1.7*									-1.7*	-1.7*
White	1975-02	0.4*	2002-15	-1.1*									-1.1*	-1.1*
Black	1975-92	-0.2	1992-15	-4.1*									-4.1*	-4.1*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	0.0	2001-15	-1.7*									-1.7*	-1.7*
White	1992-01	0.8*	2001-15	-1.1*									-1.1*	-1.1*
White NH ^{ef}	1992-01	1.0*	2001-15	-0.9*									-0.9*	-0.9*
Black	1992-15	-4.1*											-4.1*	-4.1*
Black NH ^{ef}	1992-15	-4.0*											-4.0*	-4.0*
API ^e	1992-15	-1.9*											-1.9*	-1.9*
AI/AN ^{eg}	1992-15	1.5											1.5	1.5
Hispanic ^f	1992-96	10.0	1996-15	-2.2*									-2.2*	-2.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.


* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Figure 8.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Esophagus, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 8.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Esophagus, Male, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 8.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Esophagus, Female, by Race



^a Source: SEER 9 areas and SEER 13 areas.
Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
* The APC is significantly different from zero ($p < 0.05$).

Table 9.1
Hodgkin LymphomaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-03	-0.2*	2003-06	3.0	2006-15	-1.6*							-1.6*	-1.6*
White	1975-88	0.6*	1988-01	-0.9*	2001-07	1.9	2007-15	-2.1*					-1.7*	-2.1*
Black	1975-15	1.1*											1.1*	1.1*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-07	0.5*	2007-15	-1.6*									-1.4*	-1.6*
White	1992-02	-0.2	2002-07	1.6	2007-15	-2.3*							-1.8*	-2.3*
Black	1992-15	0.8*											0.8*	0.8*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-03	-0.2*	2003-06	3.1	2006-15	-1.8*							-1.8*	-1.8*
White	1975-88	0.6*	1988-01	-0.9*	2001-07	1.9	2007-15	-2.3*					-1.8*	-2.3*
Black	1975-15	1.0*											1.0*	1.0*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-07	0.5*	2007-15	-1.8*									-1.5*	-1.8*
White	1992-07	0.4*	2007-15	-2.0*									-1.7*	-2.0*
White NH ^{ef}	1992-07	0.6*	2007-15	-2.3*									-2.0*	-2.3*
Black	1992-15	0.7*											0.7*	0.7*
Black NH ^{ef}	1992-15	0.8*											0.8*	0.8*
API ^e	1992-15	1.3*											1.3*	1.3*
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	0.0											0.0	0.0
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-79	-6.5*	1979-95	-3.5*	1995-15	-2.8*							-2.8*	-2.8*
White	1975-79	-6.3*	1979-95	-3.6*	1995-15	-2.7*							-2.7*	-2.7*
Black	1975-15	-2.9*											-2.9*	-2.9*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-2.8*											-2.8*	-2.8*
White	1992-94	-7.1	1994-15	-2.6*									-2.6*	-2.6*
White NH ^{ef}	1992-15	-2.8*											-2.8*	-2.8*
Black	1992-15	-2.9*											-2.9*	-2.9*
Black NH ^{ef}	1992-15	-2.8*											-2.8*	-2.8*
API ^e	1992-15	-1.8*											-1.8*	-1.8*
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-2.3*											-2.3*	-2.3*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 9.2
Hodgkin LymphomaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-15	-0.4*											-0.4*	-0.4*
White	1975-15	-0.4*											-0.4*	-0.4*
Black	1975-15	0.7*											0.7*	0.7*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.0											0.0	0.0
White	1992-15	-0.1											-0.1	-0.1
Black	1992-15	1.1*											1.1*	1.1*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	-0.4*											-0.4*	-0.4*
White	1975-15	-0.4*											-0.4*	-0.4*
Black	1975-15	0.6*											0.6*	0.6*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.1											-0.1	-0.1
White	1992-07	0.4	2007-15	-1.7*									-1.5*	-1.7*
White NH ^{ef}	1992-08	0.6*	2008-15	-2.2*									-1.6*	-2.2*
Black	1992-15	1.0*											1.0*	1.0*
Black NH ^{ef}	1992-15	1.1*											1.1*	1.1*
API ^e	1992-15	1.2											1.2	1.2
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-0.5											-0.5	-0.5
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-96	-4.0*	1996-15	-2.5*									-2.5*	-2.5*
White	1975-96	-4.1*	1996-15	-2.4*									-2.4*	-2.4*
Black	1975-15	-3.2*											-3.2*	-3.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	-10.2	1994-15	-2.5*									-2.5*	-2.5*
White	1992-94	-10.5	1994-15	-2.3*									-2.3*	-2.3*
White NH ^{ef}	1992-15	-2.6*											-2.6*	-2.6*
Black	1992-15	-3.3*											-3.3*	-3.3*
Black NH ^{ef}	1992-15	-3.2*											-3.2*	-3.2*
API ^e	1992-15	-2.3*											-2.3*	-2.3*
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-2.4*											-2.4*	-2.4*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).^b Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).^c Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.^d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.^e API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic^f Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^g Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 9.3
Hodgkin LymphomaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-07	0.4*	2007-15	-1.9*									-1.7*	-1.9*
White	1975-07	0.4*	2007-15	-2.1*									-1.8*	-2.1*
Black	1975-15	1.6*											1.6*	1.6*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-08	0.3	2008-15	-2.5*									-1.9*	-2.5*
White	1992-08	0.3	2008-15	-2.6*									-2.0*	-2.6*
Black	1992-15	0.3											0.3	0.3
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-07	0.4*	2007-15	-2.1*									-1.8*	-2.1*
White	1975-07	0.4*	2007-15	-2.2*									-2.0*	-2.2*
Black	1975-15	1.6*											1.6*	1.6*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-08	0.3	2008-15	-2.7*									-2.0*	-2.7*
White	1992-08	0.3	2008-15	-2.8*									-2.1*	-2.8*
White NH ^{ef}	1992-07	0.6	2007-15	-2.9*									-2.5*	-2.9*
Black	1992-15	0.2											0.2	0.2
Black NH ^{ef}	1992-15	0.3											0.3	0.3
API ^e	1992-10	3.3*	2010-15	-8.9									-3.7	-8.9
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	0.5											0.5	0.5
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-84	-4.9*	1984-15	-2.8*									-2.8*	-2.8*
White	1975-85	-5.0*	1985-09	-2.5*	2009-15	-5.3*							-4.3*	-5.3*
Black	1975-15	-2.5*											-2.5*	-2.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-3.0*											-3.0*	-3.0*
White	1992-15	-3.0*											-3.0*	-3.0*
White NH ^{ef}	1992-09	-2.5*	2009-15	-6.2*									-5.0*	-6.2*
Black	1992-15	-2.5*											-2.5*	-2.5*
Black NH ^{ef}	1992-15	-2.4*											-2.4*	-2.4*
API ^e	-	-											-	-
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-2.1*											-2.1*	-2.1*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Figure 9.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Hodgkin Lymphoma, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 9.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Hodgkin Lymphoma, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 9.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Hodgkin Lymphoma, Female, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 11.1
Cancer of the Kidney and Renal Pelvis (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-94	2.3*	1994-97	-0.7	1997-07	3.3*	2007-15	0.5					0.8*	0.5
White	1975-92	2.3*	1992-97	0.3	1997-07	3.2*	2007-15	0.4					0.8*	0.4
Black	1975-09	3.0*	2009-15	0.3									1.2	0.3
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-97	1.0	1997-05	2.7*	2005-08	4.5*	2008-11	-1.5	2011-15	1.4*			1.1	1.4*
White	1992-97	1.0	1997-08	3.1*	2008-15	0.3							0.9*	0.3
Black	1992-09	3.0*	2009-15	0.1									1.1	0.1
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-94	2.3*	1994-97	-0.5	1997-08	3.1*	2008-15	-0.1					0.6*	-0.1
White	1975-92	2.3*	1992-97	0.3	1997-07	3.2*	2007-15	0.1					0.4	0.1
Black	1975-09	3.0*	2009-15	-0.4									0.7	-0.4
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-99	1.4*	1999-08	3.2*	2008-15	-0.1							0.6*	-0.1
White	1992-97	0.9	1997-08	3.1*	2008-15	-0.1							0.6*	-0.1
White NH ^{ef}	1992-97	0.8	1997-08	3.0*	2008-15	-0.2							0.5*	-0.2
Black	1992-09	2.9*	2009-15	-0.5									0.6	-0.5
Black NH ^{ef}	1992-09	3.0*	2009-15	-0.4									0.7	-0.4
API ^e	1992-15	2.4*											2.4*	2.4*
AI/AN ^{eg}	1992-15	0.9											0.9	0.9
Hispanic ^f	1992-07	3.1*	2007-15	0.7									1.0	0.7
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-95	1.0*	1995-15	-0.7*									-0.7*	-0.7*
White	1975-95	0.9*	1995-15	-0.6*									-0.6*	-0.6*
Black	1975-94	2.3*	1994-15	-0.9*									-0.9*	-0.9*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-02	-0.2	2002-15	-0.8*									-0.8*	-0.8*
White	1992-02	-0.1	2002-15	-0.7*									-0.7*	-0.7*
White NH ^{ef}	1992-02	-0.1	2002-15	-0.8*									-0.8*	-0.8*
Black	1992-95	3.0	1995-15	-0.9*									-0.9*	-0.9*
Black NH ^{ef}	1992-95	2.9	1995-15	-0.8*									-0.8*	-0.8*
API ^e	1992-15	-0.1											-0.1	-0.1
AI/AN ^{eg}	1992-15	-0.2											-0.2	-0.2
Hispanic ^f	1992-15	-0.1											-0.1	-0.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 11.2
Cancer of the Kidney and Renal Pelvis (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-04	1.8*	2004-07	4.8	2007-15	0.6							1.1*	0.6
White	1975-03	1.7*	2003-07	3.8	2007-15	0.6							1.0*	0.6
Black	1975-15	2.8*											2.8*	2.8*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-99	1.2*	1999-08	3.1*	2008-15	0.4							1.0*	0.4
White	1992-97	0.9	1997-08	2.8*	2008-15	0.5							1.0*	0.5
Black	1992-15	2.4*											2.4*	2.4*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-04	1.8*	2004-07	4.8	2007-15	0.3							0.8	0.3
White	1975-89	2.2*	1989-97	0.8	1997-08	2.7*	2008-15	0.2					0.7*	0.2
Black	1975-15	2.7*											2.7*	2.7*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-99	1.2*	1999-08	3.1*	2008-15	0.1							0.8*	0.1
White	1992-97	0.9	1997-08	2.8*	2008-15	0.1							0.7*	0.1
White NH ^{ef}	1992-99	1.4*	1999-08	3.0*	2008-15	0.1							0.7*	0.1
Black	1992-15	2.3*											2.3*	2.3*
Black NH ^{ef}	1992-15	2.3*											2.3*	2.3*
API ^e	1992-15	2.4*											2.4*	2.4*
AI/AN ^{eg}	1992-15	0.9											0.9	0.9
Hispanic ^f	1992-07	3.0*	2007-15	0.2									0.5	0.2
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-91	1.1*	1991-01	-0.1	2001-15	-0.7*							-0.7*	-0.7*
White	1975-91	1.0*	1991-01	-0.1	2001-15	-0.6*							-0.6*	-0.6*
Black	1975-91	3.1*	1991-15	-0.6*									-0.6*	-0.6*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-0.5*											-0.5*	-0.5*
White	1992-01	-0.1	2001-15	-0.6*									-0.6*	-0.6*
White NH ^{ef}	1992-01	-0.1	2001-15	-0.6*									-0.6*	-0.6*
Black	1992-15	-0.6*											-0.6*	-0.6*
Black NH ^{ef}	1992-15	-0.6*											-0.6*	-0.6*
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-15	-0.3											-0.3	-0.3
Hispanic ^f	1992-15	-0.2											-0.2	-0.2

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 11.3
Cancer of the Kidney and Renal Pelvis (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program, 1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints, Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-93	2.7*	1993-99	0.6	1999-06	4.2*	2006-15	0.0					0.0	0.0
White	1975-92	2.8*	1992-99	0.7	1999-06	4.3*	2006-15	-0.1					-0.1	-0.1
Black	1975-08	3.0*	2008-15	-0.4									0.4	-0.4
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-08	2.6*	2008-15	0.0									0.6	0.0
White	1992-97	0.5	1997-07	3.3*	2007-15	0.0							0.4	0.0
Black	1992-09	2.8*	2009-15	-0.8									0.4	-0.8
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-93	2.7*	1993-99	0.6	1999-06	4.2*	2006-15	-0.3					-0.3	-0.3
White	1975-92	2.8*	1992-99	0.7	1999-06	4.3*	2006-15	-0.5					-0.5	-0.5
Black	1975-08	3.0*	2008-15	-0.9									-0.1	-0.9
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-08	2.6*	2008-15	-0.3									0.3	-0.3
White	1992-97	0.5	1997-07	3.3*	2007-15	-0.3							0.1	-0.3
White NH ^{ef}	1992-97	0.2	1997-07	3.3*	2007-15	-0.8							-0.3	-0.8
Black	1992-09	2.8*	2009-15	-1.4									0.0	-1.4
Black NH ^{ef}	1992-08	3.0*	2008-15	-0.8									0.0	-0.8
API ^e	1992-07	3.8*	2007-15	0.3									0.7	0.3
AI/AN ^{eg}	1992-15	0.8											0.8	0.8
Hispanic ^f	1992-15	2.5*											2.5*	2.5*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-95	1.1*	1995-15	-1.1*									-1.1*	-1.1*
White	1975-95	1.1*	1995-15	-1.0*									-1.0*	-1.0*
Black	1975-95	2.3*	1995-15	-1.2*									-1.2*	-1.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-05	-0.4*	2005-15	-1.6*									-1.6*	-1.6*
White	1992-05	-0.4*	2005-15	-1.5*									-1.5*	-1.5*
White NH ^{ef}	1992-05	-0.4*	2005-15	-1.6*									-1.6*	-1.6*
Black	1992-15	-0.9*											-0.9*	-0.9*
Black NH ^{ef}	1992-15	-0.8*											-0.8*	-0.8*
API ^e	1992-15	-0.4											-0.4	-0.4
AI/AN ^{eg}	1992-15	-0.6											-0.6	-0.6
Hispanic ^f	1992-15	-0.1											-0.1	-0.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 11.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Kidney and Renal Pelvis, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 11.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Kidney and Renal Pelvis, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 11.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Kidney and Renal Pelvis, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 12.1
Cancer of the Larynx (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-88	0.0	1988-15	-2.4*									-2.4*	-2.4*
White	1975-87	0.1	1987-15	-2.3*									-2.3*	-2.3*
Black	1975-90	1.2*	1990-15	-2.9*									-2.9*	-2.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-2.6*											-2.6*	-2.6*
White	1992-15	-2.4*											-2.4*	-2.4*
Black	1992-15	-3.2*											-3.2*	-3.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-88	0.0	1988-15	-2.4*									-2.4*	-2.4*
White	1975-87	0.1	1987-15	-2.3*									-2.3*	-2.3*
Black	1975-90	1.2*	1990-15	-2.9*									-2.9*	-2.9*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-2.7*											-2.7*	-2.7*
White	1992-15	-2.4*											-2.4*	-2.4*
White NH ^{ef}	1992-15	-2.4*											-2.4*	-2.4*
Black	1992-15	-3.3*											-3.3*	-3.3*
Black NH ^{ef}	1992-15	-3.3*											-3.3*	-3.3*
API ^e	1992-15	-3.3*											-3.3*	-3.3*
AI/AN ^{eg}	1992-15	0.1											0.1	0.1
Hispanic ^f	1992-15	-2.0*											-2.0*	-2.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-94	-0.4*	1994-15	-2.3*									-2.3*	-2.3*
White	1975-94	-0.7*	1994-15	-1.9*									-1.9*	-1.9*
Black	1975-94	1.1*	1994-15	-3.6*									-3.6*	-3.6*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-2.2*											-2.2*	-2.2*
White	1992-15	-1.9*											-1.9*	-1.9*
White NH ^{ef}	1992-15	-1.8*											-1.8*	-1.8*
Black	1992-15	-3.4*											-3.4*	-3.4*
Black NH ^{ef}	1992-15	-3.3*											-3.3*	-3.3*
API ^e	1992-15	-1.9*											-1.9*	-1.9*
AI/AN ^{eg}	1992-15	-2.7*											-2.7*	-2.7*
Hispanic ^f	1992-15	-2.6*											-2.6*	-2.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 12.2
Cancer of the Larynx (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-87	-0.2	1987-15	-2.5*									-2.5*	-2.5*
White	1975-85	0.1	1985-15	-2.4*									-2.4*	-2.4*
Black	1975-90	0.9	1990-15	-2.8*									-2.8*	-2.8*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-2.7*											-2.7*	-2.7*
White	1992-03	-3.1*	2003-10	-1.1	2010-15	-4.0*							-2.7*	-4.0*
Black	1992-15	-3.2*											-3.2*	-3.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-87	-0.2	1987-15	-2.5*									-2.5*	-2.5*
White	1975-85	0.2	1985-15	-2.4*									-2.4*	-2.4*
Black	1975-90	1.0	1990-15	-2.8*									-2.8*	-2.8*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-03	-3.2*	2003-10	-1.5*	2010-15	-4.3*							-3.1*	-4.3*
White	1992-03	-3.1*	2003-12	-1.4*	2012-15	-6.2*							-3.1*	-5.0*
White NH ^{ef}	1992-15	-2.5*											-2.5*	-2.5*
Black	1992-15	-3.3*											-3.3*	-3.3*
Black NH ^{ef}	1992-15	-3.3*											-3.3*	-3.3*
API ^e	1992-15	-3.3*											-3.3*	-3.3*
AI/AN ^{eg}	1992-15	0.8											0.8	0.8
Hispanic ^f	1992-15	-1.9*											-1.9*	-1.9*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-94	-0.8*	1994-15	-2.5*									-2.5*	-2.5*
White	1975-93	-1.1*	1993-15	-2.2*									-2.2*	-2.2*
Black	1975-94	1.2*	1994-15	-3.6*									-3.6*	-3.6*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-2.5*											-2.5*	-2.5*
White	1992-15	-2.2*											-2.2*	-2.2*
White NH ^{ef}	1992-15	-2.1*											-2.1*	-2.1*
Black	1992-15	-3.4*											-3.4*	-3.4*
Black NH ^{ef}	1992-15	-3.3*											-3.3*	-3.3*
API ^e	1992-15	-1.6*											-1.6*	-1.6*
AI/AN ^{eg}	1992-15	-1.7											-1.7	-1.7
Hispanic ^f	1992-15	-2.7*											-2.7*	-2.7*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 12.3
Cancer of the Larynx (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-89	1.8*	1989-15	-2.6*									-2.6*	-2.6*
White	1975-89	1.7*	1989-15	-2.4*									-2.4*	-2.4*
Black	1975-89	4.0*	1989-15	-2.9*									-2.9*	-2.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-3.1*											-3.1*	-3.1*
White	1992-15	-2.9*											-2.9*	-2.9*
Black	1992-15	-3.2*											-3.2*	-3.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-89	1.9*	1989-15	-2.6*									-2.6*	-2.6*
White	1975-89	1.7*	1989-15	-2.4*									-2.4*	-2.4*
Black	1975-89	4.0*	1989-15	-3.0*									-3.0*	-3.0*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-3.1*											-3.1*	-3.1*
White	1992-15	-2.9*											-2.9*	-2.9*
White NH ^{ef}	1992-15	-2.6*											-2.6*	-2.6*
Black	1992-15	-3.3*											-3.3*	-3.3*
Black NH ^{ef}	1992-15	-3.2*											-3.2*	-3.2*
API ^e	1992-15	-2.7*											-2.7*	-2.7*
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-3.0*											-3.0*	-3.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-93	1.4*	1993-15	-2.2*									-2.2*	-2.2*
White	1975-92	1.5*	1992-15	-1.8*									-1.8*	-1.8*
Black	1975-93	2.4*	1993-15	-3.4*									-3.4*	-3.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-2.2*											-2.2*	-2.2*
White	1992-15	-1.9*											-1.9*	-1.9*
White NH ^{ef}	1992-15	-1.6*											-1.6*	-1.6*
Black	1992-15	-3.3*											-3.3*	-3.3*
Black NH ^{ef}	1992-15	-3.2*											-3.2*	-3.2*
API ^e	1992-15	-3.2*											-3.2*	-3.2*
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-2.8*											-2.8*	-2.8*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 12.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Larynx, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 12.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Larynx, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 12.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Larynx, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 13.1
LeukemiaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-99	0.2*	1999-15	0.9*									0.9*	0.9*
White	1975-99	0.3*	1999-15	1.1*									1.1*	1.1*
Black	1975-15	0.2											0.2	0.2
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.6*											0.6*	0.6*
White	1992-15	0.8*											0.8*	0.8*
Black	1992-15	0.6*											0.6*	0.6*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	0.3*											0.3*	0.3*
White	1975-15	0.4*											0.4*	0.4*
Black	1975-15	0.0											0.0	0.0
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-12	0.4*	2012-15	-2.1									-0.4	-1.5
White	1992-11	0.6*	2011-15	-1.3									-0.3	-1.3
White NH ^{ef}	1992-15	0.5*											0.5*	0.5*
Black	1992-15	0.2											0.2	0.2
Black NH ^{ef}	1992-15	0.2											0.2	0.2
API ^e	1992-15	-0.1											-0.1	-0.1
AI/AN ^{eg}	1992-15	1.7*											1.7*	1.7*
Hispanic ^f	1992-13	0.7*	2013-15	-6.5									-0.9	-3.0
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-80	0.6	1980-88	-0.6*	1988-91	0.7	1991-01	-0.6*	2001-13	-0.9*	2013-15	-3.4*	-1.5*	-2.2*
White	1975-80	0.5	1980-88	-0.7*	1988-91	0.7	1991-01	-0.5*	2001-13	-0.8*	2013-15	-3.1*	-1.3*	-2.0*
Black	1975-93	0.5*	1993-15	-1.4*									-1.4*	-1.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	-0.6*	2001-13	-0.9*	2013-15	-3.4*							-1.5*	-2.2*
White	1992-01	-0.4*	2001-13	-0.8*	2013-15	-3.1*							-1.3*	-2.0*
White NH ^{ef}	1992-01	-0.3*	2001-13	-0.8*	2013-15	-3.1*							-1.3*	-2.0*
Black	1992-15	-1.3*											-1.3*	-1.3*
Black NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
API ^e	1992-07	-1.4*	2007-11	3.0	2011-15	-4.9*							-1.1	-4.9*
AI/AN ^{eg}	1992-15	-0.4											-0.4	-0.4
Hispanic ^f	1992-96	2.4	1996-10	-0.9*	2010-13	2.2	2013-15	-7.0*					-1.3	-2.5

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 13.2
LeukemiaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-98	0.0	1998-15	0.8*									0.8*	0.8*
White	1975-98	0.1	1998-15	0.9*									0.9*	0.9*
Black	1975-15	0.2											0.2	0.2
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.5*											0.5*	0.5*
White	1992-15	0.6*											0.6*	0.6*
Black	1992-15	0.5*											0.5*	0.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	0.2*											0.2*	0.2*
White	1975-15	0.3*											0.3*	0.3*
Black	1975-15	0.0											0.0	0.0
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-12	0.3*	2012-15	-2.3									-0.6	-1.7
White	1992-11	0.4*	2011-15	-1.6									-0.5	-1.6
White NH ^{ef}	1992-15	0.3*											0.3*	0.3*
Black	1992-15	0.1											0.1	0.1
Black NH ^{ef}	1992-15	0.2											0.2	0.2
API ^e	1992-15	-0.1											-0.1	-0.1
AI/AN ^{eg}	1992-15	2.6*											2.6*	2.6*
Hispanic ^f	1992-15	0.2											0.2	0.2
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-96	-0.2*	1996-13	-0.8*	2013-15	-3.7*							-1.5*	-2.3*
White	1975-00	-0.3*	2000-13	-0.8*	2013-15	-3.2							-1.4*	-2.0*
Black	1975-93	0.8*	1993-15	-1.5*									-1.5*	-1.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-02	-0.6*	2002-13	-0.9*	2013-15	-3.4*							-1.5*	-2.2*
White	1992-13	-0.7*	2013-15	-3.6*									-1.4*	-2.1*
White NH ^{ef}	1992-12	-0.6*	2012-15	-2.6*									-1.3*	-2.1*
Black	1992-15	-1.4*											-1.4*	-1.4*
Black NH ^{ef}	1992-15	-1.3*											-1.3*	-1.3*
API ^e	1992-15	-0.6*											-0.6*	-0.6*
AI/AN ^{eg}	1992-15	-0.1											-0.1	-0.1
Hispanic ^f	1992-15	-0.6*											-0.6*	-0.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 13.3
LeukemiaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-99	0.2	1999-15	1.0*									1.0*	1.0*
White	1975-99	0.3*	1999-15	1.1*									1.1*	1.1*
Black	1975-15	0.3											0.3	0.3
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.7*											0.7*	0.7*
White	1992-15	0.9*											0.9*	0.9*
Black	1992-15	0.7*											0.7*	0.7*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	0.3*											0.3*	0.3*
White	1975-15	0.5*											0.5*	0.5*
Black	1975-15	0.1											0.1	0.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	0.3*											0.3*	0.3*
White	1992-15	0.5*											0.5*	0.5*
White NH ^{ef}	1992-15	0.5*											0.5*	0.5*
Black	1992-15	0.3											0.3	0.3
Black NH ^{ef}	1992-15	0.3											0.3	0.3
API ^e	1992-15	-0.1											-0.1	-0.1
AI/AN ^{eg}	1992-15	0.9											0.9	0.9
Hispanic ^f	1992-15	0.7*											0.7*	0.7*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-80	0.8	1980-00	-0.4*	2000-15	-1.3*							-1.3*	-1.3*
White	1975-00	-0.3*	2000-15	-1.2*									-1.2*	-1.2*
Black	1975-96	0.2	1996-15	-1.5*									-1.5*	-1.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-00	-0.5*	2000-13	-1.1*	2013-15	-3.5*							-1.7*	-2.4*
White	1992-01	-0.4*	2001-15	-1.2*									-1.2*	-1.2*
White NH ^{ef}	1992-01	-0.3	2001-15	-1.2*									-1.2*	-1.2*
Black	1992-15	-1.3*											-1.3*	-1.3*
Black NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
API ^e	1992-15	-1.0*											-1.0*	-1.0*
AI/AN ^{eg}	1992-15	-1.1											-1.1	-1.1
Hispanic ^f	1992-15	-0.6*											-0.6*	-0.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Figure 13.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Leukemia, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 13.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Leukemia, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 13.5

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Leukemia, Female, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero (p < 0.05).

Table 14.1
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-84	1.3*	1984-98	4.8*	1998-01	0.2	2001-09	4.7*	2009-15	2.0*			2.9*	2.0*
White	1975-87	1.6*	1987-98	5.4*	1998-01	-0.8	2001-11	4.9*	2011-15	2.0			3.6*	2.0
Black	1975-02	2.9*	2002-09	6.1*	2009-15	0.5							2.3*	0.5
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-99	5.1*	1999-02	0.5	2002-09	4.5*	2009-15	1.2*					2.3*	1.2*
White	1992-99	5.5*	1999-02	-0.6	2002-09	5.5*	2009-15	2.0*					3.2*	2.0*
Black	1992-15	3.5*											3.5*	3.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-84	1.3*	1984-98	4.8*	1998-01	0.2	2001-09	4.7*	2009-15	1.5*			2.6*	1.5*
White	1975-87	1.6*	1987-98	5.4*	1998-01	-0.8	2001-11	4.8*	2011-15	1.2			3.2*	1.2
Black	1975-02	2.9*	2002-09	6.1*	2009-15	0.0							2.0*	0.0
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-99	5.1*	1999-02	0.5	2002-09	4.5*	2009-15	0.6					1.9*	0.6
White	1992-99	5.5*	1999-02	-0.1	2002-11	4.9*	2011-15	0.1					2.7*	0.1
White NH ^{ef}	1992-99	5.4*	1999-02	-2.2	2002-08	5.9*	2008-15	2.1*					3.0*	2.1*
Black	1992-09	4.3*	2009-15	0.8									1.9*	0.8
Black NH ^{ef}	1992-09	4.3*	2009-15	0.9									2.0*	0.9
API ^e	1992-07	1.1*	2007-15	-2.4*									-2.0*	-2.4*
AI/AN ^{eg}	1992-15	2.5*											2.5*	2.5*
Hispanic ^f	1992-12	3.3*	2012-15	-3.7									0.9	-2.0
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-80	0.2	1980-87	2.0*	1987-95	3.8*	1995-07	1.9*	2007-12	3.3*	2012-15	1.3*	2.5*	1.8*
White	1975-78	-1.2	1978-87	1.6*	1987-96	3.9*	1996-00	0.7	2000-15	2.7*			2.7*	2.7*
Black	1975-88	0.8*	1988-95	3.2*	1995-00	0.3	2000-13	2.5*	2013-15	-0.4			1.9*	1.1
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-96	3.5*	1996-99	0.7	1999-08	2.2*	2008-12	3.3*	2012-15	1.3*			2.4*	1.8*
White	1992-96	3.5*	1996-99	0.6	1999-08	2.4*	2008-12	3.5*	2012-15	1.6*			2.6*	2.1*
White NH ^{ef}	1992-96	3.7*	1996-99	0.1	1999-08	2.2*	2008-12	3.5*	2012-15	1.6*			2.6*	2.1*
Black	1992-95	4.2	1995-00	0.1	2000-13	2.5*	2013-15	-0.4					1.9*	1.1
Black NH ^{ef}	1992-95	4.7*	1995-00	0.1	2000-13	2.6*	2013-15	-0.4					1.9*	1.1
API ^e	1992-15	-0.8*											-0.8*	-0.8*
AI/AN ^{eg}	1992-15	2.2*											2.2*	2.2*
Hispanic ^f	1992-98	3.7*	1998-15	1.4*									1.4*	1.4*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 14.2
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-84	2.2*	1984-11	3.9*	2011-15	1.0							2.6*	1.0
White	1975-87	1.8*	1987-98	5.4*	1998-01	-0.7	2001-11	5.0*	2011-15	1.2			3.3*	1.2
Black	1975-02	3.2*	2002-09	6.5*	2009-15	0.2							2.3*	0.2
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-11	3.7*	2011-15	-0.2									1.9*	-0.2
White	1992-12	4.0*	2012-15	-0.6									2.5*	0.6
Black	1992-09	4.7*	2009-15	0.7									2.0*	0.7
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-82	1.7	1982-11	3.9*	2011-15	0.3							2.3*	0.3
White	1975-87	1.5*	1987-91	6.9	1991-12	3.9*	2012-15	0.3					2.7*	1.2
Black	1975-02	3.1*	2002-09	6.6*	2009-15	-0.3							2.0	-0.3
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-11	3.6*	2011-15	-1.1									1.5*	-1.1
White	1992-99	4.8*	1999-02	-0.5	2002-06	6.8*	2006-12	3.7*	2012-15	-1.8			1.8*	-0.5
White NH ^{ef}	1992-99	4.6*	1999-02	-1.2	2002-09	5.8*	2009-15	1.2					2.7*	1.2
Black	1992-09	4.7*	2009-15	0.1									1.6*	0.1
Black NH ^{ef}	1992-09	4.7*	2009-15	0.3									1.8*	0.3
API ^e	1992-07	1.3*	2007-15	-2.3*									-1.9*	-2.3*
AI/AN ^{eg}	1992-15	3.0*											3.0*	3.0*
Hispanic ^f	1992-12	3.3*	2012-15	-5.5									0.3	-3.4
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-85	1.5*	1985-96	3.8*	1996-99	0.3	1999-13	2.7*	2013-15	0.6			2.2*	1.6*
White	1975-84	1.1*	1984-95	3.7*	1995-06	2.0*	2006-12	3.4*	2012-15	1.1			2.6*	1.7*
Black	1975-87	0.8	1987-15	2.3*									2.3*	2.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-07	2.2*	2007-12	3.3*	2012-15	0.9							2.4*	1.5*
White	1992-07	2.2*	2007-12	3.5*	2012-15	1.1							2.5*	1.7*
White NH ^{ef}	1992-96	3.6*	1996-99	0.6	1999-07	2.3*	2007-12	3.4*	2012-15	1.3*			2.6*	1.8*
Black	1992-15	2.3*											2.3*	2.3*
Black NH ^{ef}	1992-15	2.4*											2.4*	2.4*
API ^e	1992-15	-0.8*											-0.8*	-0.8*
AI/AN ^{eg}	1992-15	2.5*											2.5*	2.5*
Hispanic ^f	1992-12	1.9*	2012-15	-1.3									0.8	-0.5

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 14.3
Cancer of the Liver and Intrahepatic Bile Duct (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program, 1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints, Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-83	0.4	1983-98	4.4*	1998-01	-0.4	2001-15	3.4*					3.4*	3.4*
White	1975-85	0.8	1985-99	4.1*	1999-02	-2.4	2002-15	4.1*					4.1*	4.1*
Black	1975-15	2.8*											2.8*	2.8*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	2.9*											2.9*	2.9*
White	1992-98	6.8*	1998-03	-0.3	2003-15	4.0*							4.0*	4.0*
Black	1992-15	3.0*											3.0*	3.0*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-83	0.6	1983-96	4.2*	1996-15	2.7*							2.7*	2.7*
White	1975-85	0.8	1985-99	4.1*	1999-02	-2.1	2002-15	3.9*					3.9*	3.9*
Black	1975-15	2.8*											2.8*	2.8*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-96	6.8*	1996-15	2.4*									2.4*	2.4*
White	1992-98	6.8*	1998-03	-0.1	2003-15	3.8*							3.8*	3.8*
White NH ^{ef}	1992-98	6.6*	1998-03	-1.3	2003-15	3.7*							3.7*	3.7*
Black	1992-15	2.8*											2.8*	2.8*
Black NH ^{ef}	1992-15	2.9*											2.9*	2.9*
API ^e	1992-06	1.1*	2006-15	-2.1*									-2.1*	-2.1*
AI/AN ^{eg}	1992-15	1.2											1.2	1.2
Hispanic ^f	1992-15	2.6*											2.6*	2.6*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-78	-1.5	1978-88	1.4*	1988-95	3.9*	1995-00	0.4	2000-08	1.5*	2008-15	2.7*	2.4*	2.7*
White	1975-78	-1.9	1978-88	1.2*	1988-96	3.6*	1996-00	-0.4	2000-08	1.8*	2008-15	2.8*	2.6*	2.8*
Black	1975-15	1.4*											1.4*	1.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-95	3.8*	1995-00	0.4	2000-08	1.5*	2008-15	2.7*					2.4*	2.7*
White	1992-96	3.1*	1996-00	-0.2	2000-08	1.8*	2008-15	2.9*					2.6*	2.9*
White NH ^{ef}	1992-96	3.1*	1996-01	-0.6	2001-15	2.2*							2.2*	2.2*
Black	1992-15	1.2*											1.2*	1.2*
Black NH ^{ef}	1992-15	1.3*											1.3*	1.3*
API ^e	1992-15	-0.7*											-0.7*	-0.7*
AI/AN ^{eg}	1992-15	1.3*											1.3*	1.3*
Hispanic ^f	1992-15	1.6*											1.6*	1.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 14.1

SEER Observed Incidence and Delay Adjusted Incidence Rates^a Cancer of the Liver and Intrahepatic Bile Duct, by Race and Sex




Source: SEER 9 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 14.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Liver and Intrahepatic Bile Duct, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 14.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Liver and Intrahepatic Bile Duct, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 15.1
Cancer of the Lung and Bronchus (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-82	2.5*	1982-91	0.9*	1991-07	-0.7*	2007-15	-2.2*					-2.0*	-2.2*
White	1975-82	2.6*	1982-91	1.1*	1991-07	-0.6*	2007-15	-2.1*					-1.9*	-2.1*
Black	1975-84	3.5*	1984-03	-0.6*	2003-15	-2.3*							-2.3*	-2.3*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-07	-1.0*	2007-15	-2.3*									-2.2*	-2.3*
White	1992-07	-0.9*	2007-15	-2.3*									-2.1*	-2.3*
Black	1992-08	-1.3*	2008-15	-3.2*									-2.8*	-3.2*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-82	2.5*	1982-91	0.9*	1991-07	-0.7*	2007-15	-2.3*					-2.1*	-2.3*
White	1975-82	2.6*	1982-91	1.1*	1991-07	-0.6*	2007-15	-2.3*					-2.1*	-2.3*
Black	1975-85	3.2*	1985-08	-0.9*	2008-15	-3.5*							-3.0*	-3.5*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-07	-1.0*	2007-15	-2.5*									-2.3*	-2.5*
White	1992-07	-0.9*	2007-15	-2.5*									-2.3*	-2.5*
White NH ^{ef}	1992-07	-0.7*	2007-15	-2.3*									-2.2*	-2.3*
Black	1992-08	-1.3*	2008-15	-3.5*									-3.0*	-3.5*
Black NH ^{ef}	1992-08	-1.2*	2008-15	-3.4*									-2.9*	-3.4*
API ^e	1992-05	-0.4	2005-15	-1.5*									-1.5*	-1.5*
AI/AN ^{eg}	1992-15	-1.2*											-1.2*	-1.2*
Hispanic ^f	1992-15	-1.3*											-1.3*	-1.3*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-80	3.0*	1980-90	1.8*	1990-95	-0.2	1995-05	-1.0*	2005-12	-2.3*	2012-15	-3.4*	-2.7*	-3.1*
White	1975-80	2.9*	1980-90	1.8*	1990-95	-0.1	1995-05	-0.8*	2005-10	-2.1*	2010-15	-2.9*	-2.5*	-2.9*
Black	1975-81	3.5*	1981-90	1.9*	1990-95	-0.7*	1995-04	-1.7*	2004-13	-2.7*	2013-15	-5.0*	-3.2*	-3.8*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-04	-0.8*	2004-10	-2.1*	2010-15	-3.0*							-2.6*	-3.0*
White	1992-03	-0.7*	2003-09	-1.7*	2009-15	-2.8*							-2.4*	-2.8*
White NH ^{ef}	1992-03	-0.5*	2003-09	-1.5*	2009-15	-2.7*							-2.3*	-2.7*
Black	1992-97	-1.1*	1997-04	-1.8*	2004-13	-2.6*	2013-15	-5.0*					-3.2*	-3.8*
Black NH ^{ef}	1992-95	-0.6	1995-04	-1.7*	2004-13	-2.6*	2013-15	-5.0*					-3.1*	-3.8*
API ^e	1992-08	-1.1*	2008-15	-1.8*									-1.6*	-1.8*
AI/AN ^{eg}	1992-08	0.2	2008-15	-2.4*									-1.8*	-2.4*
Hispanic ^f	1992-95	1.4	1995-04	-1.2*	2004-15	-2.4*							-2.4*	-2.4*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 15.2
Cancer of the Lung and Bronchus (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-82	1.5*	1982-91	-0.5*	1991-08	-1.7*	2008-15	-2.9*					-2.6*	-2.9*
White	1975-81	1.5*	1981-90	-0.2	1990-08	-1.7*	2008-15	-2.8*					-2.6*	-2.8*
Black	1975-84	2.9*	1984-98	-1.4*	1998-15	-2.9*							-2.9*	-2.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-08	-1.9*	2008-15	-3.0*									-2.7*	-3.0*
White	1992-08	-1.9*	2008-15	-2.9*									-2.7*	-2.9*
Black	1992-15	-2.7*											-2.7*	-2.7*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-82	1.5*	1982-91	-0.5*	1991-08	-1.7*	2008-15	-3.1*					-2.8*	-3.1*
White	1975-81	1.5*	1981-90	-0.2	1990-09	-1.7*	2009-15	-3.3*					-2.8*	-3.3*
Black	1975-84	2.9*	1984-98	-1.3*	1998-15	-3.0*							-3.0*	-3.0*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-08	-1.9*	2008-15	-3.2*									-2.9*	-3.2*
White	1992-09	-1.9*	2009-15	-3.4*									-2.9*	-3.4*
White NH ^{ef}	1992-09	-1.8*	2009-15	-3.2*									-2.7*	-3.2*
Black	1992-09	-2.4*	2009-15	-4.8*									-4.0*	-4.8*
Black NH ^{ef}	1992-09	-2.4*	2009-15	-4.6*									-3.9*	-4.6*
API ^e	1992-05	-0.8*	2005-15	-2.2*									-2.2*	-2.2*
AI/AN ^{eg}	1992-15	-1.8*											-1.8*	-1.8*
Hispanic ^f	1992-15	-2.1*											-2.1*	-2.1*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-78	2.4*	1978-84	1.2*	1984-91	0.3*	1991-05	-1.9*	2005-12	-3.0*	2012-15	-4.0*	-3.3*	-3.7*
White	1975-80	1.9*	1980-87	0.7*	1987-91	0.1	1991-05	-1.7*	2005-12	-2.9*	2012-15	-3.9*	-3.2*	-3.6*
Black	1975-82	2.9*	1982-90	1.3*	1990-94	-1.3	1994-03	-2.7*	2003-13	-3.4*	2013-15	-5.4*	-3.9*	-4.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-05	-1.9*	2005-12	-3.0*	2012-15	-4.0*							-3.3*	-3.7*
White	1992-05	-1.8*	2005-11	-2.8*	2011-15	-3.7*							-3.2*	-3.7*
White NH ^{ef}	1992-05	-1.6*	2005-12	-2.8*	2012-15	-3.8*							-3.1*	-3.6*
Black	1992-94	-1.0	1994-03	-2.7*	2003-13	-3.4*	2013-15	-5.4*					-3.8*	-4.4*
Black NH ^{ef}	1992-94	-0.6	1994-04	-2.7*	2004-13	-3.3*	2013-15	-5.4*					-3.8*	-4.4*
API ^e	1992-08	-1.4*	2008-15	-2.7*									-2.4*	-2.7*
AI/AN ^{eg}	1992-15	-1.2*											-1.2*	-1.2*
Hispanic ^f	1992-94	3.2	1994-05	-2.0*	2005-15	-3.4*							-3.4*	-3.4*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 15.3
Cancer of the Lung and Bronchus (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-82	5.6*	1982-91	3.4*	1991-06	0.5*	2006-15	-1.5*					-1.5*	-1.5*
White	1975-88	4.8*	1988-97	1.4*	1997-07	0.2	2007-15	-1.5*					-1.3*	-1.5*
Black	1975-90	4.2*	1990-06	0.6*	2006-15	-1.8*							-1.8*	-1.8*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-06	0.1	2006-15	-1.7*									-1.7*	-1.7*
White	1992-97	0.8	1997-07	-0.2	2007-15	-1.7*							-1.6*	-1.7*
Black	1992-05	0.8*	2005-15	-1.7*									-1.7*	-1.7*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-82	5.6*	1982-91	3.4*	1991-07	0.4*	2007-15	-1.8*					-1.6*	-1.8*
White	1975-88	4.8*	1988-97	1.4*	1997-07	0.3	2007-15	-1.7*					-1.5*	-1.7*
Black	1975-90	4.2*	1990-06	0.6*	2006-15	-1.9*							-1.9*	-1.9*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-07	0.0	2007-15	-2.0*									-1.8*	-2.0*
White	1992-07	0.0	2007-15	-2.0*									-1.8*	-2.0*
White NH ^{ef}	1992-07	0.2*	2007-15	-1.9*									-1.7*	-1.9*
Black	1992-05	0.8*	2005-15	-1.9*									-1.9*	-1.9*
Black NH ^{ef}	1992-05	0.8*	2005-15	-1.9*									-1.9*	-1.9*
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-05	2.4	2005-15	-3.6*									-3.6*	-3.6*
Hispanic ^f	1992-15	-0.4*											-0.4*	-0.4*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-82	6.0*	1982-90	4.2*	1990-95	1.7*	1995-03	0.3*	2003-09	-1.1*	2009-15	-2.3*	-1.9*	-2.3*
White	1975-83	5.8*	1983-90	4.1*	1990-94	2.1*	1994-03	0.5*	2003-09	-1.0*	2009-15	-2.2*	-1.8*	-2.2*
Black	1975-82	6.3*	1982-92	3.7*	1992-04	0.3*	2004-13	-1.9*	2013-15	-4.7*			-2.5*	-3.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-95	1.3*	1995-03	0.4*	2003-09	-1.1*	2009-15	-2.3*					-1.9*	-2.3*
White	1992-95	1.4*	1995-03	0.5*	2003-09	-0.9*	2009-15	-2.2*					-1.8*	-2.2*
White NH ^{ef}	1992-95	1.7*	1995-03	0.6*	2003-09	-0.8*	2009-15	-2.1*					-1.7*	-2.1*
Black	1992-04	0.4*	2004-13	-1.9*	2013-15	-4.7*							-2.5*	-3.3*
Black NH ^{ef}	1992-97	1.3*	1997-05	0.0	2005-13	-1.8*	2013-15	-4.6*					-2.5*	-3.2*
API ^e	1992-15	-0.5*											-0.5*	-0.5*
AI/AN ^{eg}	1992-05	2.1*	2005-15	-1.7*									-1.7*	-1.7*
Hispanic ^f	1992-03	0.5	2003-15	-1.2*									-1.2*	-1.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 15.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Lung and Bronchus, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 15.3

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Lung and Bronchus, Ages 65 and Over, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).


^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).


Cancer of the Lung & Bronchus SEER Delay-Adjusted Incidence & US Death Rates, 1975-2015, All Races, By Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

Figure 15.6


Cancer of the Lung & Bronchus SEER Delay-Adjusted Incidence, 1975-2015 By Sex and Race



Source: SEER 9 areas. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

Figure 15.7

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Lung and Bronchus, Male, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero (p < 0.05).

Figure 15.8

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Lung and Bronchus, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero (p < 0.05).

Table 16.1
Melanoma of the Skin (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-81	5.9*	1981-05	2.9*	2005-15	1.7*							1.7*	1.7*
White	1975-81	6.4*	1981-08	3.2*	2008-12	0.0	2012-15	5.0*					2.4*	3.8*
Black	1975-15	-0.1											-0.1	-0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-96	4.8*	1996-15	1.8*									1.8*	1.8*
White	1992-08	2.9*	2008-12	-0.3	2012-15	4.7*							2.1	3.5*
Black	1992-94	37.2	1994-15	-0.9									-0.9	-0.9
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-81	5.9*	1981-05	2.9*	2005-15	1.5*							1.5*	1.5*
White	1975-81	6.2*	1981-05	3.2*	2005-15	1.7*							1.7*	1.7*
Black	1975-15	-0.2											-0.2	-0.2
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-96	5.0*	1996-15	1.7*									1.7*	1.7*
White	1992-08	2.9*	2008-12	-0.6	2012-15	4.0*							1.7	2.8
White NH ^{ef}	1992-05	3.5*	2005-15	1.7*									1.7*	1.7*
Black	1992-94	37.8	1994-15	-1.0									-1.0	-1.0
Black NH ^{ef}	1992-94	36.9	1994-15	-0.9									-0.9	-0.9
API ^e	1992-15	0.6											0.6	0.6
AI/AN ^{eg}	1992-15	2.9*											2.9*	2.9*
Hispanic ^f	1992-15	0.9*											0.9*	0.9*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
Note- Alternate mortality trends adjusted for ICD-9/ICD-10 coding change available at: https://surveillance.cancer.gov/joinpoint/jump.html .														
All Races	1975-89	1.5*	1989-13	0.0	2013-15	-5.3*							-1.2*	-2.7*
White	1975-88	1.8*	1988-13	0.2*	2013-15	-5.0*							-1.0	-2.5*
Black	1975-15	-0.9*											-0.9*	-0.9*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-13	0.0	2013-15	-5.4*									-1.2*	-2.7*
White	1992-13	0.2*	2013-15	-5.1									-1.0	-2.5
White NH ^{ef}	1992-13	0.4*	2013-15	-5.0									-0.8	-2.3
Black	1992-15	-1.2*											-1.2*	-1.2*
Black NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
API ^e	1992-15	-1.0											-1.0	-1.0
AI/AN ^{eg}	1992-15	0.0											0.0	0.0
Hispanic ^f	1992-15	-0.4											-0.4	-0.4

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 16.2
Melanoma of the Skin (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-86	5.4*	1986-05	3.1*	2005-15	1.8*							1.8*	1.8*
White	1975-85	5.9*	1985-05	3.4*	2005-15	2.0*							2.0*	2.0*
Black	1975-15	-0.5											-0.5	-0.5
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-96	5.1*	1996-15	1.9*									1.9*	1.9*
White	1992-05	3.1*	2005-15	1.7*									1.7*	1.7*
Black	1992-95	51.9*	1995-15	-2.4*									-2.4*	-2.4*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-86	5.4*	1986-05	3.1*	2005-15	1.6*							1.6*	1.6*
White	1975-85	5.9*	1985-05	3.4*	2005-15	1.7*							1.7*	1.7*
Black	1975-15	-0.5											-0.5	-0.5
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-96	5.3*	1996-15	1.8*									1.8*	1.8*
White	1992-05	3.2*	2005-15	1.4*									1.4*	1.4*
White NH ^{ef}	1992-05	3.5*	2005-15	1.7*									1.7*	1.7*
Black	1992-95	52.3*	1995-15	-2.5*									-2.5*	-2.5*
Black NH ^{ef}	1992-95	52.6*	1995-15	-2.4*									-2.4*	-2.4*
API ^e	1992-15	1.1*											1.1*	1.1*
AI/AN ^{eg}	1992-15	2.2											2.2	2.2
Hispanic ^f	1992-15	1.3*											1.3*	1.3*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
Note- Alternate mortality trends adjusted for ICD-9/ICD-10 coding change available at: https://surveillance.cancer.gov/joinpoint/jump.html .														
All Races	1975-87	2.4*	1987-97	0.9*	1997-00	-1.7	2000-09	0.9*	2009-13	-0.9	2013-15	-5.0*	-1.2*	-3.0*
White	1975-87	2.5*	1987-97	1.0*	1997-00	-1.6	2000-09	1.1*	2009-13	-0.7	2013-15	-4.8*	-1.1	-2.8*
Black	1975-15	-0.8*											-0.8*	-0.8*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-13	0.3*	2013-15	-6.2									-1.2	-3.0*
White	1992-13	0.4*	2013-15	-5.9									-1.0	-2.8
White NH ^{ef}	1992-13	0.6*	2013-15	-5.7									-0.8	-2.6
Black	1992-15	-0.2											-0.2	-0.2
Black NH ^{ef}	1992-15	-0.1											-0.1	-0.1
API ^e	1992-15	-1.7*											-1.7*	-1.7*
AI/AN ^{eg}	1992-15	1.5											1.5	1.5
Hispanic ^f	1992-15	-0.3											-0.3	-0.3

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 16.3
Melanoma of the Skin (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-80	5.5*	1980-09	2.4*	2009-12	-1.3	2012-15	5.4*					2.1	3.7*
White	1975-81	6.2*	1981-93	2.1*	1993-05	3.5*	2005-12	0.8	2012-15	5.2*			2.3*	4.1*
Black	1975-15	-0.1											-0.1	-0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-96	4.3*	1996-08	1.9*	2008-12	-0.4	2012-15	5.1*					1.9*	3.7*
White	1992-06	3.0*	2006-12	0.2	2012-15	5.1*							1.8*	3.9*
Black	1992-15	0.1											0.1	0.1
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-86	4.0*	1986-93	0.6	1993-96	5.5	1996-09	2.3*	2009-12	-1.4	2012-15	5.0*	1.9	3.4*
White	1975-81	6.2*	1981-93	2.1*	1993-05	3.5*	2005-12	0.7	2012-15	4.5*			1.9*	3.5*
Black	1975-15	-0.1											-0.1	-0.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-96	4.3*	1996-08	1.9*	2008-12	-0.6	2012-15	4.7*					1.7	3.4*
White	1992-06	3.0*	2006-12	0.0	2012-15	4.4*							1.4	3.3*
White NH ^{ef}	1992-06	3.6*	2006-12	0.4	2012-15	4.6*							1.8*	3.5*
Black	1992-15	0.0											0.0	0.0
Black NH ^{ef}	1992-15	0.2											0.2	0.2
API ^e	1992-15	0.2											0.2	0.2
AI/AN ^{eg}	1992-15	3.3*											3.3*	3.3*
Hispanic ^f	1992-15	0.6											0.6	0.6
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
Note- Alternate mortality trends adjusted for ICD-9/ICD-10 coding change available at: https://surveillance.cancer.gov/joinpoint/jump.html .														
All Races	1975-89	0.8*	1989-15	-0.7*									-0.7*	-0.7*
White	1975-88	1.0*	1988-15	-0.4*									-0.4*	-0.4*
Black	1975-15	-1.0*											-1.0*	-1.0*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-12	-0.5*	2012-15	-3.2									-1.4*	-2.5*
White	1992-15	-0.4*											-0.4*	-0.4*
White NH ^{ef}	1992-15	-0.2											-0.2	-0.2
Black	1992-15	-1.9*											-1.9*	-1.9*
Black NH ^{ef}	1992-15	-1.9*											-1.9*	-1.9*
API ^e	1992-15	-0.3											-0.3	-0.3
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-0.6											-0.6	-0.6

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 16.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Melanoma of the Skin, White, by Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 16.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Melanoma of the Skin, Male, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 16.5

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Melanoma of the Skin, Female, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero (p < 0.05).

Table 18.1
MyelomaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-06	0.8*	2006-15	2.2*									2.2*	2.2*
White	1975-07	0.8*	2007-15	2.1*									2.0*	2.1*
Black	1975-96	1.3*	1996-02	-1.8	2002-11	3.4*	2011-15	-1.5					1.2	-1.5
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-07	0.4*	2007-10	4.1	2010-15	0.6							1.8	0.6
White	1992-06	0.5*	2006-15	2.0*									2.0*	2.0*
Black	1992-15	1.1*											1.1*	1.1*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	0.9*											0.9*	0.9*
White	1975-15	0.8*											0.8*	0.8*
Black	1975-96	1.3*	1996-02	-1.7	2002-11	3.0*	2011-15	-3.3					0.1	-3.3
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-07	0.4*	2007-10	3.7	2010-15	-0.6							0.9	-0.6
White	1992-15	0.7*											0.7*	0.7*
White NH ^{ef}	1992-15	0.8*											0.8*	0.8*
Black	1992-15	0.7*											0.7*	0.7*
Black NH ^{ef}	1992-15	0.8*											0.8*	0.8*
API ^e	1992-15	0.7*											0.7*	0.7*
AI/AN ^{eg}	1992-15	0.8											0.8	0.8
Hispanic ^f	1992-15	0.4*											0.4*	0.4*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-93	1.5*	1993-02	-0.5*	2002-08	-2.0*	2008-15	-0.1					-0.5*	-0.1
White	1975-94	1.4*	1994-02	-0.5*	2002-08	-1.9*	2008-15	-0.1					-0.5*	-0.1
Black	1975-82	3.4*	1982-96	1.2*	1996-09	-1.8*	2009-15	0.2					-0.5	0.2
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	1.9	1994-02	-0.6*	2002-09	-1.9*	2009-12	0.9	2012-15	-1.0			-0.7	-0.5
White	1992-02	-0.3	2002-08	-2.0*	2008-15	-0.1							-0.5*	-0.1
White NH ^{ef}	1992-94	2.0	1994-02	-0.5	2002-08	-2.0*	2008-15	-0.1					-0.5*	-0.1
Black	1992-98	0.1	1998-09	-2.0*	2009-15	0.2							-0.5	0.2
Black NH ^{ef}	1992-98	0.2	1998-09	-1.9*	2009-15	0.4							-0.4	0.4
API ^e	1992-15	-0.7*											-0.7*	-0.7*
AI/AN ^{eg}	1992-15	-1.5*											-1.5*	-1.5*
Hispanic ^f	1992-15	-0.7*											-0.7*	-0.7*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 18.2
MyelomaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-06	0.8*	2006-15	2.2*									2.2*	2.2*
White	1975-06	0.9*	2006-15	2.0*									2.0*	2.0*
Black	1975-07	0.6*	2007-10	8.5	2010-15	-2.9							1.1	-2.9
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	0.4	2004-15	1.9*									1.9*	1.9*
White	1992-06	0.6*	2006-15	2.1*									2.1*	2.1*
Black	1992-15	1.4*											1.4*	1.4*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	0.9*											0.9*	0.9*
White	1975-15	0.9*											0.9*	0.9*
Black	1975-05	0.6*	2005-12	3.2	2012-15	-7.7							-0.6	-5.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-06	0.3	2006-10	3.2	2010-15	-0.5							1.1	-0.5
White	1992-15	0.8*											0.8*	0.8*
White NH ^{ef}	1992-15	0.9*											0.9*	0.9*
Black	1992-15	1.0*											1.0*	1.0*
Black NH ^{ef}	1992-15	1.0*											1.0*	1.0*
API ^e	1992-15	0.8*											0.8*	0.8*
AI/AN ^{eg}	1992-15	1.3											1.3	1.3
Hispanic ^f	1992-15	0.4											0.4	0.4
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-94	1.5*	1994-10	-1.1*	2010-15	-0.2							-0.6*	-0.2
White	1975-94	1.5*	1994-15	-0.8*									-0.8*	-0.8*
Black	1975-91	2.3*	1991-15	-1.2*									-1.2*	-1.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	3.3	1994-10	-1.1*	2010-15	-0.1							-0.6*	-0.1
White	1992-94	3.0	1994-15	-0.9*									-0.9*	-0.9*
White NH ^{ef}	1992-94	3.4	1994-15	-0.9*									-0.9*	-0.9*
Black	1992-15	-1.3*											-1.3*	-1.3*
Black NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
API ^e	1992-15	-0.4											-0.4	-0.4
AI/AN ^{eg}	1992-15	-1.3*											-1.3*	-1.3*
Hispanic ^f	1992-15	-0.6*											-0.6*	-0.6*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 18.3
MyelomaTrends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-07	0.6*	2007-15	2.1*									2.0*	2.1*
White	1975-15	0.7*											0.7*	0.7*
Black	1975-15	1.1*											1.1*	1.1*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.8*											0.8*	0.8*
White	1992-15	0.7*											0.7*	0.7*
Black	1992-15	0.8*											0.8*	0.8*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	0.7*											0.7*	0.7*
White	1975-15	0.6*											0.6*	0.6*
Black	1975-15	0.9*											0.9*	0.9*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	0.5*											0.5*	0.5*
White	1992-15	0.4*											0.4*	0.4*
White NH ^{ef}	1992-15	0.4*											0.4*	0.4*
Black	1992-15	0.4											0.4	0.4
Black NH ^{ef}	1992-15	0.5*											0.5*	0.5*
API ^e	1992-15	0.7											0.7	0.7
AI/AN ^{eg}	1992-15	0.3											0.3	0.3
Hispanic ^f	1992-15	0.2											0.2	0.2
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-93	1.5*	1993-02	-0.5	2002-09	-2.7*	2009-12	1.8	2012-15	-1.7			-0.9	-0.9
White	1975-93	1.4*	1993-02	-0.5	2002-08	-2.6*	2008-15	-0.3					-0.8*	-0.3
Black	1975-96	2.0*	1996-09	-2.1*	2009-15	0.8							-0.2	0.8
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	-0.2	2001-08	-2.5*	2008-15	-0.1							-0.7*	-0.1
White	1992-02	-0.4	2002-08	-2.7*	2008-15	-0.3							-0.8*	-0.3
White NH ^{ef}	1992-02	-0.3	2002-08	-2.7*	2008-15	-0.3							-0.8*	-0.3
Black	1992-99	0.3	1999-09	-2.5*	2009-15	1.0							-0.1	1.0
Black NH ^{ef}	1992-99	0.4	1999-09	-2.4*	2009-15	1.1							0.0	1.1
API ^e	1992-15	-0.9											-0.9	-0.9
AI/AN ^{eg}	1992-15	-1.7											-1.7	-1.7
Hispanic ^f	1992-15	-0.9*											-0.9*	-0.9*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Figure 18.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Myeloma, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 18.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Myeloma, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 18.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Myeloma, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 19.1
Non-Hodgkin Lymphoma

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-91	3.6*	1991-07	0.7*	2007-15	-0.6*							-0.4	-0.6*
White	1975-90	3.8*	1990-07	0.8*	2007-15	-0.6*							-0.4	-0.6*
Black	1975-95	4.5*	1995-15	0.2									0.2	0.2
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-08	0.5*	2008-15	-0.7									-0.4	-0.7
White	1992-07	0.7*	2007-15	-0.5									-0.4	-0.5
Black	1992-97	4.0*	1997-00	-4.3	2000-04	3.8	2004-15	-0.6					-0.6	-0.6
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-91	3.6*	1991-07	0.7*	2007-15	-0.8*							-0.7*	-0.8*
White	1975-90	3.8*	1990-07	0.8*	2007-15	-0.9*							-0.7*	-0.9*
Black	1975-95	4.7*	1995-00	-2.6	2000-04	5.1	2004-15	-1.2*					-1.2*	-1.2*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-08	0.5*	2008-15	-1.0*									-0.6*	-1.0*
White	1992-07	0.7*	2007-15	-0.8*									-0.7*	-0.8*
White NH ^{ef}	1992-07	0.8*	2007-15	-0.9*									-0.7*	-0.9*
Black	1992-97	4.0*	1997-00	-4.3	2000-04	4.0	2004-15	-1.0*					-1.0*	-1.0*
Black NH ^{ef}	1992-97	3.8*	1997-00	-4.2	2000-04	4.0	2004-15	-0.8*					-0.8*	-0.8*
API ^e	1992-15	0.2											0.2	0.2
AI/AN ^{eg}	1992-15	1.3											1.3	1.3
Hispanic ^f	1992-15	0.4*											0.4*	0.4*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-91	2.5*	1991-97	1.6*	1997-06	-3.1*	2006-15	-2.2*					-2.2*	-2.2*
White	1975-91	2.5*	1991-97	1.6*	1997-06	-3.1*	2006-15	-2.2*					-2.2*	-2.2*
Black	1975-96	2.6*	1996-15	-2.3*									-2.3*	-2.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	1.7*	1997-06	-3.1*	2006-15	-2.2*							-2.2*	-2.2*
White	1992-97	1.7*	1997-06	-3.1*	2006-15	-2.2*							-2.2*	-2.2*
White NH ^{ef}	1992-97	1.9*	1997-06	-3.1*	2006-15	-2.3*							-2.3*	-2.3*
Black	1992-96	2.4	1996-15	-2.2*									-2.2*	-2.2*
Black NH ^{ef}	1992-96	2.8	1996-15	-2.2*									-2.2*	-2.2*
API ^e	1992-15	-1.8*											-1.8*	-1.8*
AI/AN ^{eg}	1992-15	-0.7											-0.7	-0.7
Hispanic ^f	1992-97	3.2*	1997-01	-4.3	2001-15	-1.7*							-1.7*	-1.7*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 19.2
Non-Hodgkin Lymphoma

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-91	4.3*	1991-15	0.3*									0.3*	0.3*
White	1975-90	4.4*	1990-07	0.6*	2007-15	-0.3							-0.2	-0.3
Black	1975-94	5.4*	1994-15	-0.2									-0.2	-0.2
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.1											0.1	0.1
White	1992-95	2.7	1995-98	-2.2	1998-07	0.9*	2007-15	-0.5					-0.4	-0.5
Black	1992-15	-0.1											-0.1	-0.1
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-91	4.2*	1991-08	0.5*	2008-15	-0.7							-0.4	-0.7
White	1975-90	4.4*	1990-07	0.6*	2007-15	-0.6							-0.5	-0.6
Black	1975-94	5.5*	1994-15	-0.4									-0.4	-0.4
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-09	0.3	2009-15	-0.8									-0.5	-0.8
White	1992-95	2.7	1995-98	-2.2	1998-07	0.9*	2007-15	-0.8*					-0.7*	-0.8*
White NH ^{ef}	1992-95	2.9	1995-98	-2.0	1998-07	0.9*	2007-15	-0.7*					-0.5*	-0.7*
Black	1992-15	-0.2											-0.2	-0.2
Black NH ^{ef}	1992-15	-0.1											-0.1	-0.1
API ^e	1992-15	0.3*											0.3*	0.3*
AI/AN ^{eg}	1992-15	0.6											0.6	0.6
Hispanic ^f	1992-15	0.1											0.1	0.1
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-91	2.7*	1991-97	1.6*	1997-06	-2.9*	2006-15	-2.0*					-2.0*	-2.0*
White	1975-91	2.6*	1991-97	1.6*	1997-06	-2.8*	2006-15	-2.0*					-2.0*	-2.0*
Black	1975-96	2.6*	1996-15	-2.4*									-2.4*	-2.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	1.7*	1997-06	-2.9*	2006-15	-2.0*							-2.0*	-2.0*
White	1992-97	1.7*	1997-06	-2.8*	2006-15	-2.0*							-2.0*	-2.0*
White NH ^{ef}	1992-97	2.0*	1997-06	-2.8*	2006-15	-2.0*							-2.0*	-2.0*
Black	1992-96	2.5	1996-15	-2.3*									-2.3*	-2.3*
Black NH ^{ef}	1992-96	2.8	1996-15	-2.2*									-2.2*	-2.2*
API ^e	1992-15	-1.8*											-1.8*	-1.8*
AI/AN ^{eg}	1992-15	-0.4											-0.4	-0.4
Hispanic ^f	1992-96	4.4	1996-03	-3.7*	2003-15	-1.1*							-1.1*	-1.1*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 19.3
Non-Hodgkin Lymphoma

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	2006-15	2011-15								
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-88	3.0*	1988-04	1.4*	2004-15	-0.7*							-0.7*	-0.7*
White	1975-88	3.1*	1988-04	1.4*	2004-15	-0.6*							-0.6*	-0.6*
Black	1975-04	3.0*	2004-15	-1.0									-1.0	-1.0
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	1.3*	2004-15	-0.6*									-0.6*	-0.6*
White	1992-04	1.4*	2004-15	-0.5*									-0.5*	-0.5*
Black	1992-98	5.8*	1998-15	0.2									0.2	0.2
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-88	3.0*	1988-04	1.4*	2004-15	-0.9*							-0.9*	-0.9*
White	1975-88	3.0*	1988-04	1.4*	2004-15	-0.8*							-0.8*	-0.8*
Black	1975-04	3.0*	2004-15	-1.3									-1.3	-1.3
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-04	1.4*	2004-15	-0.8*									-0.8*	-0.8*
White	1992-04	1.4*	2004-15	-0.7*									-0.7*	-0.7*
White NH ^{ef}	1992-04	1.5*	2004-15	-0.9*									-0.9*	-0.9*
Black	1992-04	2.9*	2004-15	-1.0									-1.0	-1.0
Black NH ^{ef}	1992-04	2.9*	2004-15	-0.9									-0.9	-0.9
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-15	1.7											1.7	1.7
Hispanic ^f	1992-15	0.9*											0.9*	0.9*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-95	2.2*	1995-98	-0.4	1998-05	-3.6*	2005-15	-2.7*					-2.7*	-2.7*
White	1975-97	2.1*	1997-04	-3.7*	2004-15	-2.7*							-2.7*	-2.7*
Black	1975-96	2.7*	1996-15	-2.2*									-2.2*	-2.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	1.7*	1997-05	-3.4*	2005-15	-2.7*							-2.7*	-2.7*
White	1992-94	3.3*	1994-97	0.6	1997-00	-2.5	2000-03	-4.4*	2003-15	-2.8*			-2.8*	-2.8*
White NH ^{ef}	1992-94	4.0*	1994-97	0.6	1997-00	-2.4	2000-03	-4.4*	2003-15	-2.8*			-2.8*	-2.8*
Black	1992-97	1.6	1997-15	-2.2*									-2.2*	-2.2*
Black NH ^{ef}	1992-97	1.9	1997-15	-2.2*									-2.2*	-2.2*
API ^e	1992-15	-1.6*											-1.6*	-1.6*
AI/AN ^{eg}	1992-15	-1.1											-1.1	-1.1
Hispanic ^f	1992-97	3.2	1997-15	-2.3*									-2.3*	-2.3*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 19.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Non-Hodgkin Lymphoma, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 19.5

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Non-Hodgkin Lymphoma, Male, by Race




● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero ($p < 0.05$).

Figure 19.6

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Non-Hodgkin Lymphoma, Female, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 20.1
Cancer of the Oral Cavity and Pharynx (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-81	0.7	1981-03	-1.2*	2003-15	0.8*							0.8*	0.8*
White	1975-03	-0.9*	2003-15	1.2*									1.2*	1.2*
Black	1975-84	2.2*	1984-15	-2.3*									-2.3*	-2.3*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	-1.2*	2004-15	0.8*									0.8*	0.8*
White	1992-03	-1.1*	2003-15	1.1*									1.1*	1.1*
Black	1992-15	-2.3*											-2.3*	-2.3*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-81	0.7	1981-03	-1.1*	2003-15	0.7*							0.7*	0.7*
White	1975-03	-0.9*	2003-15	1.1*									1.1*	1.1*
Black	1975-84	2.3*	1984-15	-2.4*									-2.4*	-2.4*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-03	-1.3*	2003-15	0.5*									0.5*	0.5*
White	1992-03	-1.1*	2003-15	1.0*									1.0*	1.0*
White NH ^{ef}	1992-03	-0.8*	2003-15	1.2*									1.2*	1.2*
Black	1992-15	-2.4*											-2.4*	-2.4*
Black NH ^{ef}	1992-15	-2.3*											-2.3*	-2.3*
API ^e	1992-15	-0.7*											-0.7*	-0.7*
AI/AN ^{eg}	1992-15	-0.9											-0.9	-0.9
Hispanic ^f	1992-06	-1.7*	2006-09	6.5	2009-15	-2.2*							0.6	-2.2*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-79	-0.5	1979-93	-1.7*	1993-00	-2.7*	2000-09	-1.3*	2009-15	0.7			0.0	0.7
White	1975-06	-1.8*	2006-15	0.6*									0.6*	0.6*
Black	1975-80	3.3*	1980-93	-1.5*	1993-09	-4.0*	2009-15	-1.2					-2.2*	-1.2
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-00	-2.6*	2000-09	-1.4*	2009-15	0.7							0.0	0.7
White	1992-06	-1.8*	2006-15	0.6*									0.6*	0.6*
White NH ^{ef}	1992-06	-1.6*	2006-15	0.7*									0.7*	0.7*
Black	1992-09	-3.9*	2009-15	-1.3									-2.2*	-1.3
Black NH ^{ef}	1992-09	-3.9*	2009-15	-1.3									-2.1*	-1.3
API ^e	1992-13	-2.1*	2013-15	9.5									0.4	3.5
AI/AN ^{eg}	1992-15	-1.5*											-1.5*	-1.5*
Hispanic ^f	1992-08	-2.7*	2008-15	1.0									0.1	1.0

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 20.3
Cancer of the Oral Cavity and Pharynx (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-05	-1.1*	2005-15	1.3*									1.3*	1.3*
White	1975-03	-1.2*	2003-15	1.5*									1.5*	1.5*
Black	1975-87	1.9*	1987-10	-3.1*	2010-15	0.9							-0.9	0.9
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	-1.4*	2004-15	1.0*									1.0*	1.0*
White	1992-03	-1.3*	2003-15	1.3*									1.3*	1.3*
Black	1992-10	-2.9*	2010-15	0.0									-1.3	0.0
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-83	-0.2	1983-03	-1.4*	2003-15	0.9*							0.9*	0.9*
White	1975-03	-1.2*	2003-15	1.3*									1.3*	1.3*
Black	1975-86	2.1*	1986-15	-2.8*									-2.8*	-2.8*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-03	-1.6*	2003-15	0.7*									0.7*	0.7*
White	1992-03	-1.2*	2003-15	1.2*									1.2*	1.2*
White NH ^{ef}	1992-03	-1.0*	2003-15	1.4*									1.4*	1.4*
Black	1992-15	-2.7*											-2.7*	-2.7*
Black NH ^{ef}	1992-15	-2.6*											-2.6*	-2.6*
API ^e	1992-15	-0.8*											-0.8*	-0.8*
AI/AN ^{eg}	1992-15	-1.7*											-1.7*	-1.7*
Hispanic ^f	1992-15	-0.6											-0.6	-0.6
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-93	-1.9*	1993-00	-3.0*	2000-09	-1.2*	2009-15	1.0*					0.3	1.0*
White	1975-05	-2.1*	2005-15	0.9*									0.9*	0.9*
Black	1975-80	3.7*	1980-91	-1.4*	1991-15	-3.6*							-3.6*	-3.6*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-00	-2.8*	2000-09	-1.3*	2009-15	1.0*							0.3	1.0*
White	1992-05	-2.0*	2005-15	0.8*									0.8*	0.8*
White NH ^{ef}	1992-05	-1.8*	2005-15	0.9*									0.9*	0.9*
Black	1992-15	-3.6*											-3.6*	-3.6*
Black NH ^{ef}	1992-15	-3.6*											-3.6*	-3.6*
API ^e	1992-13	-2.2*	2013-15	12.3									0.9	4.8
AI/AN ^{eg}	1992-15	-1.7*											-1.7*	-1.7*
Hispanic ^f	1992-05	-3.7*	2005-13	-1.1	2013-15	5.5							0.3	2.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 20.4
Cancer of the Oral Cavity and Pharynx (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-80	2.5	1980-07	-0.9*	2007-15	0.9							0.7	0.9
White	1975-81	2.1*	1981-04	-0.9*	2004-15	0.6							0.6	0.6
Black	1975-15	-1.5*											-1.5*	-1.5*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-05	-1.1*	2005-15	0.3									0.3	0.3
White	1992-04	-1.3*	2004-15	0.5									0.5	0.5
Black	1992-15	-1.6*											-1.6*	-1.6*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-80	2.6*	1980-04	-1.0*	2004-15	0.2							0.2	0.2
White	1975-81	2.1*	1981-04	-0.9*	2004-15	0.5							0.5	0.5
Black	1975-15	-1.5*											-1.5*	-1.5*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-04	-1.2*	2004-15	0.0									0.0	0.0
White	1992-04	-1.3*	2004-15	0.3									0.3	0.3
White NH ^{ef}	1992-04	-1.0*	2004-15	0.4									0.4	0.4
Black	1992-15	-1.7*											-1.7*	-1.7*
Black NH ^{ef}	1992-15	-1.6*											-1.6*	-1.6*
API ^e	1992-15	-0.4											-0.4	-0.4
AI/AN ^{eg}	1992-15	0.2											0.2	0.2
Hispanic ^f	1992-15	-0.2											-0.2	-0.2
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-90	-0.9*	1990-06	-2.3*	2006-15	-0.7*							-0.7*	-0.7*
White	1975-89	-1.0*	1989-06	-2.1*	2006-15	-0.6*							-0.6*	-0.6*
Black	1975-91	-0.2	1991-09	-3.7*	2009-15	0.1							-1.2	0.1
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-06	-2.3*	2006-15	-0.7*									-0.7*	-0.7*
White	1992-06	-2.1*	2006-15	-0.6*									-0.6*	-0.6*
White NH ^{ef}	1992-03	-2.2*	2003-15	-0.8*									-0.8*	-0.8*
Black	1992-09	-3.7*	2009-15	0.1									-1.2	0.1
Black NH ^{ef}	1992-09	-3.7*	2009-15	0.2									-1.1	0.2
API ^e	1992-15	-1.6*											-1.6*	-1.6*
AI/AN ^{eg}	1992-15	-1.5											-1.5	-1.5
Hispanic ^f	1992-15	-0.7*											-0.7*	-0.7*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 20.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Oral Cavity and Pharynx, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 20.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Oral Cavity and Pharynx, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 20.5

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Oral Cavity and Pharynx, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 21.1
Cancer of the Ovary (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	2006-15	2011-15										
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-91	-0.1	1991-15	-1.1*									-1.1*	-1.1*
White	1975-90	0.1	1990-15	-1.1*									-1.1*	-1.1*
Black	1975-15	-0.4*											-0.4*	-0.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.0*											-1.0*	-1.0*
White	1992-15	-1.0*											-1.0*	-1.0*
Black	1992-15	-0.5*											-0.5*	-0.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-85	0.1	1985-01	-0.7*	2001-15	-1.5*							-1.5*	-1.5*
White	1975-81	-0.8	1981-84	2.2	1984-01	-0.7*	2001-15	-1.5*					-1.5*	-1.5*
Black	1975-15	-0.5*											-0.5*	-0.5*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-01	-0.6*	2001-15	-1.4*									-1.4*	-1.4*
White	1992-01	-0.4	2001-04	-2.6	2004-09	-0.3	2009-15	-2.1*					-1.5*	-2.1*
White NH ^{ef}	1992-00	-0.4	2000-15	-1.5*									-1.5*	-1.5*
Black	1992-15	-0.6*											-0.6*	-0.6*
Black NH ^{ef}	1992-15	-0.6*											-0.6*	-0.6*
API ^e	1992-15	-0.5*											-0.5*	-0.5*
AI/AN ^{eg}	1992-15	-0.9											-0.9	-0.9
Hispanic ^f	1992-15	-0.5											-0.5	-0.5
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-82	-1.2*	1982-92	0.3*	1992-98	-1.2*	1998-03	0.6	2003-15	-2.3*			-2.3*	-2.3*
White	1975-81	-1.5*	1981-92	0.3*	1992-98	-1.1*	1998-02	0.9	2002-06	-1.4	2006-15	-2.5*	-2.5*	-2.5*
Black	1975-97	0.2	1997-15	-1.3*									-1.3*	-1.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-98	-1.2*	1998-03	0.6	2003-15	-2.3*							-2.3*	-2.3*
White	1992-98	-1.1*	1998-03	0.7	2003-15	-2.3*							-2.3*	-2.3*
White NH ^{ef}	1992-05	-0.2	2005-15	-2.5*									-2.5*	-2.5*
Black	1992-15	-1.1*											-1.1*	-1.1*
Black NH ^{ef}	1992-15	-1.0*											-1.0*	-1.0*
API ^e	1992-15	-0.5*											-0.5*	-0.5*
AI/AN ^{eg}	1992-15	0.0											0.0	0.0
Hispanic ^f	1992-06	-0.1	2006-15	-1.8*									-1.8*	-1.8*

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 21.2
Cancer of the Ovary (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages <65 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-15	-1.1*											-1.1*	-1.1*
White	1975-15	-1.1*											-1.1*	-1.1*
Black	1975-15	-0.7*											-0.7*	-0.7*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.8*											-0.8*	-0.8*
White	1992-15	-0.9*											-0.9*	-0.9*
Black	1992-15	0.0											0.0	0.0
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	-1.1*											-1.1*	-1.1*
White	1975-15	-1.1*											-1.1*	-1.1*
Black	1975-15	-0.7*											-0.7*	-0.7*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.9*											-0.9*	-0.9*
White	1992-15	-1.0*											-1.0*	-1.0*
White NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
Black	1992-15	-0.2											-0.2	-0.2
Black NH ^{ef}	1992-15	-0.1											-0.1	-0.1
API ^e	1992-15	-0.2											-0.2	-0.2
AI/AN ^{eg}	1992-15	-2.5*											-2.5*	-2.5*
Hispanic ^f	1992-15	0.1											0.1	0.1
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-80	-3.2*	1980-99	-1.6*	1999-02	0.5	2002-15	-2.7*					-2.7*	-2.7*
White	1975-80	-3.3*	1980-99	-1.5*	1999-02	0.5	2002-15	-2.8*					-2.8*	-2.8*
Black	1975-96	-2.0*	1996-01	0.9	2001-15	-2.0*							-2.0*	-2.0*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-98	-2.3*	1998-02	0.5	2002-15	-2.7*							-2.7*	-2.7*
White	1992-98	-2.2*	1998-02	0.6	2002-15	-2.8*							-2.8*	-2.8*
White NH ^{ef}	1992-05	-1.1*	2005-15	-3.1*									-3.1*	-3.1*
Black	1992-03	-0.5	2003-15	-2.1*									-2.1*	-2.1*
Black NH ^{ef}	1992-15	-1.3*											-1.3*	-1.3*
API ^e	1992-15	-1.0*											-1.0*	-1.0*
AI/AN ^{eg}	1992-15	-1.3											-1.3	-1.3
Hispanic ^f	1992-15	-0.8*											-0.8*	-0.8*

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

^c Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^e API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

^f Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

^g Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero (p<.05).

- Joinpoint regression line analysis could not be performed on data series.

Table 21.3
Cancer of the Ovary (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program, 1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints, Ages 65+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-91	1.3*	1991-15	-1.2*									-1.2*	-1.2*
White	1975-91	1.4*	1991-15	-1.1*									-1.1*	-1.1*
Black	1975-94	1.3	1994-15	-1.3*									-1.3*	-1.3*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-08	-0.9*	2008-15	-2.6*									-2.2*	-2.6*
White	1992-09	-0.8*	2009-15	-2.6*									-2.0*	-2.6*
Black	1992-15	-0.9*											-0.9*	-0.9*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-90	1.4*	1990-10	-1.0*	2010-15	-3.3*							-2.2*	-3.3*
White	1975-91	1.4*	1991-15	-1.1*									-1.1*	-1.1*
Black	1975-11	0.1	2011-15	-9.4									-4.3	-9.4
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-09	-1.0*	2009-15	-3.2*									-2.4*	-3.2*
White	1992-09	-0.8*	2009-15	-3.0*									-2.3*	-3.0*
White NH ^{ef}	1992-09	-0.8*	2009-15	-2.9*									-2.2*	-2.9*
Black	1992-15	-1.1*											-1.1*	-1.1*
Black NH ^{ef}	1992-15	-1.1*											-1.1*	-1.1*
API ^e	1992-15	-1.0*											-1.0*	-1.0*
AI/AN ^{eg}	1992-15	0.1											0.1	0.1
Hispanic ^f	1992-02	1.3	2002-15	-2.5*									-2.5*	-2.5*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-85	0.7*	1985-91	1.8*	1991-06	-0.1	2006-15	-2.4*					-2.4*	-2.4*
White	1975-84	0.6*	1984-91	1.6*	1991-06	0.0	2006-15	-2.4*					-2.4*	-2.4*
Black	1975-92	2.3*	1992-15	-0.9*									-0.9*	-0.9*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-05	0.0	2005-15	-2.2*									-2.2*	-2.2*
White	1992-06	0.1	2006-15	-2.4*									-2.4*	-2.4*
White NH ^{ef}	1992-05	0.3*	2005-15	-2.2*									-2.2*	-2.2*
Black	1992-15	-1.0*											-1.0*	-1.0*
Black NH ^{ef}	1992-15	-0.9*											-0.9*	-0.9*
API ^e	1992-06	1.8*	2006-15	-2.8*									-2.8*	-2.8*
AI/AN ^{eg}	1992-15	0.5											0.5	0.5
Hispanic ^f	1992-15	-0.7*											-0.7*	-0.7*

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^b Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

^c Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^d The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^e API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

^f Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


^g Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 21.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Ovary, by Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 21.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Ovary, by Age and Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 21.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Ovary, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 22.1
Cancer of the Pancreas (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-78	-2.2	1978-84	1.3	1984-89	-1.5	1989-01	0.0	2001-08	1.7*	2008-15	0.3	0.6*	0.3
White	1975-78	-2.5	1978-84	0.9	1984-95	-0.8*	1995-15	1.1*					1.1*	1.1*
Black	1975-15	-0.1											-0.1	-0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-01	0.0	2001-06	1.8*	2006-15	0.5*							0.5*	0.5*
White	1992-99	0.2	1999-15	1.0*									1.0*	1.0*
Black	1992-01	-1.8*	2001-05	2.7	2005-15	-0.6							-0.6	-0.6
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-78	-2.2	1978-84	1.3	1984-89	-1.5	1989-01	0.0	2001-08	1.7*	2008-15	0.1	0.4	0.1
White	1975-99	-0.3*	1999-07	1.7*	2007-15	0.4							0.5	0.4
Black	1975-15	-0.1											-0.1	-0.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-01	-0.1	2001-06	1.9*	2006-15	0.3							0.3	0.3
White	1992-01	0.2	2001-06	1.9*	2006-15	0.4*							0.4*	0.4*
White NH ^{ef}	1992-00	0.1	2000-06	1.7*	2006-15	0.5*							0.5*	0.5*
Black	1992-01	-1.8*	2001-05	2.8	2005-15	-0.8*							-0.8*	-0.8*
Black NH ^{ef}	1992-01	-1.8*	2001-05	2.8	2005-15	-0.7							-0.7	-0.7
API ^e	1992-15	0.6*											0.6*	0.6*
AI/AN ^{eg}	1992-97	21.1*	1997-15	-1.1									-1.1	-1.1
Hispanic ^f	1992-15	0.7*											0.7*	0.7*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-02	-0.1*	2002-06	0.8	2006-15	0.1							0.1	0.1
White	1975-97	-0.2*	1997-08	0.5*	2008-15	0.2							0.2*	0.2
Black	1975-89	1.1*	1989-15	-0.5*									-0.5*	-0.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	-0.5*	1997-08	0.4*	2008-15	0.1							0.1	0.1
White	1992-96	-0.4	1996-15	0.4*									0.4*	0.4*
White NH ^{ef}	1992-96	-0.2	1996-15	0.4*									0.4*	0.4*
Black	1992-95	-2.5*	1995-15	-0.4*									-0.4*	-0.4*
Black NH ^{ef}	1992-15	-0.4*											-0.4*	-0.4*
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-15	1.2*											1.2*	1.2*
Hispanic ^f	1992-15	0.2*											0.2*	0.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 22.2
Cancer of the Pancreas (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-81	-1.8*	1981-85	1.2	1985-90	-2.2*	1990-03	0.2	2003-06	3.1	2006-15	0.3	0.3	0.3
White	1975-94	-1.0*	1994-15	1.0*									1.0*	1.0*
Black	1975-15	-0.3*											-0.3*	-0.3*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-03	0.0	2003-06	2.9	2006-15	0.3							0.3	0.3
White	1992-15	0.8*											0.8*	0.8*
Black	1992-15	-0.1											-0.1	-0.1
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-93	-0.8*	1993-03	0.2	2003-06	3.1	2006-15	0.1					0.1	0.1
White	1975-94	-1.0*	1994-15	0.9*									0.9*	0.9*
Black	1975-15	-0.3*											-0.3*	-0.3*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-03	0.0	2003-06	3.0	2006-15	0.1							0.1	0.1
White	1992-15	0.8*											0.8*	0.8*
White NH ^{ef}	1992-15	0.8*											0.8*	0.8*
Black	1992-15	-0.2											-0.2	-0.2
Black NH ^{ef}	1992-15	-0.1											-0.1	-0.1
API ^e	1992-15	0.1											0.1	0.1
AI/AN ^{eg}	1992-15	1.4											1.4	1.4
Hispanic ^f	1992-15	0.6*											0.6*	0.6*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-86	-0.8*	1986-00	-0.3*	2000-15	0.3*							0.3*	0.3*
White	1975-86	-1.0*	1986-97	-0.3*	1997-15	0.4*							0.4*	0.4*
Black	1975-89	0.6*	1989-15	-0.7*									-0.7*	-0.7*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-95	-1.4	1995-15	0.2*									0.2*	0.2*
White	1992-96	-0.6	1996-15	0.4*									0.4*	0.4*
White NH ^{ef}	1992-15	0.4*											0.4*	0.4*
Black	1992-95	-3.9*	1995-15	-0.5*									-0.5*	-0.5*
Black NH ^{ef}	1992-01	-1.4*	2001-15	-0.2									-0.2	-0.2
API ^e	1992-15	-0.3											-0.3	-0.3
AI/AN ^{eg}	1992-15	1.7*											1.7*	1.7*
Hispanic ^f	1992-15	0.1											0.1	0.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 22.3
Cancer of the Pancreas (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-84	1.4*	1984-96	-0.5	1996-15	1.0*							1.0*	1.0*
White	1975-84	1.3*	1984-97	-0.5	1997-15	1.2*							1.2*	1.2*
Black	1975-15	0.1											0.1	0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	0.7*											0.7*	0.7*
White	1992-15	0.8*											0.8*	0.8*
Black	1992-15	-0.2											-0.2	-0.2
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-84	1.4*	1984-96	-0.5	1996-15	0.9*							0.9*	0.9*
White	1975-84	1.3*	1984-96	-0.6	1996-15	1.0*							1.0*	1.0*
Black	1975-15	0.1											0.1	0.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	0.7*											0.7*	0.7*
White	1992-15	0.8*											0.8*	0.8*
White NH ^{ef}	1992-15	0.8*											0.8*	0.8*
Black	1992-15	-0.3											-0.3	-0.3
Black NH ^{ef}	1992-15	-0.2											-0.2	-0.2
API ^e	1992-15	1.0*											1.0*	1.0*
AI/AN ^{eg}	1992-15	-0.6											-0.6	-0.6
Hispanic ^f	1992-15	0.7*											0.7*	0.7*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-84	0.8*	1984-03	0.1	2003-06	1.0	2006-15	0.0					0.0	0.0
White	1975-83	0.6*	1983-02	0.1*	2002-08	0.8*	2008-15	-0.1					0.1	-0.1
Black	1975-86	2.5*	1986-15	-0.3*									-0.3*	-0.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	-0.5	1997-08	0.4*	2008-15	-0.1							0.0	-0.1
White	1992-97	-0.4	1997-08	0.6*	2008-15	0.0							0.1	0.0
White NH ^{ef}	1992-02	0.2	2002-08	0.9*	2008-15	-0.1							0.1	-0.1
Black	1992-15	-0.4*											-0.4*	-0.4*
Black NH ^{ef}	1992-15	-0.3*											-0.3*	-0.3*
API ^e	1992-15	0.5*											0.5*	0.5*
AI/AN ^{eg}	1992-15	0.8											0.8	0.8
Hispanic ^f	1992-15	0.3*											0.3*	0.3*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 22.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Pancreas, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 22.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Pancreas, Male, by Race




● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero ($p < 0.05$).

Figure 22.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Pancreas, Female, by Race



Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 23.1
Cancer of the Prostate (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-88	2.6*	1988-92	16.5*	1992-95	-11.5*	1995-00	2.2	2000-09	-1.6*	2009-15	-7.4*	-5.5*	-7.4*
White	1975-88	2.8*	1988-92	16.3*	1992-95	-12.5*	1995-00	2.6	2000-09	-1.8*	2009-15	-7.4*	-5.6*	-7.4*
Black	1975-89	2.1*	1989-92	19.0*	1992-09	-1.5*	2009-15	-6.7*					-5.0*	-6.7*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-95	-11.1*	1995-00	2.0	2000-09	-1.8*	2009-15	-7.5*					-5.6*	-7.5*
White	1992-95	-11.9*	1995-00	2.4	2000-09	-1.9*	2009-15	-7.6*					-5.7*	-7.6*
Black	1992-09	-1.6*	2009-15	-6.8*									-5.1*	-6.8*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-88	2.6*	1988-92	16.5*	1992-95	-11.5*	1995-00	2.2	2000-09	-1.7*	2009-15	-7.7*	-5.7*	-7.7*
White	1975-88	2.8*	1988-92	16.3*	1992-95	-12.5*	1995-00	2.6	2000-09	-1.8*	2009-15	-8.0*	-6.0*	-8.0*
Black	1975-89	2.1*	1989-92	19.2*	1992-09	-1.6*	2009-15	-7.5*					-5.5*	-7.5*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-95	-11.1*	1995-00	2.0	2000-09	-1.8*	2009-15	-7.9*					-5.9*	-7.9*
White	1992-95	-11.9*	1995-00	2.4	2000-09	-1.9*	2009-15	-8.2*					-6.2*	-8.2*
White NH ^{ef}	1992-94	-15.4*	1994-07	-0.2	2007-15	-6.9*							-6.2*	-6.9*
Black	1992-09	-1.7*	2009-15	-7.8*									-5.8*	-7.8*
Black NH ^{ef}	1992-09	-1.6*	2009-15	-7.7*									-5.7*	-7.7*
API ^e	1992-96	-5.4	1996-07	-0.1	2007-15	-8.0*							-7.2*	-8.0*
AI/AN ^{eg}	1992-09	-1.5*	2009-15	-9.2*									-6.7*	-9.2*
Hispanic ^f	1992-95	-4.9	1995-02	0.9	2002-10	-2.8*	2010-15	-9.6*					-6.6*	-9.6*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-87	0.9*	1987-91	3.0*	1991-94	-0.5	1994-98	-4.2*	1998-13	-3.5*	2013-15	-0.8	-2.9*	-2.2*
White	1975-87	0.8*	1987-91	3.1*	1991-94	-0.7	1994-99	-4.3*	1999-13	-3.3*	2013-15	-0.4	-2.7*	-1.9*
Black	1975-88	1.9*	1988-93	3.2*	1993-01	-2.5*	2001-15	-4.2*					-4.2*	-4.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	-0.6	1994-98	-4.2*	1998-13	-3.5*	2013-15	-0.8					-2.9*	-2.2*
White	1992-94	-0.7	1994-99	-4.3*	1999-13	-3.3*	2013-15	-0.4					-2.7*	-1.9*
White NH ^{ef}	1992-94	-0.2	1994-98	-4.5*	1998-12	-3.4*	2012-15	-1.4					-2.7*	-1.9*
Black	1992-01	-2.2*	2001-15	-4.2*									-4.2*	-4.2*
Black NH ^{ef}	1992-01	-2.1*	2001-15	-4.2*									-4.2*	-4.2*
API ^e	1992-15	-3.2*											-3.2*	-3.2*
AI/AN ^{eg}	1992-15	-1.5*											-1.5*	-1.5*
Hispanic ^f	1992-94	4.6	1994-15	-2.8*									-2.8*	-2.8*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 23.2
Cancer of the Prostate (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages <65 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-88	3.1*	1988-92	23.4*	1992-01	3.3*	2001-09	0.0	2009-15	-7.4*			-5.0*	-7.4*
White	1975-88	3.5*	1988-92	23.0*	1992-01	3.1*	2001-09	0.0	2009-15	-7.8*			-5.3*	-7.8*
Black	1975-89	2.0*	1989-93	26.2*	1993-00	2.9*	2000-09	-0.1	2009-15	-5.4*			-3.7*	-5.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-01	3.3*	2001-09	-0.2	2009-15	-7.4*							-5.1*	-7.4*
White	1992-01	3.2*	2001-09	-0.3	2009-15	-8.0*							-5.5*	-8.0*
Black	1992-00	3.8*	2000-10	-0.6	2010-15	-6.7*							-4.0*	-6.7*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-88	3.1*	1988-92	23.4*	1992-01	3.3*	2001-09	0.0	2009-15	-7.6*			-5.2*	-7.6*
White	1975-88	3.5*	1988-92	23.0*	1992-01	3.1*	2001-09	-0.1	2009-15	-8.3*			-5.6*	-8.3*
Black	1975-89	2.0*	1989-93	26.2*	1993-00	2.9*	2000-09	-0.2	2009-15	-6.1*			-4.2*	-6.1*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-01	3.3*	2001-09	-0.2	2009-15	-7.8*							-5.3*	-7.8*
White	1992-01	3.2*	2001-09	-0.3	2009-15	-8.5*							-5.8*	-8.5*
White NH ^{ef}	1992-01	3.2*	2001-09	-0.2	2009-15	-8.3*							-5.7*	-8.3*
Black	1992-00	3.9*	2000-10	-0.8	2010-15	-7.6*							-4.6*	-7.6*
Black NH ^{ef}	1992-00	3.9*	2000-10	-0.7	2010-15	-7.5*							-4.5*	-7.5*
API ^e	1992-01	6.1*	2001-09	0.4	2009-15	-7.6*							-5.0*	-7.6*
AI/AN ^{eg}	1992-06	-0.2	2006-15	-7.6*									-7.6*	-7.6*
Hispanic ^f	1992-02	4.6*	2002-10	-1.5*	2010-15	-8.0*							-5.2*	-8.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-87	0.5*	1987-90	3.0	1990-96	-3.1*	1996-00	-6.0*	2000-15	-1.2*			-1.2*	-1.2*
White	1975-87	0.7*	1987-90	3.8	1990-01	-4.5*	2001-15	-1.1*					-1.1*	-1.1*
Black	1975-92	0.6*	1992-06	-3.1*	2006-15	-2.2*							-2.2*	-2.2*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-01	-4.5*	2001-15	-1.2*									-1.2*	-1.2*
White	1992-01	-4.8*	2001-15	-1.0*									-1.0*	-1.0*
White NH ^{ef}	1992-01	-4.7*	2001-15	-1.1*									-1.1*	-1.1*
Black	1992-04	-3.4*	2004-15	-2.2*									-2.2*	-2.2*
Black NH ^{ef}	1992-04	-3.4*	2004-15	-2.1*									-2.1*	-2.1*
API ^e	1992-15	-0.3											-0.3	-0.3
AI/AN ^{eg}	1992-99	-10.8*	1999-15	0.7									0.7	0.7
Hispanic ^f	1992-15	-1.7*											-1.7*	-1.7*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

a
b
c
d
e
f
g
*
-

Table 23.3
Cancer of the Prostate (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages 65+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-88	2.5*	1988-92	14.8*	1992-95	-13.2*	1995-07	-0.9*	2007-15	-6.7*			-6.0*	-6.7*
White	1975-88	2.7*	1988-92	14.9*	1992-95	-15.7*	1995-00	1.7	2000-09	-2.6*	2009-15	-7.2*	-5.7*	-7.2*
Black	1975-89	2.0*	1989-92	20.3*	1992-95	-9.0*	1995-09	-2.2*	2009-15	-7.9*			-6.0*	-7.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-95	-14.1*	1995-00	0.9	2000-09	-2.5*	2009-15	-7.6*					-5.9*	-7.6*
White	1992-95	-15.0*	1995-00	1.5	2000-10	-2.7*	2010-13	-11.2	2013-15	-0.6			-5.2*	-6.1
Black	1992-95	-8.2*	1995-09	-2.4*	2009-15	-7.7*							-5.9*	-7.7*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-88	2.5*	1988-92	14.8*	1992-95	-13.2*	1995-07	-0.9*	2007-15	-7.0*			-6.3*	-7.0*
White	1975-88	2.7*	1988-92	14.9*	1992-95	-15.7*	1995-00	1.7	2000-09	-2.7*	2009-15	-7.8*	-6.1*	-7.8*
Black	1975-89	2.0*	1989-92	20.3*	1992-95	-8.9*	1995-09	-2.2*	2009-15	-8.7*			-6.6*	-8.7*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-95	-14.1*	1995-00	0.9	2000-09	-2.5*	2009-15	-8.0*					-6.2*	-8.0*
White	1992-94	-18.9*	1994-07	-1.2*	2007-15	-7.2*							-6.5*	-7.2*
White NH ^{ef}	1992-94	-19.5*	1994-07	-1.1*	2007-15	-7.0*							-6.3*	-7.0*
Black	1992-95	-8.0*	1995-09	-2.5*	2009-15	-8.7*							-6.7*	-8.7*
Black NH ^{ef}	1992-95	-8.1*	1995-09	-2.5*	2009-15	-8.7*							-6.6*	-8.7*
API ^e	1992-96	-6.9*	1996-07	-0.9	2007-15	-8.6*							-7.8*	-8.6*
AI/AN ^{eg}	1992-10	-1.8*	2010-15	-10.2*									-6.6*	-10.2*
Hispanic ^f	1992-96	-5.7*	1996-00	1.7	2000-10	-3.0*	2010-15	-10.4*					-7.2*	-10.4*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-87	0.9*	1987-90	3.3*	1990-93	1.3	1993-13	-3.6*	2013-15	-0.9			-3.0*	-2.3*
White	1975-87	0.8*	1987-91	3.2*	1991-94	-0.5	1994-98	-4.4*	1998-13	-3.5*	2013-15	-0.4	-2.8*	-2.0*
Black	1975-94	2.3*	1994-02	-2.8*	2002-15	-4.4*							-4.4*	-4.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	-1.2	1994-13	-3.7*	2013-15	-0.8							-3.0*	-2.2*
White	1992-94	-0.6	1994-98	-4.3*	1998-13	-3.5*	2013-15	-0.4					-2.8*	-2.0*
White NH ^{ef}	1992-94	-0.2	1994-98	-4.3*	1998-12	-3.5*	2012-15	-1.5					-2.9*	-2.0*
Black	1992-01	-2.1*	2001-15	-4.4*									-4.4*	-4.4*
Black NH ^{ef}	1992-01	-2.0*	2001-15	-4.3*									-4.3*	-4.3*
API ^e	1992-15	-3.3*											-3.3*	-3.3*
AI/AN ^{eg}	1992-15	-1.5*											-1.5*	-1.5*
Hispanic ^f	1992-95	2.2	1995-15	-2.9*									-2.9*	-2.9*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 23.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Prostate, by Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 23.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Prostate, by Age and Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.


The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 23.4

Cancer of the Prostate


Delay-Adjusted SEER Incidence & US Mortality 1975-2015



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

Figure 23.5


Delay-Adjusted Incidence Rates for Cancer of the Prostate and Cancer of the Lung and Bronchus All Races, Males 1975-2015



Source: SEER 9 areas. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

Figure 23.6

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Prostate, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero (p < 0.05).

Table 24.1
Cancer of the Stomach (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-15	-1.5*											-1.5*	-1.5*
White	1975-97	-2.0*	1997-15	-1.2*									-1.2*	-1.2*
Black	1975-82	1.0	1982-15	-1.6*									-1.6*	-1.6*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.2*											-1.2*	-1.2*
White	1992-15	-1.1*											-1.1*	-1.1*
Black	1992-15	-1.7*											-1.7*	-1.7*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-15	-1.5*											-1.5*	-1.5*
White	1975-97	-2.0*	1997-15	-1.3*									-1.3*	-1.3*
Black	1975-82	1.1	1982-15	-1.6*									-1.6*	-1.6*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.3*											-1.3*	-1.3*
White	1992-15	-1.1*											-1.1*	-1.1*
White NH ^{ef}	1992-15	-1.5*											-1.5*	-1.5*
Black	1992-15	-1.8*											-1.8*	-1.8*
Black NH ^{ef}	1992-15	-1.7*											-1.7*	-1.7*
API ^e	1992-15	-3.0*											-3.0*	-3.0*
AI/AN ^{eg}	1992-15	-0.4											-0.4	-0.4
Hispanic ^f	1992-15	-1.5*											-1.5*	-1.5*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-87	-2.5*	1987-90	-0.3	1990-09	-3.0*	2009-15	-1.9*					-2.3*	-1.9*
White	1975-87	-2.8*	1987-90	-0.8	1990-08	-3.2*	2008-15	-1.8*					-2.1*	-1.8*
Black	1975-95	-1.3*	1995-15	-3.3*									-3.3*	-3.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-09	-3.0*	2009-15	-1.9*									-2.3*	-1.9*
White	1992-08	-3.2*	2008-15	-1.9*									-2.2*	-1.9*
White NH ^{ef}	1992-94	-1.2	1994-08	-3.7*	2008-15	-2.5*							-2.7*	-2.5*
Black	1992-95	-1.2	1995-15	-3.3*									-3.3*	-3.3*
Black NH ^{ef}	1992-95	-0.8	1995-15	-3.2*									-3.2*	-3.2*
API ^e	1992-15	-3.7*											-3.7*	-3.7*
AI/AN ^{eg}	1992-15	-2.1*											-2.1*	-2.1*
Hispanic ^f	1992-15	-2.2*											-2.2*	-2.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 24.2
Cancer of the Stomach (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-80	-0.2	1980-15	-1.8*									-1.8*	-1.8*
White	1975-15	-1.8*											-1.8*	-1.8*
Black	1975-82	2.4	1982-15	-2.1*									-2.1*	-2.1*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.6*											-1.6*	-1.6*
White	1992-15	-1.5*											-1.5*	-1.5*
Black	1992-15	-2.1*											-2.1*	-2.1*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-80	-0.2	1980-15	-1.8*									-1.8*	-1.8*
White	1975-15	-1.8*											-1.8*	-1.8*
Black	1975-82	2.5	1982-15	-2.1*									-2.1*	-2.1*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-1.7*											-1.7*	-1.7*
White	1992-15	-1.6*											-1.6*	-1.6*
White NH ^{ef}	1992-15	-1.8*											-1.8*	-1.8*
Black	1992-15	-2.2*											-2.2*	-2.2*
Black NH ^{ef}	1992-15	-2.1*											-2.1*	-2.1*
API ^e	1992-15	-3.0*											-3.0*	-3.0*
AI/AN ^{eg}	1992-15	-0.5											-0.5	-0.5
Hispanic ^f	1992-15	-2.0*											-2.0*	-2.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-87	-2.3*	1987-91	-1.0	1991-08	-3.5*	2008-15	-2.3*					-2.5*	-2.3*
White	1975-88	-2.5*	1988-91	-0.9	1991-08	-3.7*	2008-15	-2.2*					-2.5*	-2.2*
Black	1975-93	-1.1*	1993-15	-3.3*									-3.3*	-3.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	-1.4	1994-06	-3.6*	2006-15	-2.5*							-2.5*	-2.5*
White	1992-09	-3.6*	2009-15	-2.0*									-2.5*	-2.0*
White NH ^{ef}	1992-94	-0.9	1994-08	-4.1*	2008-15	-2.6*							-3.0*	-2.6*
Black	1992-15	-3.2*											-3.2*	-3.2*
Black NH ^{ef}	1992-15	-3.1*											-3.1*	-3.1*
API ^e	1992-15	-3.8*											-3.8*	-3.8*
AI/AN ^{eg}	1992-15	-2.0*											-2.0*	-2.0*
Hispanic ^f	1992-15	-2.5*											-2.5*	-2.5*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 24.3
Cancer of the Stomach (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-95	-1.8*	1995-15	-0.9*									-0.9*	-0.9*
White	1975-95	-2.3*	1995-15	-1.1*									-1.1*	-1.1*
Black	1975-15	-0.9*											-0.9*	-0.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.8*											-0.8*	-0.8*
White	1992-15	-0.7*											-0.7*	-0.7*
Black	1992-15	-1.1*											-1.1*	-1.1*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-95	-1.8*	1995-15	-1.0*									-1.0*	-1.0*
White	1975-95	-2.3*	1995-15	-1.1*									-1.1*	-1.1*
Black	1975-15	-0.9*											-0.9*	-0.9*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	-0.9*											-0.9*	-0.9*
White	1992-15	-0.8*											-0.8*	-0.8*
White NH ^{ef}	1992-15	-1.4*											-1.4*	-1.4*
Black	1992-15	-1.2*											-1.2*	-1.2*
Black NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
API ^e	1992-15	-2.9*											-2.9*	-2.9*
AI/AN ^{eg}	1992-15	-0.1											-0.1	-0.1
Hispanic ^f	1992-15	-1.0*											-1.0*	-1.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-87	-2.8*	1987-90	-0.3	1990-09	-2.7*	2009-15	-1.8*					-2.1*	-1.8*
White	1975-87	-3.1*	1987-90	-1.0	1990-08	-2.9*	2008-15	-1.6*					-1.9*	-1.6*
Black	1975-98	-1.2*	1998-15	-3.5*									-3.5*	-3.5*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-09	-2.7*	2009-15	-1.8*									-2.1*	-1.8*
White	1992-09	-2.8*	2009-15	-1.5*									-1.9*	-1.5*
White NH ^{ef}	1992-15	-3.1*											-3.1*	-3.1*
Black	1992-15	-3.1*											-3.1*	-3.1*
Black NH ^{ef}	1992-00	-2.0*	2000-15	-3.5*									-3.5*	-3.5*
API ^e	1992-15	-3.6*											-3.6*	-3.6*
AI/AN ^{eg}	1992-15	-2.4*											-2.4*	-2.4*
Hispanic ^f	1992-15	-2.0*											-2.0*	-2.0*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 24.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Stomach, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 24.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Stomach, Male, by Race




● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero (p < 0.05).

Figure 24.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Stomach, Female, by Race



● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero (p < 0.05).

Table 25.1
Cancer of the Testis (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
All Ages by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-89	2.3*	1989-15	0.8*									0.8*	0.8*
White	1975-89	2.7*	1989-15	1.0*									1.0*	1.0*
Black	1975-15	2.0*											2.0*	2.0*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	1.1*											1.1*	1.1*
White	1992-15	1.2*											1.2*	1.2*
Black	1992-94	47.3	1994-15	0.9									0.9	0.9
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-89	2.4*	1989-15	0.8*									0.8*	0.8*
White	1975-89	2.7*	1989-15	1.0*									1.0*	1.0*
Black	1975-15	2.0*											2.0*	2.0*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	1.0*											1.0*	1.0*
White	1992-15	1.2*											1.2*	1.2*
White NH ^{ef}	1992-05	1.8*	2005-15	-0.2									-0.2	-0.2
Black	1992-94	47.6	1994-15	0.8									0.8	0.8
Black NH ^{ef}	1992-94	44.3	1994-15	1.0									1.0	1.0
API ^e	1992-15	1.6*											1.6*	1.6*
AI/AN ^{eg}	1992-15	3.1*											3.1*	3.1*
Hispanic ^f	1992-15	3.0*											3.0*	3.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-81	-10.5*	1981-95	-3.1*	1995-15	-0.2							-0.2	-0.2
White	1975-81	-10.8*	1981-95	-3.0*	1995-15	0.0							0.0	0.0
Black	1975-15	-1.8*											-1.8*	-1.8*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-0.5*											-0.5*	-0.5*
White	1992-15	-0.3											-0.3	-0.3
White NH ^{ef}	1992-15	-0.5*											-0.5*	-0.5*
Black	1992-15	-0.9											-0.9	-0.9
Black NH ^{ef}	1992-15	-1.0											-1.0	-1.0
API ^e	1992-15	-0.8											-0.8	-0.8
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	0.6											0.6	0.6

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 25.2
Cancer of the Testis (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages <50 by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-89	2.8*	1989-15	0.7*									0.7*	0.7*
White	1975-89	3.2*	1989-15	0.9*									0.9*	0.9*
Black	1975-15	1.9*											1.9*	1.9*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	1.0*											1.0*	1.0*
White	1992-15	1.2*											1.2*	1.2*
Black	1992-15	1.3*											1.3*	1.3*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-89	2.8*	1989-15	0.7*									0.7*	0.7*
White	1975-89	3.2*	1989-15	0.9*									0.9*	0.9*
Black	1975-15	1.8*											1.8*	1.8*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	1.0*											1.0*	1.0*
White	1992-15	1.1*											1.1*	1.1*
White NH ^{ef}	1992-05	1.8*	2005-15	-0.5									-0.5	-0.5
Black	1992-15	1.3*											1.3*	1.3*
Black NH ^{ef}	1992-15	1.4*											1.4*	1.4*
API ^e	1992-15	1.5*											1.5*	1.5*
AI/AN ^{eg}	1992-15	3.0*											3.0*	3.0*
Hispanic ^f	1992-15	3.0*											3.0*	3.0*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-81	-11.3*	1981-95	-2.8*	1995-15	-0.6*							-0.6*	-0.6*
White	1975-81	-11.5*	1981-95	-2.7*	1995-15	-0.4							-0.4	-0.4
Black	1975-15	-1.4*											-1.4*	-1.4*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-0.8*											-0.8*	-0.8*
White	1992-15	-0.6*											-0.6*	-0.6*
White NH ^{ef}	1992-15	-1.1*											-1.1*	-1.1*
Black	1992-15	-1.6*											-1.6*	-1.6*
Black NH ^{ef}	1992-15	-1.7*											-1.7*	-1.7*
API ^e	1992-15	0.7											0.7	0.7
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	1.4*											1.4*	1.4*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 25.3
Cancer of the Testis (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Ages 50+ by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-99	-0.3	1999-15	2.2*									2.2*	2.2*
White	1975-98	-0.3	1998-15	2.2*									2.2*	2.2*
Black	-	-											-	-
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-15	1.6*											1.6*	1.6*
White	1992-15	1.7*											1.7*	1.7*
Black	-	-											-	-
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-99	-0.3	1999-15	2.2*									2.2*	2.2*
White	1975-98	-0.3	1998-15	2.1*									2.1*	2.1*
Black	-	-											-	-
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-15	1.6*											1.6*	1.6*
White	1992-15	1.6*											1.6*	1.6*
White NH ^{ef}	1992-15	1.8*											1.8*	1.8*
Black	-	-											-	-
Black NH ^{ef}	-	-											-	-
API ^e	1992-15	1.8											1.8	1.8
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	1.9											1.9	1.9
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-91	-5.8*	1991-15	0.4									0.4	0.4
White	1975-91	-5.8*	1991-15	0.6									0.6	0.6
Black	1975-15	-2.3*											-2.3*	-2.3*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	0.4											0.4	0.4
White	1992-15	0.6											0.6	0.6
White NH ^{ef}	1992-15	0.8*											0.8*	0.8*
Black	1992-15	0.0											0.0	0.0
Black NH ^{ef}	1992-15	-0.2											-0.2	-0.2
API ^e	-	-											-	-
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-1.8											-1.8	-1.8

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 25.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Testis, by Race



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 25.2

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Testis, by Age and Race



Regression line cannot be calculated for Black Male Age 50+ Incidence and Delay-adjusted Incidence due to years with zero count.

Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 25.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Testis, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 26.1
Cancer of the Thyroid (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-77	6.3	1977-80	-6.5	1980-97	2.4*	1997-09	6.7*	2009-15	1.5*			3.2*	1.5*
White	1975-77	7.5	1977-80	-6.7	1980-97	2.7*	1997-09	7.0*	2009-15	1.4*			3.2*	1.4*
Black	1975-95	0.8	1995-12	6.3*	2012-15	-1.9							3.5*	0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-98	2.9*	1998-09	6.6*	2009-15	1.6*							3.3*	1.6*
White	1992-98	3.6*	1998-09	6.8*	2009-15	1.5*							3.2*	1.5*
Black	1992-95	-4.4	1995-11	6.8*	2011-15	-0.5							3.5*	-0.5
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-77	6.3	1977-80	-6.5	1980-97	2.4*	1997-09	6.7*	2009-15	1.3*			3.1*	1.3*
White	1975-77	7.5	1977-80	-6.7	1980-97	2.7*	1997-09	7.0*	2009-15	1.2*			3.1*	1.2*
Black	1975-95	0.9	1995-12	6.2*	2012-15	-2.6							3.2*	-0.4
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-98	2.9*	1998-09	6.6*	2009-15	1.5*							3.1*	1.5*
White	1992-98	3.6*	1998-09	6.8*	2009-15	1.3*							3.1*	1.3*
White NH ^{ef}	1992-98	4.4*	1998-09	7.0*	2009-15	1.1*							3.0*	1.1*
Black	1992-95	-4.2	1995-11	6.7*	2011-15	-1.0							3.2*	-1.0
Black NH ^{ef}	1992-95	-4.6	1995-11	6.9*	2011-15	-0.9							3.3*	-0.9
API ^e	1992-04	1.9*	2004-09	7.6*	2009-15	1.0							3.1*	1.0
AI/AN ^{eg}	1992-15	3.7*											3.7*	3.7*
Hispanic ^f	1992-95	-5.3	1995-13	5.0*	2013-15	-1.7							3.5*	1.6
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-88	-2.1*	1988-15	0.7*									0.7*	0.7*
White	1975-88	-2.2*	1988-15	0.7*									0.7*	0.7*
Black	1975-15	0.1											0.1	0.1
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	0.7*											0.7*	0.7*
White	1992-15	0.7*											0.7*	0.7*
White NH ^{ef}	1992-15	0.7*											0.7*	0.7*
Black	1992-15	0.9*											0.9*	0.9*
Black NH ^{ef}	1992-15	1.0*											1.0*	1.0*
API ^e	1992-15	-0.5											-0.5	-0.5
AI/AN ^{eg}	1992-15	0.8											0.8	0.8
Hispanic ^f	1992-15	0.1											0.1	0.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 26.2
Cancer of the Thyroid (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-80	-4.6	1980-97	1.8*	1997-12	5.5*	2012-15	-1.1					3.2*	0.5
White	1975-77	10.4	1977-80	-9.1	1980-98	2.6*	1998-09	6.2*	2009-15	2.3*			3.6*	2.3*
Black	1975-15	3.0*											3.0*	3.0*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-96	-1.4	1996-09	5.6*	2009-15	2.5*							3.6*	2.5*
White	1992-96	-0.3	1996-09	5.8*	2009-15	2.3*							3.5*	2.3*
Black	1992-15	3.8*											3.8*	3.8*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-80	-4.6	1980-97	1.8*	1997-12	5.4*	2012-15	-1.4					3.1*	0.3
White	1975-77	10.4	1977-80	-9.1	1980-98	2.6*	1998-09	6.2*	2009-15	2.1*			3.4*	2.1*
Black	1975-15	3.0*											3.0*	3.0*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-96	-1.3	1996-09	5.6*	2009-15	2.4*							3.4*	2.4*
White	1992-96	-0.3	1996-09	5.8*	2009-15	2.1*							3.3*	2.1*
White NH ^{ef}	1992-96	0.1	1996-09	6.2*	2009-15	2.1*							3.4*	2.1*
Black	1992-15	3.7*											3.7*	3.7*
Black NH ^{ef}	1992-15	3.7*											3.7*	3.7*
API ^e	1992-03	0.6	2003-15	5.5*									5.5*	5.5*
AI/AN ^{eg}	1992-15	2.5											2.5	2.5
Hispanic ^f	1992-15	3.9*											3.9*	3.9*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-83	-3.1*	1983-15	1.2*									1.2*	1.2*
White	1975-84	-2.5*	1984-15	1.2*									1.2*	1.2*
Black	1975-15	0.4											0.4	0.4
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	1.3*											1.3*	1.3*
White	1992-15	1.4*											1.4*	1.4*
White NH ^{ef}	1992-15	1.4*											1.4*	1.4*
Black	1992-15	1.3*											1.3*	1.3*
Black NH ^{ef}	1992-15	1.4*											1.4*	1.4*
API ^e	1992-15	0.3											0.3	0.3
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	1.2*											1.2*	1.2*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 26.3
Cancer of the Thyroid (Invasive)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-77	6.5	1977-80	-5.2	1980-93	2.3*	1993-99	4.4*	1999-09	7.1*	2009-15	1.2*	3.1*	1.2*
White	1975-81	-1.1	1981-93	2.4*	1993-99	5.1*	1999-09	7.4*	2009-15	1.1*			3.2*	1.1*
Black	1975-89	-1.1	1989-15	5.5*									5.5*	5.5*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-99	4.1*	1999-09	6.9*	2009-15	1.4*							3.2*	1.4*
White	1992-98	4.4*	1998-09	7.0*	2009-15	1.4*							3.2*	1.4*
Black	1992-98	2.9	1998-11	7.5*	2011-15	-0.6							3.8*	-0.6
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-77	6.5	1977-80	-5.2	1980-93	2.3*	1993-99	4.5*	1999-09	7.1*	2009-15	1.0*	3.0*	1.0*
White	1975-81	-1.1	1981-93	2.4*	1993-99	5.2*	1999-09	7.4*	2009-15	0.9			3.0*	0.9
Black	1975-93	0.2	1993-12	6.3*	2012-15	-1.4							3.7*	0.5
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-99	4.1*	1999-09	6.9*	2009-15	1.2*							3.1*	1.2*
White	1992-98	4.4*	1998-09	7.0*	2009-15	1.2*							3.1*	1.2*
White NH ^{ef}	1992-99	5.6*	1999-09	7.3*	2009-15	0.8							2.9*	0.8
Black	1992-98	3.0	1998-11	7.4*	2011-15	-1.1							3.5*	-1.1
Black NH ^{ef}	1992-98	2.8	1998-11	7.6*	2011-15	-1.1							3.6*	-1.1
API ^e	1992-99	0.3	1999-10	5.3*	2010-15	0.0							2.3*	0.0
AI/AN ^{eg}	1992-15	4.0*											4.0*	4.0*
Hispanic ^f	1992-95	-4.2	1995-13	5.1*	2013-15	-1.3							3.7*	1.9
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-88	-2.5*	1988-15	0.3*									0.3*	0.3*
White	1975-88	-2.8*	1988-15	0.3*									0.3*	0.3*
Black	1975-15	0.0											0.0	0.0
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-94	-4.9	1994-15	0.5*									0.5*	0.5*
White	1992-15	0.2											0.2	0.2
White NH ^{ef}	1992-15	0.2											0.2	0.2
Black	1992-15	0.8*											0.8*	0.8*
Black NH ^{ef}	1992-15	0.9*											0.9*	0.9*
API ^e	1992-15	-1.0											-1.0	-1.0
AI/AN ^{eg}	1992-15	-1.2											-1.2	-1.2
Hispanic ^f	1992-15	-0.5											-0.5	-0.5

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 26.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Thyroid, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 26.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Thyroid, Male, by Race




● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero ($p < 0.05$).

Figure 26.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Thyroid, Female, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Table 27.1
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Both Sexes by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-86	0.7*	1986-07	0.2*	2007-15	-0.9*							-0.8*	-0.9*
White	1975-86	0.9*	1986-07	0.3*	2007-15	-0.8*							-0.7*	-0.8*
Black	1975-15	0.3*											0.3*	0.3*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-07	0.1	2007-15	-1.2*									-1.1*	-1.2*
White	1992-07	0.2*	2007-15	-1.1*									-1.0*	-1.1*
Black	1992-15	0.3											0.3	0.3
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-86	0.7*	1986-07	0.2*	2007-15	-1.1*							-1.0*	-1.1*
White	1975-86	0.9*	1986-07	0.3*	2007-15	-1.1*							-0.9*	-1.1*
Black	1975-15	0.2*											0.2*	0.2*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-07	0.1	2007-15	-1.5*									-1.3*	-1.5*
White	1992-07	0.3*	2007-15	-1.4*									-1.2*	-1.4*
White NH ^{ef}	1992-07	0.4*	2007-15	-1.2*									-1.0*	-1.2*
Black	1992-15	0.1											0.1	0.1
Black NH ^{ef}	1992-15	0.1											0.1	0.1
API ^e	1992-08	0.4	2008-15	-2.3*									-1.7*	-2.3*
AI/AN ^{eg}	1992-15	1.4											1.4	1.4
Hispanic ^f	1992-15	-0.4*											-0.4*	-0.4*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-77	0.2	1977-87	-2.2*	1987-15	0.0*							0.0*	0.0*
White	1975-77	0.3	1977-87	-2.2*	1987-15	0.1*							0.1*	0.1*
Black	1975-15	-1.0*											-1.0*	-1.0*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	0.0											0.0	0.0
White	1992-15	0.1*											0.1*	0.1*
White NH ^{ef}	1992-15	0.3*											0.3*	0.3*
Black	1992-15	-0.9*											-0.9*	-0.9*
Black NH ^{ef}	1992-15	-0.8*											-0.8*	-0.8*
API ^e	1992-15	-0.5											-0.5	-0.5
AI/AN ^{eg}	1992-15	2.3*											2.3*	2.3*
Hispanic ^f	1992-15	-0.2											-0.2	-0.2

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 27.2
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Males by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-86	0.8*	1986-07	0.1	2007-15	-1.0*							-0.9*	-1.0*
White	1975-86	1.0*	1986-07	0.2	2007-15	-0.9*							-0.8*	-0.9*
Black	1975-15	0.4*											0.4*	0.4*
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-07	0.1	2007-15	-1.4*									-1.2*	-1.4*
White	1992-07	0.2	2007-15	-1.3*									-1.1*	-1.3*
Black	1992-15	0.5*											0.5*	0.5*
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-86	0.8*	1986-07	0.1	2007-15	-1.2*							-1.1*	-1.2*
White	1975-87	1.2*	1987-94	-0.4	1994-07	0.4*	2007-15	-1.4*					-1.2*	-1.4*
Black	1975-15	0.4*											0.4*	0.4*
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-07	0.1	2007-15	-1.6*									-1.4*	-1.6*
White	1992-07	0.2	2007-15	-1.5*									-1.4*	-1.5*
White NH ^{ef}	1992-07	0.3*	2007-15	-1.3*									-1.2*	-1.3*
Black	1992-15	0.4											0.4	0.4
Black NH ^{ef}	1992-15	0.4											0.4	0.4
API ^e	1992-07	0.9*	2007-15	-2.5*									-2.1*	-2.5*
AI/AN ^{eg}	1992-15	1.1											1.1	1.1
Hispanic ^f	1992-15	-0.4											-0.4	-0.4
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-83	-1.4*	1983-87	-2.8*	1987-93	0.2	1993-97	-1.1	1997-13	0.0	2013-15	-1.4	-0.3	-0.7
White	1975-83	-1.4*	1983-87	-2.8*	1987-93	0.3	1993-97	-0.9	1997-13	0.1*	2013-15	-1.3	-0.2	-0.6
Black	1975-15	-0.9*											-0.9*	-0.9*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-97	-0.8	1997-15	-0.1									-0.1	-0.1
White	1992-15	0.0											0.0	0.0
White NH ^{ef}	1992-02	-0.3*	2002-05	1.2	2005-15	0.0							0.0	0.0
Black	1992-15	-0.7*											-0.7*	-0.7*
Black NH ^{ef}	1992-15	-0.6*											-0.6*	-0.6*
API ^e	1992-15	0.0											0.0	0.0
AI/AN ^{eg}	1992-15	2.7*											2.7*	2.7*
Hispanic ^f	1992-15	-0.4*											-0.4*	-0.4*

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska

Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.

Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Table 27.3
Cancer of the Urinary Bladder (Invasive and *In Situ*)

Trends in SEER Incidence^{ab} and U.S. Mortality^c Using the Joinpoint Regression Program,
1975-2015 With up to Five Joinpoints, 1992-2015 With up to Four Joinpoints,
Females by Race/Ethnicity

	JP Trend 1		JP Trend 2		JP Trend 3		JP Trend 4		JP Trend 5		JP Trend 6		AAPC ^d	
	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	Years	APC	2006-15	2011-15
<u>SEER 9 Delay-Adjusted Incidence^a, 1975-2015</u>														
All Races	1975-02	0.2*	2002-15	-1.0*									-1.0*	-1.0*
White	1975-02	0.4*	2002-15	-0.9*									-0.9*	-0.9*
Black	1975-15	0.1											0.1	0.1
<u>SEER 13 Delay-Adjusted Incidence^b, 1992-2015</u>														
All Races	1992-04	-0.2	2004-15	-1.3*									-1.3*	-1.3*
White	1992-04	0.0	2004-15	-1.2*									-1.2*	-1.2*
Black	1992-15	-0.3											-0.3	-0.3
<u>SEER 9 Observed Incidence^a, 1975-2015</u>														
All Races	1975-03	0.2*	2003-15	-1.2*									-1.2*	-1.2*
White	1975-02	0.4*	2002-15	-1.0*									-1.0*	-1.0*
Black	1975-15	0.1											0.1	0.1
<u>SEER 13 Observed Incidence^b, 1992-2015</u>														
All Races	1992-04	-0.1	2004-15	-1.4*									-1.4*	-1.4*
White	1992-04	0.0	2004-15	-1.4*									-1.4*	-1.4*
White NH ^{ef}	1992-04	0.2	2004-15	-1.2*									-1.2*	-1.2*
Black	1992-15	-0.4											-0.4	-0.4
Black NH ^{ef}	1992-15	-0.4											-0.4	-0.4
API ^e	1992-15	-0.2											-0.2	-0.2
AI/AN ^{eg}	1992-15	4.3*											4.3*	4.3*
Hispanic ^f	1992-15	-0.7*											-0.7*	-0.7*
<u>U.S. Cancer Mortality^c, 1975-2015</u>														
All Races	1975-86	-1.6*	1986-15	-0.4*									-0.4*	-0.4*
White	1975-86	-1.7*	1986-15	-0.3*									-0.3*	-0.3*
Black	1975-02	-0.8*	2002-15	-1.7*									-1.7*	-1.7*
<u>U.S. Cancer Mortality^c, 1992-2015</u>														
All Races	1992-15	-0.5*											-0.5*	-0.5*
White	1992-15	-0.3*											-0.3*	-0.3*
White NH ^{ef}	1992-15	-0.2*											-0.2*	-0.2*
Black	1992-15	-1.3*											-1.3*	-1.3*
Black NH ^{ef}	1992-15	-1.2*											-1.2*	-1.2*
API ^e	1992-15	-1.1*											-1.1*	-1.1*
AI/AN ^{eg}	-	-											-	-
Hispanic ^f	1992-15	-0.1											-0.1	-0.1

Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute. (<https://surveillance.cancer.gov/joinpoint/>).

^a The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
^b Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

^c Trends are from the SEER 13 areas (SEER 9 Areas, Los Angeles, San Jose-Monterey, Rural Georgia, and the Alaska Native Registry).

^d Trends are from US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^e The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.

^f API - Asian/Pacific Islander, AI/AN - American Indian/Alaska Native, NH - Non-Hispanic

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

The Hispanic and Non-Hispanic mortality trends exclude deaths from New Hampshire and Oklahoma.


^g Data for American Indian/Alaska Native are based on the CHSDA (Contract Health Service Delivery Area) counties.

* The APC/AAPC is significantly different from zero ($p < .05$).

- Joinpoint regression line analysis could not be performed on data series.

Figure 27.1

SEER Observed Incidence, SEER Delay Adjusted Incidence and US Death Rates^a Cancer of the Urinary Bladder, by Race and Sex



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 27.3

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Urinary Bladder, Male, by Race



● SEER 9 Delay-Adj. Incidence ○ SEER 9 Observed Incidence ■ SEER 13 Delay-Adj. Incidence □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).


Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

Figure 27.4

SEER Observed Incidence and SEER Delay Adjusted Incidence Rates^a SEER 9 Areas Compared to SEER 13 Areas Cancer of the Urinary Bladder, Female, by Race




● SEER 9 Delay-Adj. Incidence
 ○ SEER 9 Observed Incidence
 ■ SEER 13 Delay-Adj. Incidence
 □ SEER 13 Observed Incidence

Source: SEER 9 areas and SEER 13 areas.
^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
 Regression lines and APCs are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.
 The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.
 * The APC is significantly different from zero ($p < 0.05$).


Figure 28.1

SEER Delay-Adjusted Incidence and US Mortality All Childhood Cancers, Under 20 Years of Age Both Sexes, All Races, 1975-2015



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC).
Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).
Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018,
National Cancer Institute.

SEER Delay-Adjusted Incidence and US Mortality Brain and CNS, Under 15 Years of Age Both Sexes, All Races, 1975-2015



Source: SEER 9 areas and US Mortality Files (National Center for Health Statistics, CDC). Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines are calculated using the Joinpoint Regression Program Version 4.6, February 2018, National Cancer Institute.