

Table 28.1

Age-Adjusted SEER Cancer Incidence^a and U.S. Death^b Rates, 2011-2015

By Primary Cancer Site

All Races, Males and Females

Site	Ages 0-14						Ages 0-19					
	Incidence			Mortality			Incidence			Mortality		
	Total	Males	Females	Total	Males	Females	Total	Males	Females	Total	Males	Females
All Sites												
All Races	16.4	17.0	15.9	2.1	2.3	1.9	18.2	18.7	17.6	2.3	2.5	2.1
Whites	17.3	17.9	16.6	2.2	2.3	2.0	19.3	19.9	18.7	2.4	2.6	2.1
Blacks	13.1	13.0	13.2	2.0	2.1	1.9	13.7	13.8	13.6	2.2	2.4	2.0
Bone & Joint	0.8	0.8	0.8	0.1	0.1	0.1	1.0	1.1	0.9	0.2	0.3	0.2
Brain & Other nervous	3.4	3.5	3.2	0.7	0.8	0.7	3.1	3.3	3.0	0.7	0.7	0.6
Hodgkin lymphoma	0.5	0.7	0.4	-	-	-	1.2	1.3	1.1	0.0	0.0	0.0
Kidney & Renal pelvis	0.9	0.8	1.0	0.1	0.1	0.1	0.7	0.7	0.8	0.1	0.0	0.1
Leukemia	5.3	5.5	5.0	0.6	0.6	0.5	4.7	5.1	4.4	0.6	0.7	0.5
Acute lymphocytic	4.2	4.3	4.0	0.2	0.3	0.2	3.6	3.8	3.3	0.3	0.3	0.2
Non-Hodgkin lymphoma	1.1	1.4	0.7	0.1	0.1	0.0	1.3	1.7	0.9	0.1	0.1	0.1
Soft tissue	1.1	1.0	1.2	0.2	0.2	0.1	1.1	1.1	1.2	0.2	0.2	0.2

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

- Statistic could not be calculated. Rate based on less than 16 cases for the time interval.