

Table 5.7
Cancer of the Cervix Uteri (Invasive)

SEER Incidence^a and U.S. Death^b Rates, Age-Adjusted and Age-Specific Rates, by Race

	SEER Incidence ^a			U.S. Mortality ^b		
	All Races Females	White Females	Black Females	All Races Females	White Females	Black Females
<u>Age-Adjusted Rates, 2012-2016</u>						
Age at Diagnosis/Death:						
All ages	7.3	7.2	8.7	2.3	2.2	3.5
Under 65	6.9	6.9	7.3	1.8	1.8	2.4
65 and over	10.4	9.1	18.9	5.6	5.0	11.1
All ages (WHO world std) ^c	6.3	6.3	7.3	1.7	1.6	2.5
<u>Age-Specific Rates, 2012-2016</u>						
Age at Diagnosis/Death:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	0.9	0.9	1.0	0.1	0.1	-
25-29	4.7	5.0	4.5	0.6	0.7	0.6
30-34	9.3	10.0	7.6	1.3	1.3	1.7
35-39	13.1	14.0	11.4	2.3	2.3	2.8
40-44	14.2	14.6	13.7	3.1	3.2	3.5
45-49	13.1	12.7	14.2	4.0	3.9	5.1
50-54	12.3	11.6	14.9	4.4	4.2	6.0
55-59	11.9	11.1	16.5	4.8	4.5	7.2
60-64	11.9	10.9	16.4	5.0	4.5	8.3
65-69	11.1	10.2	16.6	5.2	4.7	9.4
70-74	10.6	9.3	18.6	5.4	4.8	10.6
75-79	10.8	9.2	20.5	5.7	5.1	11.3
80-84	10.0	8.3	21.5	5.9	5.2	12.7
85+	8.2	6.9	19.1	6.3	5.6	13.5

^a SEER 21 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

- Rates are per 100,000 and are age-adjusted to the world (WHO 2000-2025) standard million.

- Statistic not shown. Rate based on less than 16 cases for the time interval.