Table 29.5

Age-Adjusted SEER Cancer Incidence Trends, 1975-2016a By International Classification of Childhood Cancer(ICCC)^b Selected Group and Subgroup^c Excluding benign brain and myelodysplastic syndromes All Races, Females

	Ages 0-14				Ages 0-19			
ICCC Group and Subgroup	% Change 1975-2016	APC 1975-2016	APC 1975-1995	APC 1996-2016	% Change 1975-2016	APC 1975-2016	APC 1975-1995	APC 1996-2016
All ICCC Groups Combined	49.3	0.7*	0.6*	0.7*	43.3	0.7*	0.7*	0.9*
I Leukemia I(a) Lymphoid leukemia I(b) Acute myeloid leukemia	91.1 102.2 55.4	0.7* 0.7* 0.6*	0.7 0.7 1.2	0.7* 0.6 0.4	81.9 101.0 47.7	0.8* 0.8* 0.8*	0.9 0.8 1.8*	0.9* 0.8* 0.7
II Lymphomas and reticuloendothelial neo. II(a) Hodgkin lymphoma II(b,c,e) Non-Hodgkin lymphoma	0.0 -30.5 42.0	0.4 - -	-1.0 - -	0.6 - -	-4.5 -24.8 42.8	0.1 -0.6* 1.6*	0.1 -0.4 1.0	0.2 -0.5 1.5
III CNS & misc intracranial & intraspinal neo. III(a) Ependymomas and choroid plexus tumors III(b) Astrocytoma III(c) Intracranial and intraspinal embryonal tum. III(d) Other gliomas	7.3 - 40.1 -	0.8* - 1.1* -	1.5* - 2.6* -	0.5 - 0.8 -	4.0 - 29.7 - -58.2	0.8* - 1.0* -	1.3* - 2.1* -	0.7 - 1.2 -0.6
<pre>IV Neuroblastoma and other peripheral nervous cell tumors IV(a) Neuroblastoma and ganglioneuroblastoma</pre>	-6.1 -6.1	0.3	0.0	0.2	-6.1 -6.1	0.3	-0.2 -0.3	0.2
V Retinoblastoma	-	-	-	-	-	-	-	-
VI Renal tumors VI(a) Nephroblastoma and other nonepithelial renal tumors	94.8 88.9	0.2 0.1	-0.2 -0.4	0.4 0.4	101.0 88.9	0.3 0.1	0.3 -0.2	0.6 0.3
VII Hepatic tumors VII(a) Hepatoblastoma	- -	- -	- -	- -	- -	- -	- -	-
VIII Malignant bone tumors VIII(a) Osteosarcoma VIII(c) Ewing tum. and related sarcomas of bone	118.8	- - -	0.7 - -	- - -	37.0 46.0 -	-0.1 0.4 -	0.1 0.6 -	1.2* 1.9* -
IX Soft tissue and other extraosseous sarcomas IX(a) Rhabdomyosarcoma	7.4	0.4	0.5	-0.3 -	39.4 5.0	0.4	0.2	-0.5 -
X Germ cell & trophoblastic tum. & neo. of gonads $X(a)$ Intracranial and intraspinal germ-cell tum. $X(c)$ Malignant gonadal germ cell tum. (ovary)	30.2	- - -	- - -	1.4 - -	-33.7 - -33.1	0.2 - -	1.2	0.7 - -0.2
XI Other malignant epithelial neo. and melanomas $^{\rm d}$ XI(b) Thyroid carcinoma XI(d) Malignant melanoma	135.1 - -	- - -	- - -	2.9* - -	134.3 188.2 108.8	1.6* 2.5* -	0.8 1.4 -	3.0* [#] 4.6* [#] -1.0

- The APC is the Annual Percent Change over the time interval.
- SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).
- International Classification of Childhood Cancer is based on ICD-O-3. Stellarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, Third Edition. Cancer. April 1, 2005: Vol 103, No. 7, pg 1457-1467.
- Classifications are shown for invasive cases only except as noted. For ICCC groups and subgroups that include in situ
- behavior, only invasive cases are shown except as noted.
- Includes in situ urinary bladder tumors.
- The APC is significantly different from zero (p<.05)
- The APC for 1996-2016 is significantly different from the APC for 1975-1995 (p<.05). @
- The APC for 1996-2016 is significantly different from the APC for 1975-1995 (p<.10).
- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval. Neoplasms are abbreviated as neo. Tumors are abbreviated as tum.