

Table 1.8  
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex  
Whites, 2008-2017

Site	Incidence <sup>a</sup>			US Mortality <sup>b</sup>		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.1*	-1.9*	-0.4*	-1.4*	-1.7*	-1.3*
Oral Cavity & Pharynx:						
Lip	0.6*	0.8*	0.2	1.0*	1.3*	0.2
Tongue	-0.9	-1.3	0.0	-1.3	-2.4	0.8
Salivary gland	1.7*	1.8*	1.4*	2.0*	2.4*	1.1
Floor of mouth	-0.4	-0.7	0.0	1.5*	1.7*	0.8
Gum & other oral cavity	-3.7*	-3.8*	-3.5*	-3.8*	-3.0	-5.8
Nasopharynx	-0.3	-0.2	-0.4	0.7	1.2	0.0
Tonsil	-2.0*	-1.6*	-3.0*	-1.3*	-1.2*	-1.4
Oropharynx	2.4*	2.6*	1.4	2.8*	3.0*	1.6
Hypopharynx	2.6*	2.9*	1.1	3.9*	4.4*	2.4*
Other oral cavity & pharynx	-2.5*	-2.5*	-3.2	1.5	1.9*	-1.1
Digestive System:	-3.5*	3.9*	1.8	-2.2*	-2.2*	-2.6*
Esophagus	-0.9*	-1.1*	-0.8*	-0.5*	-0.5*	-0.5*
Stomach	-1.2*	-1.4*	-1.0	-0.6*	-0.7*	-1.0*
Small intestine	-0.7*	-1.4*	0.2	-1.9*	-2.3*	-1.6*
Colon & Rectum:	1.1*	0.7	1.4*	1.8*	2.2*	1.3
Colon	-2.2*	-2.4*	-2.0*	-2.0*	-2.2*	-1.9*
Rectum	-2.5*	-2.8*	-2.2*	-	-	-
Anus, anal canal & anorectum	-1.5*	-1.7*	-1.5*	-	-	-
Liver & intrahepatic bile duct	1.3*	0.3	1.9*	3.3*	3.3*	3.4*
Gallbladder	1.9*	1.5*	2.8*	2.3*	2.0*	2.7*
Other biliary	-0.5	-0.3	-0.4	-1.3*	-1.3	-1.2*
Pancreas	-0.4	-0.5	-0.6	0.8	0.9	0.6
Retroperitoneum	0.5*	0.6*	0.4	0.3*	0.4*	0.1
Peritoneum, omentum & mesentery	-1.3	-1.2	-1.3	1.5	2.3	0.4
Other digestive system	-4.5*	-2.5	-4.6*	0.6	2.9*	0.2
Respiratory System:	6.1*	6.2*	5.7*	3.6*	2.9*	4.1*
Nose, nasal cavity & middle ear	-2.2*	-2.9*	-1.5*	-3.0*	-3.6*	-2.5*
Larynx	-0.3	-0.9	0.4	-1.0	-0.8	-1.6
Lung & bronchus	-3.0*	-3.1*	-2.9*	-2.0*	-2.1*	-2.4*
Pleura	-2.2*	-2.9*	-1.5*	-3.1*	-3.7*	-2.5*
Trachea & other respiratory organs	-5.9*	-7.1	-5.1	3.9*	3.4*	4.3*
Bones & joints	-1.6*	-1.2	-2.3*	1.3	0.4	2.5
Soft tissue (including heart)	0.3	0.5	0.1	1.1*	1.5*	0.5
Skin (excl. basal & squamous):	0.1	0.4	-0.4	0.2	0.2	0.2
Melanoma of the skin	1.2*	1.1*	1.3*	-1.3*	-1.2*	-1.6*
Other non-epithelial skin	1.3*	1.2*	1.4*	-2.6*	-2.7*	-2.7*
Breast	0.5	0.9	-0.3	2.6*	2.5*	2.3*
Breast (in situ)	-0.2	0.1	0.0	-1.6*	-0.8	-1.4*
	-1.9*	-0.5	-1.8*	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>a</sup> SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

\* The APC is significantly different from zero ( $p<.05$ ).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.8 - continued  
 SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex  
 Whites, 2008-2017

Site	Incidence <sup>a</sup>			US Mortality <sup>b</sup>		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:	-0.2	-	0.0	-0.6*	-	-0.4*
Cervix uteri	-0.8	-	-0.7	0.1	-	0.2
Corpus uteri	0.5*	-	0.7*	3.7*	-	4.0*
Uterus, NOS	2.1*	-	2.5*	-0.1	-	0.2
Ovary <sup>c</sup>	-2.6*	-	-2.4*	-2.5*	-	-2.2*
Vagina	-1.8*	-	-1.5*	-2.2*	-	-1.9*
Vulva	0.5	-	0.8*	1.7*	-	2.0*
Other female genital system	9.0*	-	9.1*	7.3*	-	7.6*
Male Genital System:	-4.4*	-4.7*	-	-1.2*	-1.9*	-
Prostate	-4.8*	-5.0*	-	-1.3*	-2.0*	-
Testis	0.5*	0.4*	-	0.6	0.4	-
Penis	0.2	0.0	-	3.1*	2.9*	-
Other male genital system	-1.0	-1.5	-	3.4	2.7	-
Urinary System:	-0.6*	-0.8*	-0.5*	-0.5*	-0.6*	-0.9*
Urinary bladder	-1.4*	-1.6*	-1.5*	-0.1	-0.3*	-0.3
Kidney & renal pelvis	0.5*	0.4	0.4	-1.0*	-1.0*	-1.4*
Ureter	-1.2*	-1.3	-1.1	0.2	0.6	-0.3
Other urinary system	2.0	0.9	4.2*	0.9	1.2	-0.3
Eye & Orbit	-0.4	-0.9	0.1	1.1	1.3	0.8
Brain & Nervous System: <sup>d</sup>	-0.7*	-0.8*	-0.6	0.6*	0.5*	0.6*
Brain	-0.7*	-0.8*	-0.6	-	-	-
Cranial nerves & other nervous system	-1.1	-0.3	-1.8*	-	-	-
Endocrine System:	0.7	1.2*	0.5	-0.1	0.4*	-0.4
Thyroid	0.7	1.4*	0.6	0.0	0.7	-0.5
Other endocrine & thymus	-1.0	-0.7	-1.3	-0.1	0.0	-0.3
Lymphoma:	-0.9*	-0.9*	-1.0*	-2.3*	-2.1*	-2.6*
Hodgkin lymphoma	-1.7*	-1.8*	-1.6*	-4.1*	-3.7*	-4.8*
Non-Hodgkin lymphoma	-0.8*	-0.8*	-0.9*	-2.1*	-2.0*	-2.5*
Myeloma	0.6	0.5	0.5	-0.4	-0.5*	-0.6
Leukemia:	-0.4	-0.5	-0.4	-1.5*	-1.6*	-1.5*
Lymphocytic:	-0.9*	-1.1*	-0.9*	-1.9*	-2.1*	-1.9*
Acute lymphocytic	0.3	0.6*	-0.2	-0.4	0.0	-1.1
Chronic lymphocytic	-1.3*	-1.5*	-1.3*	-2.6*	-2.8*	-2.4*
Other lymphocytic	-1.1	-1.4*	-0.6	-0.8	-1.4	0.3
Myeloid & Monocytic:	0.8	0.7	0.7	-0.1	-0.2	-0.1
Acute myeloid	1.1	1.1	1.0	-0.3	-0.4*	-0.2
Chronic myeloid	0.9	0.8	1.0*	0.0	-0.4	0.4
Acute monocytic	-4.2*	-4.4*	-4.1*	-1.8	-1.7	-2.3
Other myeloid & monocytic	-4.2*	-4.4*	-4.5*	2.1*	2.2*	1.6
Other leukemia:	-5.4*	-4.8*	-6.4*	-4.0*	-4.1*	-4.3*
Other acute leukemia	-3.0*	-2.5*	-3.6*	-6.2*	-6.1*	-6.4*
Aleukemic, subleukemic & NOS	-7.3*	-6.5*	-8.5*	-2.9*	-3.0*	-3.0*
Kaposi Sarcoma <sup>e</sup>	-5.0*	-5.1*	-5.2	-	-	-
Mesothelioma <sup>e</sup>	-2.9*	-3.9*	-0.2	-	-	-
Ill-defined & unspecified	-3.0*	-2.8*	-3.2*	-1.3*	-1.5*	-1.3*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

<sup>a</sup> SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

<sup>b</sup> US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

<sup>c</sup> Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

<sup>d</sup> Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

<sup>e</sup> Trend not shown for mortality. Category did not exist in mortality coding until 1999.

\* The APC is significantly different from zero ( $p<.05$ ).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.