

Table 1.9
SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
Blacks, 2008-2017

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
All Sites	-1.6*	-2.7*	-0.4*	-2.1*	-2.7*	-1.6*
Oral Cavity & Pharynx:						
Lip	-1.2*	-1.3*	-1.0	-1.7*	-2.0*	-1.4
Tongue	-5.3	-4.2	-5.3	-	-	-
Salivary gland	-2.1*	-1.8*	-2.7*	-1.7	-2.3*	-0.8
Floor of mouth	1.0	-0.4	2.0	0.3	-0.9	1.4
Gum & other oral cavity	-4.4*	-4.1*	-4.7	-	-	-
Nasopharynx	-1.6	-1.6	-1.6	-0.9	-1.6	0.3
Tonsil	0.1	0.1	-0.6	-1.4	-0.1	-4.0*
Oropharynx	0.4	0.7	-0.6	-1.4	-1.2	-0.9
Hypopharynx	0.0	0.6	-1.4	-0.4	-0.2	-1.2
Other oral cavity & pharynx	-3.1*	-4.2*	1.6	0.7	2.7	-
Digestive System:	-1.6*	-1.8*	-1.5*	-1.5*	-1.7*	-1.5*
Esophagus	-4.1*	-4.5*	-3.8*	-4.6*	-5.1*	-3.7*
Stomach	-2.1*	-2.3*	-2.1*	-3.4*	-3.2*	-3.9*
Small intestine	1.9*	1.4*	2.3*	1.9	0.6	3.4
Colon & Rectum:	-3.0*	-3.0*	-3.0*	-2.8*	-2.8*	-2.9*
Colon	-3.3*	-3.5*	-3.2*	-	-	-
Rectum	-2.0*	-1.8*	-2.3*	-	-	-
Anus, anal canal & anorectum	0.3	-0.4	1.1	1.5*	0.9	2.6
Liver & intrahepatic bile duct	0.7	0.7	0.8	1.4*	1.1*	2.1*
Gallbladder	0.1	-0.8	1.1	-0.9	-2.6*	-0.2
Other biliary	1.0	0.8	1.1	1.9	1.6	2.3
Pancreas	0.1	-0.1	0.3	-0.3*	-0.2	-0.4
Retroperitoneum	-1.6	-0.8	-1.7	7.1*	-	-
Peritoneum, omentum & mesentery	0.5	2.9	0.4	5.6*	-	6.4*
Other digestive system	3.2*	0.7	4.9*	1.9	1.7	1.8
Respiratory System:	-2.7*	-3.4*	-2.0*	-3.5*	-4.2*	-2.8*
Nose, nasal cavity & middle ear	0.1	-0.1	0.9	-1.5	-1.5	-0.9
Larynx	-2.9*	-3.1*	-3.1*	-2.9*	-3.2*	-2.2
Lung & bronchus	-2.7*	-3.5*	-2.0*	-3.6*	-4.3*	-2.9*
Pleura	-6.8	-	-	-	-	-
Trachea & other respiratory organs	-2.8	-4.8	-0.5	2.1	-	-
Bones & joints	0.4	1.2	-0.3	0.6	1.0	0.8
Soft tissue (including heart)	0.5	1.3	-0.2	0.4	0.4	0.5
Skin (excl. basal & squamous):	-0.6	-0.9	-0.2	-2.1*	-3.0*	-0.9
Melanoma of the skin	-1.1	0.1	-1.9	-3.3*	-3.4	-3.1
Other non-epithelial skin	-0.3	-1.5	1.3	-1.1	-2.8*	2.4
Breast	0.2	1.7	0.3	-1.7*	-0.9	-1.6*
Breast (<i>in situ</i>)	0.6	-0.4	0.7	-	-	-

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^a SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

* The APC is significantly different from zero ($p<.05$).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.

Table 1.9 - continued
 SEER Incidence and U.S. Mortality Trends by Primary Cancer Site and Sex
 Blacks, 2008-2017

Site	Incidence ^a			US Mortality ^b		
	Total APC	Males APC	Females APC	Total APC	Males APC	Females APC
Female Genital System:						
Cervix uteri	0.2	-	0.4	-0.4	-	-0.2
Corpus uteri	-2.6*	-	-2.4*	-3.0*	-	-2.9*
Uterus, NOS	1.4*	-	1.6*	4.3*	-	4.5*
Ovary ^c	2.6*	-	2.8*	0.6	-	0.9
Vagina	-1.7*	-	-1.4*	-2.1*	-	-1.9*
Vulva	-0.2	-	0.0	-1.3	-	-1.2
Other female genital system	0.5	-	0.8	-1.8	-	-1.6
	8.2*	-	8.5*	6.2*	-	6.4*
Male Genital System:						
Prostate	-4.1*	-4.4*	-	-3.0*	-3.6*	-
Testis	-4.2*	-4.5*	-	-3.0*	-3.7*	-
Penis	2.6*	2.4*	-	2.0	2.3	-
Other male genital system	-1.4	-1.9	-	-3.6	-3.7	-
	1.1	0.6	-	-	-	-
Urinary System:						
Urinary bladder	-0.3	-0.5*	-0.2	-0.8*	-0.3	-1.6*
Kidney & renal pelvis	-0.8*	-1.1*	-0.4	-0.3	0.0	-0.9
Ureter	0.1	0.1	-0.2	-1.3*	-0.8	-2.2*
Other urinary system	-1.7	-5.6*	1.8	-3.8	-	-
	1.5	-1.7	4.7	1.4	-	-2.0
Eye & Orbit	-0.5	-1.7	0.4	-	-	-
Brain & Nervous System: ^d						
Brain	-1.2*	-1.1	-1.6	0.7	0.5	0.8
Cranial nerves & other nervous system	-1.2	-1.1	-1.5	-	-	-
	-1.9	-1.5	-2.0	-	-	-
Endocrine System:						
Thyroid	1.1	0.4	1.4	-0.7	-0.8	-0.7
Other endocrine & thymus	1.3	0.7	1.5	-0.6	0.0	-1.1
	-0.2	-0.6	-0.2	-0.8	-1.4	0.1
Lymphoma:						
Hodgkin lymphoma	-0.7*	-1.1*	-0.2	-1.8*	-1.8*	-1.9*
Non-Hodgkin lymphoma	-1.7*	-1.9	-1.3*	-5.0*	-4.3*	-5.3*
	-0.5*	-1.0*	0.0	-1.6*	-1.6*	-1.6*
Myeloma	0.3	0.2	0.3	-0.4	-0.6	-0.3
Leukemia:						
Lymphocytic:	0.3	-0.1	0.7	-2.0*	-2.7*	-1.4*
Acute lymphocytic	-0.5	-1.2	0.4	-3.0*	-4.2*	-1.4
Chronic lymphocytic	0.2	-0.9	1.8	-0.1	-1.2	1.3
Other lymphocytic	-0.4	-0.7	-0.3	-4.0*	-4.8*	-2.9*
Myeloid & Monocytic:	-4.0*	-9.4*	4.8	0.6	-	-
Acute myeloid	1.5*	1.4	1.4	0.3	-0.3	0.8
Chronic myeloid	2.0*	2.0	1.8	0.2	-0.5	0.8
Acute monocytic	0.9	0.9	0.9	-1.1	-2.3	-0.2
Other myeloid & monocytic	-0.5	0.8	-1.9	-	-	-
Other leukemia:	-2.7	-3.5	-1.4	3.7*	5.0*	2.3
Other acute leukemia	-3.7*	-3.9	-3.3	-5.0*	-5.0*	-5.3*
Aleukemic, subleukemic & NOS	1.8	1.4	3.6	-7.0*	-7.4*	-6.8*
	-5.6*	-6.5*	-4.8	-4.2*	-4.0*	-4.6*
Kaposi Sarcoma ^e	-3.0*	-3.0	-4.0	-	-	-
Mesothelioma ^e	-1.4	-1.7	-0.1	-	-	-
Ill-defined & unspecified	-2.7*	-2.7*	-2.6*	-2.4*	-3.0*	-1.8*

The APC is the Annual Percent Change over the time interval.

Trends are based on rates age-adjusted to the 2000 US Std Population
 (19 age groups - Census P25-1130).

^a SEER 21 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey, Georgia excluding ATL/RG, Idaho, New York and Massachusetts).

^b US Mortality Files, National Center for Health Statistics, Centers for Disease Control and Prevention.

^c Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.

^d Due to coding changes, Brain & Nervous System mortality are no longer shown separately.

^e Trend not shown for mortality. Category did not exist in mortality coding until 1999.

* The APC is significantly different from zero ($p<.05$).

- Statistic could not be calculated. Trend based on less than 10 cases for at least one year within the time interval.