Table 28.8 5-Year Relative Survival (Percent)

## By Selected Primary Cancer Site and Year of Diagnosis

All Races, Males and Females

	Ages 0-14											
	1975-	1978-	1981-	1984-	1987-	1990-	1993-	1996-	1999-	2002-	2005-	2010-
Site	1977	1980	1983	1986	1989	1992	1995	1998	2001	2004	2009	2016
All Sites												
All Races	58.0	62.5	67.1	68.2	71.7	75.8	77.4	79.2	80.8	82.8	83.5	85.6°
Whites	58.0	63.1	67.8	69.9	72.5	76.7	78.3	80.6	81.9	84.8	84.9	86.5°
Blacks	57.3	57.7	62.2	56.6	65.3	70.8	73.3	75.6	74.1	73.5	78.3	80.9°
Bone & Joint	50.1 <sup>d</sup>	47.8	56.8 <sup>d</sup>	57.3 <sup>d</sup>	66.8 <sup>d</sup>	67.8	74.2	70.3	69.3	77.7	78.5	73.8°
Brain & ONS	57.2	57.9	57.2	61.8	64.5	64.3	70.7	75.3	74.4	75.5	75.0	76.3°
Hodgkin lymphoma	81.1	86.8	88.1	89.9	88.1	96.8	94.6	96.1	94.4	97.5	97.6	99.5°
Leukemia	49.7	58.0	62.8	63.7	70.9	75.5	75.9	80.3	82.8	86.0	86.7	88.9°
Acute lymphocytic	57.2	65.7	71.1	72.2	77.7	83.1	83.9	87.0	88.5	91.8	91.1	92.5°
Acute myeloid	18.8	26.3	27.8 <sup>d</sup>	30.6 <sup>d</sup>	37.1 <sup>d</sup>	42.2	40.6 <sup>d</sup>	48.7	58.6	61.5	65.9	70.6°
Neuroblastoma <sup>a</sup>	52.5	56.6	54.8	52.3	63.2	76.0	66.6	66.7	72.3	73.8	75.3	79.5°
Non-Hodgkin lymphoma	43.2	53.0	67.0	69.8	70.8	76.9	80.7	83.2	89.8	84.6	87.2	90.1°
Soft tissue	61.3	74.2	69.2	73.0	66.4	79.7	76.7	70.6	76.9	84.8	80.6	83.0°
Wilms' tumor <sup>b</sup>	73.1	79.0	86.7	90.7	92.2	91.9	91.7	91.6	93.8	89.5	92.4	96.3°
	Ages 0-19											
	1975-	1978-	1981-	1984-	1987-	1990-	1993-	1996-	1999-	2002-	2005-	2010-
Site	1977	1980	1983	1986	1989	1992	1995	1998	2001	2004	2009	2016
All Sites						<b>5</b> 6.0						0= 00
All Races	61.6	65.3	68.4	70.6	73.6	76.3	77.8	79.8	80.4	82.6	84.1	85.9°
Whites	61.4	65.9	69.3	72.5	75.1	77.6	78.5	81.1	81.5	84.1	85.4	87.0°
Blacks	59.2	60.3	62.5	58.3	64.3	68.7	74.2	73.9	74.8	73.3	78.5	80.3°
Bone & Joint	50.5	48.1	51.2	56.1	64.2	68.9	69.0	67.0	66.4	73.5	75.1	70.8°
Brain & ONS	59.1	58.3	58.7	64.0	66.1	66.3	71.5	76.1	74.8	75.5	75.6	76.5°
Hodgkin lymphoma	86.2	88.3	85.5	90.6	89.0	94.3	93.9	95.2	95.1	96.3	97.2	98.2°
Leukemia	44.9	53.2	58.0	60.4	67.5	71.3	71.9	76.6	77.7	81.8	85.0	86.2°
Acute lymphocytic	53.8	61.9	66.9	69.9	75.0	79.8	81.5	84.4	84.8	88.0	90.0	90.4°
Acute myeloid	18.7	26.5	27.2	31.4	37.6	41.9	38.6	46.3	51.9	59.8	64.8	69.4°
Neuroblastoma <sup>a</sup>	53.1	56.7	53.7	52.5	61.8	75.7	66.6	66.7	72.1	73.5	74.9	79.2°
Non-Hodgkin lymphoma	44.6	53.6	63.7	67.6	70.4	72.7	78.0	81.1	84.3	84.5	85.5	89.5°
Soft tissue	65.4	68.9	68.2	72.7	68.0	69.5	73.8	72.4	72.5	77.0	78.5	79.6°
Wilms' tumor <sup>b</sup>	72.6	78.4	86.8	91.0	92.2	91.3	91.8	91.6	93.8	89.6	92.0	96.3°

Source: SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Based on follow-up of patients into 2016. Expected survival rates are derived from the U.S. Annual Life Tables.

Note: Neuroblastoma and Wilms' tumor are not mutually exclusive from the other tumors presented in table.

Neuroblastoma is defined as histologies 9490-9509.

Wilms' tumor is defined as histologies 8959-8960.

The difference between 1975-1977 and 2010-2016 is statistically significant (p<.05).

The standard error is between 5 and 10 percentage points.

The standard error is greater than 10 percentage points.

Statistic gould not be calculated due to fewer than 25 cases during the time period.

Statistic could not be calculated due to fewer than 25 cases during the time period.