

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

November 2011 Data Submission					
	Item # refers to the NAACCR item number - see https://www.naacr.org/StandardsandRegistryOperations/Volumell.aspx				
	CS= Collaborative Staging				
	SSF = Site-specific Factor				
Field number	Name	NAACCR Item #	Description	Category name	Category number
1	Age recode with <1 year olds		<p>The age recode variable is based on Age at Diagnosis (single-year ages). The groupings used in the age recode variable are determined by the age groupings in the population data. This recode has 19 age groups in the age recode variable (< 1 year, 1-4 years, 5-9 years, ..., 85+ years). See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf.</p> <p>Will be in Race and Age (case data only) in rate or prevalence sessions if an alternate age is used as the population age variable.</p>	Age at Diagnosis (or Race and Age (case data only))	1 (or 13)
2	Race recode (White, Black, Other)		Race recode is based on the race variables and the American Indian/Native American IHS link variable. This recode should be used to link to the populations for white, black and other. It is independent of Hispanic ethnicity. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity .	Race, Sex, Year Dx, Registry, County	2
3	Sex	220	Includes 1= Male and 2=Female from Sex [NAACCR Item #220] plus a total of male and female. This is used to link to the correct populations for males and females when calculating sex-specific rates.	Race, Sex, Year Dx, Registry, County	2
4	Year of diagnosis	390	Year of Diagnosis: values are 1973-2009 but may be a subset depending on the file that is used and the registry that is selected. There are no unknown values on the file.	Race, Sex, Year Dx, Registry, County	2
5	SEER registry	40	This field show the SEER registries which contribute data to this file. After the name of the registry, there is the beginning year of diagnosis for that registry. This data item varies by which data file is selected. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Race, Sex, Year Dx, Registry, County	2
6	Louisiana 2005 - 1st vs 2nd half of year		This field is used to separate Louisiana cases diagnosed in the first half of 2005 from those diagnosed in the second half of 2005 to link to different population estimates used to account for displaced persons due to Katrina/Rita. See https://seer.cancer.gov/data/hurricane.html	Race, Sex, Year Dx, Registry, County	2
7	County	90	County of residence at diagnosis. This must be used in conjunction with SEER registry or use State-county variable. For more information, see https://seer.cancer.gov/manuals/2004Revision%201/SPM_AppendixA.pdf .	Race, Sex, Year Dx, Registry, County	2
8	State-county		State and county at diagnosis. Can be used to link to the populations to produce rates at the state/county level.	Race, Sex, Year Dx, Registry, County	2

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
9	In research data		Flag indicates whether it is supplemental data or not. In research databases which include Louisiana, July-December 2005 cases are considered supplemental data. These cases/populations are set to "No" for this field and are typically excluded in SEER analyses. This field is associated with the Research data check box on the selection tab in all SEER*Stat sessions other than prevalence. For more information, see: https://seer.cancer.gov/data/hurricane.html	Race, Sex, Year Dx, Registry, County	2
10	In research data for prevalence		Flag indicates whether it is supplemental data or not. Only available in prevalence sessions when using a research databases which includes Louisiana. All Louisiana cases are considered supplemental data in this situation. These cases/populations are set to "No" for this field and are typically excluded in SEER analyses. This field is associated with the Research data check box on the selection tab SEER*Stat prevalence sessions. For more information, see: https://seer.cancer.gov/data/hurricane.html	Race, Sex, Year Dx, Registry, County	2
11	CHSDA 2006		This data item identifies whether or not the county of diagnosis is served by CHSDA. The primary use of this field is to be able to limit analyses of AI/AN race to areas served by CHSDA. See https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity OR https://seer.cancer.gov/seerstat/variables/countyattribs/	Race, Sex, Year Dx, Registry, County	2
12	CHSDA Region		This data item is a grouping of counties that is primarily used when working with CHSDA 2006. See https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity OR https://seer.cancer.gov/seerstat/variables/countyattribs/	Race, Sex, Year Dx, Registry, County	2
13	Site rec with Kaposi and mesothelioma		A recode based on Primary Site and Histology in order to make analyses of site/histology groups easier. For example, the lymphomas are excluded from stomach and Kaposi and mesothelioma are separate categories based on histology. For more information, see https://seer.cancer.gov/siterecode/icdo3_d01272003 .	Site and Morphology	3
14	Behavior recode for analysis		This recode was created so that data analyses could eliminate major groups of histologies/behaviors that weren't collected consistently over time, for example benign brain, myelodysplastic syndromes, and borderline tumors of the ovary. Created from ICD-O-3 behavior and histology. For more information, see https://seer.cancer.gov/behavrecode .	Site and Morphology	3
15	AYA site recode		A site/histology recode that is mainly used to analyze data on adolescent and young adults. The recode was applied to all cases no matter the age in order that age comparisons can be made with these groupings. For more information, see https://seer.cancer.gov/ayarecode/index.html .	Site and Morphology	3
16	Lymphoma subtype recode		A site/histology recode that is mainly used to analyze data on lymphoma sub-types. Based on ICD-O-3. Note that cases diagnosed before 2001 were not coded under ICD-O-3 and were converted to ICD-O-3 and may not have the specificity of cases after 2000 that were coded directly under ICD-O-3. For more information, see https://seer.cancer.gov/lymphomarecode/index.html .	Site and Morphology	3

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
17	ICCC site recode ICD-O-3		A site/histology recode that is mainly used to analyze data on children. The recode was applied to all cases no matter the age in order that age comparisons can be made with these groupings. Based on ICD-O-3. Note that cases diagnosed before 2001 were not coded under ICD-O-3 and were converted to ICD-O-3 and may not have the specificity of cases after 2000 that were coded directly under ICD-O-3. For more information on this International Classification of Childhood Cancer (ICCC) site recode, see https://seer.cancer.gov/iccc .	Site and Morphology	3
18	CS Schema - AJCC 6th ed		CS information is collected under the specifications of a particular schema based on site and histology. This recode should used in any analysis of AJCC 6th ed stage and T, N, M. Based on CS version 1, it should not be used for SSFs collected or modified under CS v02. https://cancerstaging.org/cstage/schema.html . For more information see ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf	Site and Morphology	3
19	CS Schema - v0203		CS information is collected under the specifications of a particular schema based on site and histology. This recode should used in any analysis of AJCC 7th ed stage and T, N, M. For more information see ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf	Site and Morphology	3
20	Primary Site - labeled	400	This provides the primary site code in ICD-O-3 and a descriptive primary site label. Note that the label is the preferred ICD-O-3 bolded name and there may be other sites or sub-sites included in the code but not reflected in the preferred term. Refer to ICD-O-3 for further information. Cases with years of diagnosis before 1992 were converted to ICD-O-3 from earlier versions.	Site and Morphology	3
21	Primary Site	400	Codes are found in the Topography section of the International Classification of Diseases for Oncology (ICD-O) 3rd edition. ICD-O-2 codes used for 1992-2000 are similar. Primary site codes for 1973-1991 were converted to ICD-O-3 and may lack the specificity of the ICD-O-3 primary site codes.	Site and Morphology	3
22	Histologic Type ICD-O-3	522	Based on histology codes in ICD-O-3. Cases diagnosed in 1973-2000 were coded in earlier versions and converted to ICD-O-3 and coded directly for 2001+.	Site and Morphology	3
23	Behavior code ICD-O-3 (1973+)		Based on behavior codes in ICD-O-3. Behavior code has had the same definitions in previous versions but may be associated differently with histologies over time. For example, borderline of the ovary were considered malignant in ICD-O-2 but benign in ICD-O-3. cases diagnosed in 1973-2000 were coded in earlier versions and converted to ICD-O-3 and coded directly for 2001+. In situ bladder cases have been converted to malignant in this field. See Behavior recode for consistency over time. For more information see ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=22	Site and Morphology	3

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
24	Grade	440	Based on grade codes in ICD-O-3. Cases diagnosed in 1973-2000 were coded in earlier versions and may lack the specificity of the 2001+ cases that were coded directly, especially for lymphomas/leukemias. For more information see ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=23	Site and Morphology	3
25	Diagnostic Confirmation	490	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=23	Site and Morphology	3
26	ICD-O-3 Hist/behav		Labeled version of ICD-O-3 values for all behaviors. See SEER*Stat dictionary for labels.	Site and Morphology	3
27	ICD-O-3 Hist/behav, malignant		Labeled version of ICD-O-3 values for malignant tumors. All non-malignant tumors are grouped into one value. See SEER*Stat dictionary for labels.	Site and Morphology	3
28	Histology recode - broad groupings		Based on Histologic type ICD-O-3. For more information see ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=63	Site and Morphology	3
29	Histology recode - brain groupings		Based on Histologic type ICD-O-3. For more information see ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=65	Site and Morphology	3
30	Site rec B with Kaposi and mesothelioma		A recode based on Primary Site and Histology in order to make analyses of site/histology groups easier for multiple primary analyses. For example, the lymphomas are excluded from stomach and Kaposi and mesothelioma are separate categories based on histology. For more information, see https://seer.cancer.gov/siterecode_b/icdo3_d12192003 .	Site and Morphology	3
31	Site recode		A recode based on Primary Site and Histology in order to make analyses of site/histology groups easier. For example, the lymphomas are excluded from stomach and included with Lymphoma. Note: In this recode Kaposi and mesothelioma are not separate categories based on histology. To match the site groupings in the CSR, do NOT use this recode -- use 'Site rec with Kaposi and mesothelioma'. For more information, see https://seer.cancer.gov/siterecode/icdo3_d01272003 .	Site and Morphology	3
32	ICCC site recode extended ICD-O-3		Based on ICD-O-3. For more information on this International Classification of Childhood Cancer (ICCC) site/histology recode, see https://seer.cancer.gov/iccc . While the recode is normally used for childhood cancers, it is on the file for all ages so that childhood cancers could be compared across age groups.	Site and Morphology	3
33	Derived AJCC Stage Group, 6th ed (2004+)	3000	The stage category for AJCC 6th edition is derived from Collaborative Staging data elements for 2004+ cases. See the CS site-specific schema for details (https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage) and the ASCII text file description for allowable values for display codes and storage values (https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf).	Stage - AJCC	4

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
34	Derived AJCC - Flag (2004+)	3030	This flag currently reflects only when AJCC stage is derived based on CS. If AJCC stage is derived based on cases before 2004, it will currently be found in a separate field and not overlaid in the Derived AJCC fields. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Stage - AJCC	4
35	AJCC stage 3rd edition (1988-2003)		Derived by algorithm from extent of disease (EOD). Not available for all years or for all sites. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Stage - AJCC	4
36	SEER modified AJCC stage 3rd (1988-2003)		Derived by algorithm from extent of disease (EOD). Not available for all years or for all sites. The modified version stages cases that would be unstaged under strict AJCC staging rules. For example, it assumes NX is N0. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Stage - AJCC	4
37	Derived AJCC T, 6th ed (2004+)	2940	The T category for AJCC 6th edition is derived from Collaborative Staging data elements for 2004+ cases. See the CS site-specific schema for details (https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage) and the ASCII test file description for allowable values for display codes and storage values (https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf).	Stage - TNM	5
38	Derived AJCC N, 6th ed (2004+)	2960	The N category for AJCC 6th edition is derived from Collaborative Staging data elements for 2004+ cases. See the CS site-specific schema for details (https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage) and the ASCII test file description for allowable values for display codes and storage values (https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf).	Stage - TNM	5
39	Derived AJCC M, 6th ed (2004+)	2980	The M category for AJCC 6th edition is derived from Collaborative Staging data elements for 2004+ cases. See the CS site-specific schema for details (https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage) and the ASCII test file description for allowable values for display codes and storage values (https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf).	Stage - TNM	5
40	Derived SS1977 (2004+)	3010	Derived Summary Stage 1977 is derived from Collaborative Staging data elements for 2004+ cases. See the CS site-specific schema for details (https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage) and the ASCII test file description for allowable values for display codes and storage values (https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf).	Stage - LRD (Summary and Historic)	6

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
41	Derived SS2000 (2004+)	3020	Derived Summary Stage 2000 is derived from Collaborative Staging data elements for 2004+ cases. See the CS site-specific schema for details (https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage) and the ASCII test file description for allowable values for display codes and storage values (https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf).	Stage - LRD (Summary and Historic)	6
42	Summary stage 2000 (1998+)		Summary Stage 2000 is derived from Collaborative Stage (CS) for 2004+ and Extent of Disease (EOD) from 1998-2003. It is a simplified version of stage: in situ, localized, regional, distant, & unknown. Used in the SEER CSR and more recent SEER publications. For more information including sites and years for which it isn't calculated, see https://seer.cancer.gov/seerstat/variables/seer/lrd-stage .	Stage - LRD (Summary and Historic)	6
43	SEER historic stage A		SEER Historic Stage A is Derived from Collaborative Stage (CS) for 2004+ and Extent of Disease (EOD) from 1973-2003. It is a simplified version of stage: in situ, localized, regional, distant, & unknown. For more information including sites and years for which it isn't calculated, see https://seer.cancer.gov/seerstat/variables/seer/lrd-stage .	Stage - LRD (Summary and Historic)	6
44	SEER summary stage 2000 (2001-2003)		Summary Stage 2000 for 2001-2003 is based on SEER Extent of Disease (EOD) following a SEER algorithm. This variable is provided on the NAACCR call for data. For more information, see https://seer.cancer.gov/tools/ssm .	Stage - LRD (Summary and Historic)	6
45	SEER summary stage 1977 (1995-2000)		SEER summary stage 1977 (1995-2000) is based on SEER Extent of Disease (EOD) following a SEER algorithm. This variable is provided on the NAACCR call for data. For more information, see https://seer.cancer.gov/manuals/historic/ssm_1977.pdf .	Stage - LRD (Summary and Historic)	6
46	Derived SS1977 - Flag (2004+)	3040	Flag used to indicate if derivation was based on CS or EOD. Currently only based on CS.	Stage - LRD (Summary and Historic)	6
47	Derived SS2000 - Flag (2004+)	3050	Flag used to indicate if derivation was based on CS or EOD. Currently only based on CS.	Stage - LRD (Summary and Historic)	6
48	RX Summ--Surg Prim Site (1998+)	1290	NAACCR Name=RX Summ--Surg Prim Site, Item #=1290. The information in this field is site-specific. The amount/detail of information has varied over time and caution should be used when looking at trends over time. For site-specific codes, see Appendix C of https://seer.cancer.gov/tools/codingmanuals . Data for 1998-2002 are converted from Surgery of primary site (1998-2002), documented here: https://seer.cancer.gov/manuals/historic/AppendC.pdf . See here for changes between the two coding systems: https://seer.cancer.gov/tools/SEER2003.code.changes.122302.pdf .	Therapy	7

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
49	RX Summ--Scope Reg LN Sur (2003+)	1292	Changed to Unknown or not applicable for breast cancer cases. See https://seer.cancer.gov/seerstat/variables/seer/regional_ln/ for more information. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
50	RX Summ--Surg Oth Reg/Dis (2003+)	1294	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
51	Radiation sequence with surgery	1380	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
52	Reason no cancer-directed surgery	1340	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
53	Radiation	1360	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
54	Surgery of primary site (1998-2002)	1646	NAACCR Name=RX Summ--Surg Site 98-02, Item #=1646. For more information, see https://seer.cancer.gov/manuals/historic/AppendC.pdf .	Therapy	7
55	Scope of reg lymph nd surg (1998-2002)	1647	Changed to Unknown or not applicable for breast cancer cases. See https://seer.cancer.gov/seerstat/variables/seer/regional_ln/ for more information. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
56	Num of regional lym nd exam (1998-2002)	1296	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
57	Surgery of oth reg/dis sites (1998-2002)	1648	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Therapy	7
58	First course of reconstruct (1998-2002)	1330	NAACCR Name=RX Summ--Reconstruct 1st, Item #=1330. For more information, see https://seer.cancer.gov/manuals/historic/AppendC.pdf .	Therapy	7
59	Site specific surgery (1983-1997)	1640	NAACCR Name=RX Summ--Surgery Type, Item #=1640. For more information, see https://seer.cancer.gov/manuals/historic/AppendD.pdf .	Therapy	7
60	Radiation to Brain or CNS (1988-1997)	1370	NAACCR Name=RX Summ--Rad to CNS, Item #=1370	Therapy	7

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
61	CS tumor size (2004+)	2800	Information on tumor size. Available for 2004+. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
62	CS extension (2004+)	2810	Information on extension of the tumor. Available for 2004+. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage . Note: this item was originally a 2 digit field and was expanded to 3 digits during conversion. Generally, a zero was added to the right of the existing 2 digit field except for 99 which became 999.	Extent of Disease - CS	8
63	CS lymph nodes (2004+)	2830	Information on involvement of lymph nodes. Available for 2004+. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage . Note: this item was originally a 2 digit field and was expanded to 3 digits during conversion. Generally, a zero was added to the right of the existing 2 digit field except for 99 which became 999.	Extent of Disease - CS	8
64	CS mets at dx (2004+)	2850	Information on distant metastasis. Available for 2004+. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
65	CS site-specific factor 1 (2004+)	2880	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
66	ER Status Recode Breast Cancer (1990+)		Created by combining information from Tumor marker 1 (1990-2003) (NAACCR Item #=1150), with information from CS site-specific factor 1 (2004+) (NAACCR Item #=2880). This field is blank for non-breast cases and cases diagnosed before 1990.	Extent of Disease - CS	8
67	CS site-specific factor 2 (2004+)	2890	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
68	PR Status Recode Breast Cancer (1990+)		Created by combining information from Tumor marker 2 (1990-2003) (NAACCR Item #=1150), with information from CS site-specific factor 2 (2004+) (NAACCR Item #=2880). This field is blank for non-breast cases and cases diagnosed before 1990.	Extent of Disease - CS	8

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
69	CS site-specific factor 3 (2004+)	2900	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
70	CS site-specific factor 4 (2004+)	2910	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
71	CS site-specific factor 5 (2004+)	2920	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
72	CS site-specific factor 6 (2004+)	2930	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
73	CS site-specific factor 8 (2004+)	2862	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
74	CS site-specific factor 10 (2004+)	2864	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
75	CS site-specific factor 11 (2004+)	2865	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
76	CS site-specific factor 13 (2004+)	2867	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
77	CS site-specific factor 15 (2004+)	2869	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
78	CS site-specific factor 16 (2004+)	2870	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
79	CS site-specific factor 25 (2004+)	2879	Each CS site-specific factor (SSF) is schema dependent. They can provide information needed to stage the case, clinically relevant information, or prognostic information. Available for varying years and schemas depending on standard setter requirements. Earlier cases may be converted and new codes added which weren't available for use prior to the current version of CS. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
80	CS Tumor Size/Ext Eval (2004+)	2820	Available for 2004+, but not required for the entire timeframe. Will be blank in cases not collected. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
81	CS Reg Node Eval (2004+)	2840	Available for 2004+, but not required for the entire timeframe. Will be blank in cases not collected. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
82	CS Mets Eval (2004+)	2860	Available for 2004+, but not required for the entire timeframe. Will be blank in cases not collected. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
83	Regional nodes examined (1988+)	830	Currently coded under CS - https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage . Cases coded 1988-2003 used slightly different definitions - see SEER coding manual for those years. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=27	Extent of Disease - CS	8
84	Regional nodes positive (1988+)	820	Currently coded under CS - see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage . Cases coded 1988-2003 used slightly different definitions - see SEER coding manual for 1988. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=26	Extent of Disease - CS	8

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
85	Laterality (1973+)	410	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Extent of Disease - CS	8
86	Lymph-vascular Invasion (2004+)	1182	Required for cases originally coded under CSv2 or diagnosed 2010+ for the schemas for penis and testis only. On the research file LVI is shown only for testis because it is needed for AJCC 6th ed tagging. For penis, LVI is needed for AJCC 7th ed staging only which isn't calculated until 2010. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Extent of Disease - CS	8
87	CS version input current (2004+)	2937	Data item shows what version was in effect the last time that any CS field was updated for this case. Special codes for converted cases. This data item along with CS version input original gives info on what document to use for the CS codes. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
88	CS version input original (2004+)	2935	Data item shows what version was in effect the first time that CS was coded for this case. This data item along with CS version input current gives info on what document to use for the CS codes. For more information, see https://seer.cancer.gov/seerstat/variables/seer/ajcc-stage .	Extent of Disease - CS	8
89	CS version derived (2004+)	2936	Data item shows what CS version was used to derive the CS Derived item fields including T, N, M, AJCC stage and SEER Summary stages 1977 and 2000.	Extent of Disease - CS	8
90	EOD 10 - Prostate path ext (1995-2003)	800	Information on extension of the tumor from the primary based on information from the prostatectomy for prostate cancer only. For more information, see https://seer.cancer.gov/manuals/EOD10Dig.pub.pdf . Note: for 2004+ similar type of information was collected in CS SSF 3 in the collaborative stage variables.	Extent of Disease - Historic	9
91	EOD 10 - extent (1988-2003)	790	Information on extension of the tumor from the primary and distant metastases. For more information, see https://seer.cancer.gov/manuals/EOD10Dig.pub.pdf . Note: for 2004+ similar type of information was collected in CS extension (2004+) and CS Mets at DX (2004+) in the collaborative stage variables.	Extent of Disease - Historic	9
92	EOD 10 - nodes (1988-2003)	810	Information on lymph node involvement. Information is site-specific. For more information, see https://seer.cancer.gov/manuals/EOD10Dig.pub.pdf . Note: for 2004+ similar type of information was collected in CS Lymph Nodes (2004+) in the collaborative stage variables.	Extent of Disease - Historic	9
93	EOD 10 - size (1988-2003)	780	Information on size of tumor. For more information, see https://seer.cancer.gov/manuals/EOD10Dig.pub.pdf . Note: for 2004+ similar type of information was collected in CS Tumor Size (2004+) in the collaborative stage variables.	Extent of Disease - Historic	9

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
94	Tumor marker 1 (1990-2003)	1150	This data item records prognostic indicators for breast cases (ERA 1990-2003), prostate cases (PAP 1998-2003) and testis cases (AFP 1998-2003). Please see CS SSFs for similar information for 2004+. For breast cancer cases ERA over time is available in ER Status Recode Breast Cancer (1990+). https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf	Extent of Disease - Historic	9
95	Tumor marker 2 (1990-2003)	1160	This data item records prognostic indicators for breast cases (PRA 1990-2003), prostate cases (PSA 1998-2003) and testis cases (hCG 1998-2003). Please see CS SSFs for similar information for 2004+. For breast cancer cases PRA over time is available in PR Status Recode Breast Cancer (1990+). https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf	Extent of Disease - Historic	9
96	Tumor marker 3 (1998-2003)	1170	This data item records prognostic indicators for testis cases (LDH 1998-2003). Please see CS SSFs for similar information for 2004+. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf	Extent of Disease - Historic	9
97	Coding system-EOD (1973-2003)	870	Flag to indicate which type of EOD was coded: 0 Non-specific (N); 1 two-digit; 2 Expanded (EEOD)	Extent of Disease - Historic	9
98	2-Digit NS EOD part 1 (1973-1982)	850	This field is used in conjunction with 2-digit NS EOD part 2 to describe a very rudimentary stage. Used in 1973-1982 for some years and some sites/histologies. Use Coding system-EOD with a 0 to use these definitions. For more information see the introduction of https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
99	2-Digit NS EOD part 2 (1973-1982)	850	See the description for 2-Digit NS EOD part 1 (1973-1982).	Extent of Disease - Historic	9
100	2-Digit SS EOD part 1 (1973-1982)	850	This field is used in conjunction with 2-digit SS EOD part 2 to describe a two-digit site-specific EOD. Used in 1973-1982 for some years and some sites/histologies. Use Coding system-EOD with a 1 to use these definitions. For more information see the appropriate site-specific page of https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
101	2-Digit SS EOD part 2 (1973-1982)	850	See the description for 2-Digit SS EOD part 1 (1973-1982).	Extent of Disease - Historic	9
102	Expanded EOD(1) - CP53 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--1st Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
103	Expanded EOD(2) - CP54 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--2nd Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
104	Expanded EOD(1,2) - CP53,54 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--1st and 2nd Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
105	Expanded EOD(3) - CP55 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--3rd Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
106	Expanded EOD(4) - CP56 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--4th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
107	Expanded EOD(5) - CP57 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--5th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
108	Expanded EOD(6) - CP58 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--6th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
109	Expanded EOD(7) - CP59 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--7th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
110	Expanded EOD(8) - CP60 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--8th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
111	Expanded EOD(9) - CP61 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--9th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
112	Expanded EOD(10) - CP62 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--10th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
113	Expanded EOD(11) - CP63 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--11th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
114	Expanded EOD(12) - CP64 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--12th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
115	Expanded EOD(13) - CP65 (1973-1982)	840	NAACCR Name=EOD--Old 13 Digit--13th Digit, Item #=840. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1977.pdf .	Extent of Disease - Historic	9
116	EOD 4 - extent (1983-1987)	860	NAACCR Name=EOD--Old 4 Digit--3rd Digit, Item #=860. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1984.pdf .	Extent of Disease - Historic	9
117	EOD 4 - nodes (1983-1987)	860	NAACCR Name=EOD--Old 4 Digit--4th Digit, Item #=860. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1984.pdf .	Extent of Disease - Historic	9
118	EOD 4 - size (1983-1987)	860	NAACCR Name=EOD--Old 4 Digit--1st and 2nd Digit, Item #=860. For more information, see https://seer.cancer.gov/manuals/historic/EOD_1984.pdf .	Extent of Disease - Historic	9
119	COD to site recode		The underlying cause of death from the death certificate was grouped into a recode similar to the incidence site recode. For more information, see https://seer.cancer.gov/codrecode/1969_d04162012 . Study cutoff date has been applied, i.e. coded as alive if death occurred after study cutoff.	Cause of Death (COD) and Follow-up	10

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
120	SEER cause-specific death classification		Created for use in cause-specific survival. This variable designates that the person died of their cancer for cause-specific survival. For more information, see https://seer.cancer.gov/causespecific .	Cause of Death (COD) and Follow-up	10
121	SEER other cause of death classification		Created for use in left-truncated life table session. This variable designates that the person died of causes other than their cancer. For more information, see https://seer.cancer.gov/causespecific .	Cause of Death (COD) and Follow-up	10
122	COD to site rec KM		This is a recode based on underlying cause of death to designate cause of death into groups similar to the incidence site recode with KS and mesothelioma. For more information, see https://seer.cancer.gov/codrecode/1969_d04162012 . Study cutoff date has been applied, i.e. coded as alive if death occurred after study cutoff.	Cause of Death (COD) and Follow-up	10
123	Survival time recode (total # of months)		The Survival Time Recode is calculated using the date of diagnosis and one of the following: date of death, date last known to be alive, or follow-up cutoff date used for this file. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=77	Cause of Death (COD) and Follow-up	10
124	Survival time recode (month)		The Survival Time Recode is calculated using the date of diagnosis and one of the following: date of death, date last known to be alive, or follow-up cutoff date used for this file. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=77	Cause of Death (COD) and Follow-up	10
125	Survival time recode (year)		The Survival Time Recode is calculated using the date of diagnosis and one of the following: date of death, date last known to be alive, or follow-up cutoff date used for this file. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=77	Cause of Death (COD) and Follow-up	10
126	Vital status recode (study cutoff used)		Any patient that dies after the follow-up cut-off date is recoded to alive as of the cut-off date. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf#page=78	Cause of Death (COD) and Follow-up	10
127	Type of follow-up expected	2180	NAACCR Name=SEER Type of Follow-Up, Item #=2180	Cause of Death (COD) and Follow-up	10
128	Sequence number	380	NAACCR Name=Sequence Number--Central, Item #=380	Multiple Primary Fields	11
129	Number of primaries		Based on the total number of tumors in SEER, not necessarily this database/file. This value is the same across all tumors for a person.	Multiple Primary Fields	11
130	First malignant primary indicator		Based on all the tumors in SEER. Tumors not reported to SEER are assumed malignant.	Multiple Primary Fields	11
131	Primary by international rules		Created using IARC multiple primary rules. Did not include benign tumors or non-bladder in situ tumors in algorithm. No tumor information was modified on any records. https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf	Multiple Primary Fields	11

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
132	Record number	2190	NAACCR Name=SEER Record Number, Item #=2190. Sequentially numbers a person's tumors within each SEER submission. Order is based on sequence #. All federally reportable tumors (sequence # < 60) are prior to all state/registry reportable tumors (sequence # 60+) regardless of diagnosis date.	Multiple Primary Fields	11
133	Record number recode		Sequentially numbers a person's tumors within each SEER submission. Order is based on date of diagnosis and then sequence #.	Multiple Primary Fields	11
134	Behavior code ICD-O-2 (1973+)		Converted from ICD-O-3 for 2001+ and coded directly for 1973-2000.	Site and Morphology - Historic (ICD-O-1 and ICD-O-2)	12
135	Histology (92-00) ICD-O-2	420	NAACCR Item #=420	Site and Morphology - Historic (ICD-O-1 and ICD-O-2)	12
136	Recode ICD-O-2 to 9		Primary site/type recoded into ICD-9. An underscore in the right-most position of an unlabeled value represents a blank. i.e., C00_ C009 is all values starting with C00. All tumors not originally coded in ICD-O-2 were first converted from ICD-O-1 or ICD-O-3 and then converted to ICD-9.	Site and Morphology - Historic (ICD-O-1 and ICD-O-2)	12
137	Recode ICD-O-2 to 10		Primary site/type recoded into ICD-10. An underscore in the right-most position of an unlabeled value represents a blank. i.e., C00_ C009 is all values starting with C00. All tumors not originally coded in ICD-O-2 were first converted from ICD-O-1 or ICD-O-3 and then converted to ICD-10.	Site and Morphology - Historic (ICD-O-1 and ICD-O-2)	12
138	ICCC site recode ICD-O-2		A site/histology recode based on the International Classification of Childhood Cancer (ICCC) is mainly used to analyze data on children. The recode was applied to all cases no matter the age in order that age comparisons can be made with these groupings. Based on ICD-O-2. Note that cases diagnosed before 1992 were not coded under ICD-O-2 and were converted to ICD-O-2 and may not have the specificity of cases after 1991 that were coded directly under ICD-O-2 or 3. For more information on this site recode, see https://seer.cancer.gov/iccc .	Site and Morphology - Historic (ICD-O-1 and ICD-O-2)	12
139	SEER modified ICDC site recode ICD-O-2		A site/histology recode based on the International Classification of Childhood Cancer (ICCC) with slight modifications is mainly used to analyze data on children. The recode was applied to all cases no matter the age in order that age comparisons can be made with these groupings. Based on ICD-O-2. Note that cases diagnosed before 1992 were not coded under ICD-O-2 and were converted to ICD-O-2 and may not have the specificity of cases after 1991 that were coded directly under ICD-O-2 or 3. For more information on this site recode, see https://seer.cancer.gov/iccc .	Site and Morphology - Historic (ICD-O-1 and ICD-O-2)	12
140	Age recode with single ages and 85+		Patient's age at diagnosis with all ages over 85 grouped together. In rate and prevalence sessions, this field can be selected as the population age variable (on the data tab), in which case it will be in the Age at Diagnosis category (1)	Race and Age (case data only) (or Age at Diagnosis)	13 (or 1)
141	Race recode (W, B, AI, API)		Caution should be exercised when using this variable. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity .	Race and Age (case data only)	13
142	Origin recode NHIA (Hispanic, Non-Hisp)		Caution should be exercised when using this variable. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity .	Race and Age (case data only)	13

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
143	Age at diagnosis	230	Age at Diagnosis is the patient's age at diagnosis of this cancer and is a three digit field based on single year of age. This can not be used in a rate or prevalence session to link to populations. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Race and Age (case data only)	13
144	Race/ethnicity		Recode which gives priority to non-white races for persons of mixed races. Note that not all codes were in effect for all years. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity .	Race and Age (case data only)	13
145	Spanish surname or origin	190	NAACCR Name=Spanish/Hispanic Origin, Item #=190. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity .	Race and Age (case data only)	13
146	NHIA Derived Hisp Origin	191	NAACCR Item #=191. For more information, see https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity .	Race and Age (case data only)	13
147	IHS Link	192	Incidence files are periodically linked with Indian Health Service (IHS) files to identify Native Americans. The race recode uses information from this field and race to determine if a person is Native American or not. See https://seer.cancer.gov/seerstat/variables/seer/race_ethnicity/	Race and Age (case data only)	13
148	Place of birth	250	Place of birth includes state/province/country of birth. Caution should be used since this variable may be biased in that the death certificate is a major source for this information. For more information, see https://seer.cancer.gov/manuals/2004Revision%201/SPM_AppendixB_r1.pdf .	Geographic Locations	14
149	Year of birth	240	NAACCR Name=Birth Date--Year, Item #=240. The SEER dates on this file do not have the corresponding date flag field included. Blank means unknown.	Dates	15
150	Month of diagnosis	390	NAACCR Name=Date of Diagnosis--Month, Item #=390. SEER dates on this file do not have the corresponding date flag field included. Blank means unknown.	Dates	15
151	Month of diagnosis recode		Estimates month of diagnosis, based on other known dates for that patient, when actual month of diagnosis is unknown.	Dates	15
152	Age-site edit override	1990	Flag is set to '1' if there is an unusual combination of Age/Site/Morph to indicate that the case has been reviewed and the information is okay. Code 2 indicates that case was diagnosed <i>in utero</i> and code 3 that both conditions hold.	Overrides	17
153	Histology edit override	2040	Flag is set to '1' if there is an unusual combination of histology and behavior to indicate that the case has been reviewed and the information is okay.	Overrides	17
154	Leuk-Lymph dx confirmation override	2070	Flag is set to '1' if there is an unusual combination of Leuk, Lymphoma and Diagnostic Confirmation to indicate that the case has been reviewed and the information is okay.	Overrides	17

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
155	Report source sequence override	2050	Flag is set to '1' if there is an unusual combination of Report Source and sequence number to indicate that the case has been reviewed and the information is okay.	Overrides	17
156	Seq-ill-defined site override	2060	Flag is set to '1' if there is an unusual combination of Ill-defined Site and Sequence Number to indicate that the case has been reviewed and the information is okay.	Overrides	17
157	Sequence number-dx conf override	2000	Flag is set to '1' if there is an unusual combination of SeqNo/DxConf to indicate that the case has been reviewed and the information is okay.	Overrides	17
158	Site-type edit override	2030	Flag is set to '1' if there is an unusual combination of Site/Type to indicate that the case has been reviewed and the information is okay.	Overrides	17
159	Site-type-lat-seq override	2010	Flag is set to '1' if there is an unusual combination of Site/Lat/SeqNo to indicate that the case has been reviewed and the information is okay.	Overrides	17
160	Surgery-diagnostic conf override	2020	Flag is set to '1' if there is an unusual combination of Surg/DxConf to indicate that the case has been reviewed and the information is okay.	Overrides	17
161	Site-behavior override	2071	Flag is set to '1' if there is an unusual combination of Site/Behavior to indicate that the case has been reviewed and the information is okay.	Overrides	17
162	Site-EOD-dx date override	2072	Flag is set to '1' if there is an unusual combination of Site/EOD/DX Dt to indicate that the case has been reviewed and the information is okay.	Overrides	17
163	Site-laterality-EOD override	2073	Flag is set to '1' if there is an unusual combination of Site/Lat/EOD to indicate that the case has been reviewed and the information is okay.	Overrides	17
164	Site-laterality-morph override	2074	Flag is set to '1' if there is an unusual combination of Site/Lat/Morph to indicate that the case has been reviewed and the information is okay.	Overrides	17
165	SS seq # - mal+ins (most detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (most detail). Based on all the tumors in SEER.	Site Specific Sequence Numbers	18
166	SS seq # 1975+ - mal+ins (most detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (most detail). Based on tumors diagnosed 1975+.	Site Specific Sequence Numbers	18
167	SS seq # 1992+ - mal+ins (most detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (most detail). Based on tumors diagnosed 1992+.	Site Specific Sequence Numbers	18
168	Site - mal+ins (most detail)		Should be used in conjunction with and only with the variables SS seq # - mal+ins (most detail), SS seq # 1975+ - mal+ins (most detail), or SS seq # 1992+ - mal+ins (most detail). Groupings should not be created.	Site Specific Sequence Numbers	18
169	SS seq # - mal (most detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (most detail). Based on all the tumors in SEER.	Site Specific Sequence Numbers	18

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
170	SS seq # 1975+ - mal (most detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (most detail). Based on tumors diagnosed 1975+.	Site Specific Sequence Numbers	18
171	SS seq # 1992+ - mal (most detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (most detail). Based on tumors diagnosed 1992+.	Site Specific Sequence Numbers	18
172	Site - malignant (most detail)		Should be used in conjunction with and only with the variables SS seq # - mal (most detail), SS seq # 1975+ - mal (most detail), or SS seq # 1992+ - mal (most detail). Groupings should not be created.	Site Specific Sequence Numbers	18
173	SS seq # - mal+ins (mid detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (mid detail). Based on all the tumors in SEER.	Site Specific Sequence Numbers	18
174	SS seq # 1975+ - mal+ins (mid detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (mid detail). Based on tumors diagnosed 1975+.	Site Specific Sequence Numbers	18
175	SS seq # 1992+ - mal+ins (mid detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (mid detail). Based on tumors diagnosed 1992+.	Site Specific Sequence Numbers	18
176	Site - mal+ins (mid detail)		Should be used in conjunction with and only with the variables SS seq # - mal+ins (mid detail), SS seq # 1975+ - mal+ins (mid detail), or SS seq # 1992+ - mal+ins (mid detail). Groupings should not be created.	Site Specific Sequence Numbers	18
177	SS seq # - mal (mid detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (mid detail). Based on all the tumors in SEER.	Site Specific Sequence Numbers	18
178	SS seq # 1975+ - mal (mid detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (mid detail). Based on tumors diagnosed 1975+.	Site Specific Sequence Numbers	18
179	SS seq # 1992+ - mal (mid detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (mid detail). Based on tumors diagnosed 1992+.	Site Specific Sequence Numbers	18
180	Site - malignant (mid detail)		Should be used in conjunction with and only with the variables SS seq # - mal (mid detail), SS seq # 1975+ - mal (mid detail), or SS seq # 1992+ - mal (mid detail). Groupings should not be created.	Site Specific Sequence Numbers	18
181	SS seq # - mal+ins (least detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (least detail). Based on all the tumors in SEER.	Site Specific Sequence Numbers	18
182	SS seq # 1975+ - mal+ins (least detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (least detail). Based on tumors diagnosed 1975+.	Site Specific Sequence Numbers	18
183	SS seq # 1992+ - mal+ins (least detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - mal+ins (least detail). Based on tumors diagnosed 1992+.	Site Specific Sequence Numbers	18

Dictionary of SEER*Stat Variables
November 2011 Submission (released April 2012)

Field number	Name	NAACCR Item #	Description	Category name	Category number
184	Site - mal+ins (least detail)		Should be used in conjunction with and only with the variables SS seq # - mal+ins (least detail), SS seq # 1975+ - mal+ins (least detail), or SS seq # 1992+ - mal+ins (least detail). Groupings should not be created.	Site Specific Sequence Numbers	18
185	SS seq # - mal (least detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (least detail). Based on all the tumors in SEER.	Site Specific Sequence Numbers	18
186	SS seq # 1975+ - mal (least detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (least detail). Based on tumors diagnosed 1975+.	Site Specific Sequence Numbers	18
187	SS seq # 1992+ - mal (least detail)		Site specific sequence number of the tumor associated with the site classification scheme in the variable Site - malignant (least detail). Based on tumors diagnosed 1992+.	Site Specific Sequence Numbers	18
188	Site - malignant (least detail)		Should be used in conjunction with and only with the variables SS seq # - mal (least detail), SS seq # 1975+ - mal (least detail), or SS seq # 1992+ - mal (least detail). Groupings should not be created.	Site Specific Sequence Numbers	18
189	Patient ID	20	This field used in conjunction with SEER registry to uniquely identify a person. One person can have multiple primaries but has the same Patient ID. See the sequence number for more information about the primary. This is a dummy number and is not the number used by the registry to identify the patient. The same number is not used across all submissions for each patient.	Other	19
190	Type of Reporting Source	500	Contains information on where the information from this case came. Is used in survival analysis to eliminate cases which are found at autopsy or the only information is from a death certificate. See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Other	19
191	Marital status at diagnosis	150	See ASCII text file description: https://seer.cancer.gov/data-software/documentation/seerstat/nov2011/TextData.FileDescription.pdf .	Other	19