

# SEER Inquiry System - Report

Produced: 05/06/2024 2:44 PM

## Question 20000513

### References:

#1: SEER Program Code Man, 3rd Ed, 96, 101

#2: 2004 SEER Manual, 87, 92

### Question:

Multiple Primaries/Histology (Pre-2007)--Bladder: What code is used to represent the histology and how many primaries should be coded for a TURB specimen that demonstrates carcinoma in situ, Grade I to II papillary transitional cell carcinoma, and high grade transitional cell carcinoma? See discussion.

### Discussion:

Pathology report:

- A. Biopsy, bladder neck, @ 6:00: Carcinoma in situ
- B. Biopsy, Bladder wall, lateral, left:
  - 1. Papillary carcinoma (Grade I-II)
  - 2. Loose fragments of high-grade transitional carcinoma
- C. Biopsy, Bladder neck @ 5:00: Carcinoma in situ
- D. Biopsy, Bladder neck @ 7:00: Cystitis Glandularis
- E. Biospsy, Bladder wall, posterior: Papillary carcinoma (Grade I)

### Answer:

For tumors diagnosed prior to 2007:

Code this case as one primary and code the Histology and Grade, Differentiation fields to 8130/34 [papillary transitional cell carcinoma, high grade].

For tumors diagnosed 2007 or later, refer to the MP/H rules. If there are still questions about how this type of tumor should be coded, submit a new question to SINQ and include the difficulties you are encountering in applying the MP/H rules.

### Cancer Site Category:

Bladder

### Data Item Category:

Histology

### Other Category:

Multiple primaries

### Year:

2000