

SEER Inquiry System - Report

Produced: 05/06/2024 4:40 PM

Question 20210051

References:

#1: CAP Protocol, Ampulla of Vater. AmpullaVater 4.1.0.0, Feb 2020

#2: ICD-O-3, 49

Question:

Primary site/Biliary tract--Ampulla of Vater: What is the correct primary site code for intra-ampullary and periampullary adenocarcinoma, C241 (8144/3) or C249? See Discussion.

Discussion:

Ampulla, biopsy: High grade dysplasia with focal intramucosal carcinoma in a background of ulceration with acute and chronic inflammation.

Surgery pathology: Head of pancreas, duodenum, and distal stomach, pancreaticoduodenectomy-Ampulla, Adenocarcinoma, intestinal type, intra-ampullary and peri-ampullary (mixed type). Grade moderately differentiated, 1.5cm. Tumor invades into duodenal submucosa.

Lymphovascular Invasion: Foci suspicious for lymphovascular invasion identified. Perineural Invasion: Present.

Synoptic report: Tumor Site Intra-ampullary and peri-ampullary (mixed type). Histologic Type Adenocarcinoma, intestinal type.

There is not enough information regarding site in radiology reports or operative report. CT-A/P/C:The patient's known ampullary mass is not well visualized on this exam. No significant intrahepatic or extrahepatic biliary ductal dilation is identified. The pancreatic duct is normal caliber.

Answer:

Assign C241. Ampulla (C241) includes both periampullary and intra-ampullary.

Cancer Site Category:

N/A

Data Item Category:

Primary site

Other Category:

N/A

Year:

2021